

thanks

L'exemplaire filmé fut reproduit grâce à la
générosité de:

Dana Porter Arts Library
University of Waterloo

quality
ibility
he

Les images suivantes ont été reproduites avec le
plus grand soin, compte tenu de la condition et
de la netteté de l'exemplaire filmé, et en
conformité avec les conditions du contrat de
filmage.

e filmed
g on
impres-
. Ali
on the
pres-
printed

Les exemplaires originaux dont la couverture en
papier est imprimée sont filmés en commençant
par le premier plat et en terminant soit par la
dernière page qui comporte une empreinte
d'impression ou d'illustration, soit par le second
plat, selon le cas. Tous les autres exemplaires
originaux sont filmés en commençant par la
première page qui comporte une empreinte
d'impression ou d'illustration et en terminant par
la dernière page qui comporte une telle
empreinte.

he
CON-
ND"),

Un des symboles suivants apparaîtra sur la
dernière image de chaque microfiche, selon le
cas: le symbole → signifie "A SUIVRE", le
symbole ▽ signifie "FIN".

at
e to be
ed
left to
as
te the

Les cartes, planches, tableaux, etc., peuvent être
filmés à des taux de réduction différents.
Lorsque le document est trop grand pour être
reproduit en un seul cliché, il est filmé à partir
de l'angle supérieur gauche, de gauche à droite,
et de haut en bas, en prenant le nombre
d'images nécessaire. Les diagrammes suivants
illustrent la méthode.

	3
--	---

1
2
3

1	2	3
4	5	6

EL

ELEMENTARY ALGEBRA FOR SCHOOLS

WITH ANSWERS

BY

H. S. HALL, M.A.

FORMERLY SCHOLAR OF CHRIST'S COLLEGE, CAMBRIDGE
MASTER OF THE MILITARY SIDE, CLIFTON COLLEGE

AND

S. R. KNIGHT, B.A., M.B., CL.B.

FORMERLY SCHOLAR OF TRINITY COLLEGE, CAMBRIDGE
SOMETIME ASSISTANT MASTER AT MARLBOROUGH COLLEGE

SEVENTH EDITION
REVISED AND ENLARGED

06505

Halifax, N.S.

A. AND W. MACKINLAY

London

MACMILLAN AND CO., LIMITED

NEW YORK: THE MACMILLAN COMPANY

1900

Property of the Library
University of Waterloo

FROM

In setting
of elements
should d
symbols,
stood. T
thought t
succession
illustrate
adopted a
the comm
upon the
of frequen

As re
determined
to introdu
subject an
as easy e
opinion th
resolution
operations
simple exp
acquainted
encounterin
at the sam

EXTRACT
FROM THE PREFACE TO THE FIRST EDITION.

IN setting before a beginner the real and perplexing difficulties of elementary Algebra, there is some fear lest first lessons should degenerate into a mere mechanical manipulation of symbols, uninteresting and uninformative, because little understood. This well-known danger led us to devote special thought to the question of order; to consider, in short, what succession of the various parts of the subject would best illustrate its bearings at an early stage; and we have finally adopted an arrangement, which—if it varies somewhat from the common use of elementary text-books—is at least based upon the experience of many years, and embodies the result of frequent consultation with our colleagues and other teachers.

As regards the earlier chapters, our order has been determined mainly by two considerations: first, a desire to introduce as early as possible the practical side of the subject and some of its most interesting applications, such as easy equations and problems; and secondly, the strong opinion that all reference to compound expressions and their resolution into factors should be postponed until the usual operations of Algebra have been exemplified in the case of simple expressions. By this course the beginner soon becomes acquainted with the ordinary algebraical processes without encountering too many of their difficulties; and he is learning at the same time something of the more attractive parts of

the subject. Again, by postponing Resolution into Factors until the student has acquired some freedom and readiness in the use of symbols, we are enabled to treat this important section, and all the processes to which it gives rise, more adequately than is possible where factors are introduced and disposed of in one short early chapter.

We had originally intended to arrange the chapters in an order that might be followed without deviation from beginning to end; but in the course of the work this was found impossible without very extensive subdivision in some parts, and needless repetition in others. We have therefore marked with an asterisk all articles and examples which may conveniently be omitted by a student who is reading the subject for the first time, and we have occasionally added a note suggesting the most suitable place for a section which may have to be postponed.

June 1885.

* * * * *

The distinctive features of the Seventh Edition are:—

- (1) The definitions of *dimension*, *degree*, *homogeneous expression* are transferred from Art. 10 to Art. 24.
- (2) Greater prominence has been given to *the fundamental Laws of Algebra* (see Arts. 22, 29-32, 46-48). With this object parts of the chapters on Multiplication and Division have been re-written.
- (3) A short section on the use of *Detached Coefficients* has been given on page 37.
- (4) A fuller treatment of the *Remainder Theorem* and its applications will be found on pages 236, 237.

(5)

As
in many
without
for add
the sam
little a
of misc
chapter
editions
will all
inconve
old form
editions
complet
text-bo
all who
the Bin
I re
Knight
which v
and rev

Jan. 1

- (5) Five new sets of *Miscellaneous Examples* have been added at convenient intervals, beginning with one on page 32, which replaces Examples IV.c. With the exception of this change and a combination of Examples XI.b and XI.c, which now appear as one exercise, all the original examples will be found under their old numbers and with the answers unaltered, even where the examples themselves have been improved.

As the book has been entirely reprinted, improvement in many typographical details has been secured ; in particular, without any appearance of crowding, space has been found for additions to some of the more important exercises. At the same time it will be found that the pagination is very little altered, the only difference being that the introduction of miscellaneous examples has placed the beginning of some chapters a few pages further on in the book than in previous editions. It is believed that the few alterations in the text will all be justified by usage, and that they will cause no inconvenience to those who are familiar with the book in its old form. The very favourable reception accorded to previous editions encourages the hope that, in its present more complete form, the work will be found suitable as a first text-book for every class of student, and amply sufficient for all whose study of Algebra does not extend much beyond the Binomial Theorem.

I regret that by the early and lamented death of Dr. Knight I have been deprived of the advice and assistance which would have been invaluable in the final arrangement and revision of this our first joint work.

H. S. HALL.

Jan. 1897.

CHAP

I.

II.

III.

IV.

V.

VI.

VII.

VIII.

IX.

X.

XI.

XII.

XIII.

XIV.

XV.

XVI.

XVII.

XVIII.

XIX.

XX.

XXI.

CONTENTS.

CHAP.	PAGE
I. DEFINITIONS, SUBSTITUTIONS	1
II. NEGATIVE QUANTITIES. ADDITION OF LIKE TERMS	9
III. SIMPLE BRACKETS. ADDITION	13
IV. SUBTRACTION	20
MISCELLANEOUS EXAMPLES I.	23
V. MULTIPLICATION	26
VI. DIVISION	38
VII. REMOVAL AND INSERTION OF BRACKETS	46
VIII. SIMPLE EQUATIONS	52
IX. SYMBOLICAL EXPRESSION	61
X. PROBLEMS LEADING TO SIMPLE EQUATIONS	69
XI. HIGHEST COMMON FACTOR, LOWEST COMMON MUL- TIPLE OF SIMPLE EXPRESSIONS	74
XII. ELEMENTARY FRACTIONS	76
MISCELLANEOUS EXAMPLES II.	80
XIII. SIMULTANEOUS EQUATIONS	83
XIV. PROBLEMS LEADING TO SIMULTANEOUS EQUATIONS.	93
XV. INVOLUTION	98
XVI. EVOLUTION	102
XVII. RESOLUTION INTO FACTORS	112
MISCELLANEOUS EXAMPLES III.	126
XVIII. HIGHEST COMMON FACTOR	128
XIX. FRACTIONS	136
XX. LOWEST COMMON MULTIPLE	144
XXI. ADDITION AND SUBTRACTION OF FRACTIONS	148

CHAP.	PAGE
XXII. MISCELLANEOUS FRACTIONS	160
MISCELLANEOUS EXAMPLES IV.	172
XXIII. HARDER EQUATIONS	178
XXIV. HARDER PROBLEMS	186
XXV. QUADRATIC EQUATIONS	192
XXVI. SIMULTANEOUS QUADRATIC EQUATIONS	202
XXVII. PROBLEMS LEADING TO QUADRATIC EQUATIONS	209
XXVIII. HARDER FRACTIONS	215
XXIX. MISCELLANEOUS THEOREMS AND EXAMPLES	222
XXX. THE THEORY OF INDICES	240
XXXI. ELEMENTARY SURDS	253
XXXII. RATIO, PROPORTION, AND VARIATION	270
XXXIII. ARITHMETICAL PROGRESSION	287
XXXIV. GEOMETRICAL PROGRESSION	294
XXXV. HARMONICAL PROGRESSION	301
MISCELLANEOUS EXAMPLES V.	307
XXXVI. THEORY OF QUADRATIC EQUATIONS	311
XXXVII. PERMUTATIONS AND COMBINATIONS	319
XXXVIII. BINOMIAL THEOREM	329
XXXIX. LOGARITHMS	340
XL. SCALES OF NOTATION	349
XLI. EXPONENTIAL AND LOGARITHMIC SERIES	355
XLII. MISCELLANEOUS EQUATIONS	361
XLIII. INTEREST AND ANNUITIES	367
MISCELLANEOUS EXAMPLES VI.	371
ANSWERS	393

1. AL
greater g
processes
value, alg
have any

The sy
alphabet;
values a sy
piece of w
we say "le
always, bu
Moreover,
them any
such opera

We begi
symbols +
Arithmetic
the algebra

2. An
it may con
each other
an expressi

Note. W
E. A.

160
172
178
186
192
202
209
215
222
240
253
270
287
294
301
307
311
319
329
340
349
355
361
367
371
393

ALGEBRA.

CHAPTER I.

DEFINITIONS. SUBSTITUTIONS.

1. ALGEBRA treats of quantities as in Arithmetic, but with greater generality; for while the quantities used in arithmetical processes are denoted by *figures* which have one single definite value, algebraical quantities are denoted by *symbols* which may have any value we choose to assign to them.

The symbols employed are letters, usually those of our own alphabet; and, though there is no restriction as to the numerical values a symbol may represent, it is understood that in the same piece of work it keeps the same value throughout. Thus, when we say "let $a=1$," we do not mean that a must have the value 1 always, but only in the particular example we are considering. Moreover, we may operate with symbols without assigning to them any particular numerical value at all; indeed it is with such operations that Algebra is chiefly concerned.

We begin with the definitions of Algebra, premising that the symbols $+$, $-$, $\times$, $\div$, $=$, will have the same meanings as in Arithmetic. Also, for the present it will be assumed that all the algebraical symbols employed denote integral numbers.

2. An **algebraical expression** is a collection of symbols; it may consist of one or more **terms**, which are separated from each other by the signs $+$ and $-$. Thus $7a+5b-3c-x+2y$ is an expression consisting of five terms.

Note. When no sign precedes a term the sign $+$ is understood.

E. A.

A

E

3. **Expressions** are either **simple** or **compound**. A *simple expression* consists of *one* term, as $5a$. A *compound expression* consists of *two or more* terms. Compound expressions may be further distinguished. Thus an expression of *two* terms, as $3a - 2b$, is called a **binomial** expression; one of *three* terms, as $2a - 3b + c$, a **trinomial**; one of *more than three* terms a **multinomial**. Simple expressions are also spoken of as **monomials**.

4. When two or more quantities are multiplied together the result is called the **product**. One important difference between the notation of Arithmetic and Algebra should be here remarked. In Arithmetic the product of 2 and 3 is written 2×3 , whereas in Algebra the product of a and b may be written in any of the forms $a \times b$, $a.b$, or ab . The form ab is the most usual. Thus, if $a=2$, $b=3$, the product $ab = a \times b = 2 \times 3 = 6$; but in Arithmetic 23 means "twenty-three," or $2 \times 10 + 3$.

5. Each of the quantities multiplied together to form a product is called a **factor** of the product. Thus 5, a , b , are the factors of the product $5ab$.

6. When one of the factors of an expression is a numerical quantity, it is called the **coefficient** of the remaining factors. Thus, in the expression $5ab$, 5 is the coefficient. But the word coefficient is also used in a wider sense, and it is sometimes convenient to consider any factor, or factors, of a product as the coefficient of the remaining factors. Thus, in the product $6abc$, $6a$ may be appropriately called the coefficient of bc . A coefficient which is not merely numerical is sometimes called a **literal coefficient**.

Note. When the coefficient is unity it is usually omitted. Thus we do not write $1a$, but simply a .

7. If a quantity be multiplied by itself any number of times, the product is called a **power** of that quantity, and is expressed by writing the number of factors to the right of the quantity and above it. Thus

$a \times a$ is called the *second power* of a , and is written a^2 ;
 $a \times a \times a$*third power* of a , ... a^3 ;
 and so on.

The number which expresses the power of any quantity is called its **index** or **exponent**. Thus 2, 5, 7 are respectively the indices of a^2 , a^5 , a^7 .

Note. a^2 is usually read " a squared"; a^3 is read " a cubed"; a^4 is read " a to the fourth"; and so on.

When the index is unity it is omitted. Thus we do not write a^1 , but simply a . Thus a , $1a$, a^1 , $1a^1$ all have the same meaning.

8. The coefficient

Example

By $3a$

By a^3

qua

Thus,

Example

Here

when

Example

Here

Note. T

9. In factors of 3×4 mean in each case

In a sim

and it is product of

In like manner of the two have there acb , bac , abc product of what order however, to

Fractions kept in the

Example.

Here

A *simple expression* may be terms, as terms, as a **multi-nomials**.

gether the e between remarked. , whereas in any of ost usual. ; but in

rm a pro- , are the

numerical g factors. the word ometimes roduct as e product of *bc*. A s called a

ed. Thus

of times, expressed quantity

a^2 ;
 a^3 ;

antity is pectively

cubed";

ot write meaning.

8. The beginner must be careful to distinguish between *coefficient* and *index*.

Example 1. What is the difference in meaning between $3a$ and a^3 ?

By $3a$ we mean the product of the quantities 3 and a .

By a^3 we mean the third power of a ; that is, the product of the quantities a, a, a .

Thus, if $a=4$,

$$3a = 3 \times a = 3 \times 4 = 12;$$

$$a^3 = a \times a \times a = 4 \times 4 \times 4 = 64.$$

Example 2. If $b=5$, distinguish between $4b^2$ and $2b^4$.

Here $4b^2 = 4 \times b \times b = 4 \times 5 \times 5 = 100$;

whereas $2b^4 = 2 \times b \times b \times b \times b = 2 \times 5 \times 5 \times 5 \times 5 = 1250$.

Example 3. If $a=4$, $x=1$, find the value of $5x^a$.

Here $5x^a = 5 \times x \times x \times x \times x = 5 \times 1 \times 1 \times 1 \times 1 = 5$.

Note. The beginner should observe that every power of 1 is 1.

9. In arithmetical multiplication the order in which the factors of a product are written is immaterial. For instance 3×4 means 4 sets of 3 units, and 4×3 means 3 sets of 4 units; in each case we have 12 units in all. Thus

$$3 \times 4 = 4 \times 3.$$

In a similar way,

$$3 \times 4 \times 5 = 4 \times 3 \times 5 = 4 \times 5 \times 3;$$

and it is easy to see that the same principle holds for the product of any number of arithmetical quantities.

In like manner in Algebra ab and ba each denote the product of the two quantities represented by the letters a and b , and have therefore the same value. Again, the expressions abc , acb , bac , bca , cab , cba have the same value, each denoting the product of the three quantities a, b, c . It is immaterial in what order the factors of a product are written; it is usual, however, to arrange them in alphabetical order.

Fractional coefficients which are greater than unity are usually kept in the form of improper fractions.

Example. If $a=6$, $x=7$, $z=5$, find the value of $\frac{13}{10}axz$.

Here $\frac{13}{10}axz = \frac{13}{10} \times 6 \times 7 \times 5 = 273$.

EXAMPLES I. a.

If $a=7$, $b=2$, $c=1$, $x=5$, $y=3$, find the value of

- | | | | | |
|--------------|--------------|--------------|--------------|--------------|
| 1. $14x$. | 2. x^3 . | 3. $3ax$. | 4. a^3 . | 5. $5by$. |
| 6. b^5 . | 7. $3b^2$. | 8. $2xa$. | 9. $6c^4$. | 10. $4y^2$. |
| 11. $7c^5$. | 12. $9b^4$. | 13. $8bcy$. | 14. $7y^3$. | 15. $8x^2$. |

If $a=8$, $b=5$, $c=4$, $x=1$, $y=3$, find the value of

- | | | | | |
|-------------|--------------|--------------|-------------|--------------|
| 16. $9xy$. | 17. $8b^3$. | 18. $3x^5$. | 19. x^2 . | 20. $7y^4$. |
| 21. c^2 . | 22. b^y . | 23. y^c . | 24. x^b . | 25. y^5 . |
| 26. a^y . | 27. b^z . | 28. a^c . | 29. c^y . | 30. $6bxy$. |

If $a=5$, $b=1$, $c=6$, $x=4$, find the value of

- | | | | | |
|------------------------|-------------------------|------------------------|------------------------|------------------------|
| 31. $\frac{3}{8}x^3$. | 32. $\frac{1}{10}ax$. | 33. 3^z . | 34. 2^c . | 35. 8^b . |
| 36. 7^z . | 37. $\frac{7}{18}acx$. | 38. $\frac{1}{8}bcx$. | 39. $\frac{2}{9}c^3$. | 40. $\frac{x^5}{64}$. |

10. When several different quantities are multiplied together a notation similar to that of Art. 7 is adopted. Thus $aabbbbcd$ is written $a^2b^4cd^3$. And conversely $7a^3cd^2$ has the same meaning as $7 \times a \times a \times a \times c \times d \times d$.

Example 1. If $x=5$, $y=3$, find the value of $4x^2y^3$.

$$\begin{aligned} 4x^2y^3 &= 4 \times 5^2 \times 3^3 \\ &= 4 \times 25 \times 27 \\ &= 2700. \end{aligned}$$

Example 2. If $a=4$, $b=9$, $x=6$, find the value of $\frac{8bx^2}{27a^3}$.

$$\begin{aligned} \frac{8bx^2}{27a^3} &= \frac{8 \times 9 \times 6^2}{27 \times 4^3} = \frac{8 \times 9 \times 36}{27 \times 64} \\ &= \frac{3}{2} = 1\frac{1}{2}. \end{aligned}$$

11. If one factor of a product is equal to 0, the whole product must be equal to 0, *whatever values the other factors may have*. A factor 0 is usually called a **zero factor**.

For instance, if $x=0$ then ab^3xy^2 contains a zero factor. Therefore $ab^3xy^2=0$ when $x=0$, whatever be the values of a , b , y . Again, if $c=0$, then $c^3=0$; therefore $ab^2c^3=0$, whatever values a and b may have.

Note. Every power of 0 is 0.

EXAMPLES I. b.

If $a=7$, $b=2$, $c=0$, $x=5$, $y=3$, find the value of

- | | | | |
|------------------------|--------------------------|---------------------------|---------------------------|
| 1. $4ax^2$. | 2. a^3b . | 3. $8b^2y$. | 4. $3xy^2$. |
| 5. $\frac{3}{4}b^2x$. | 6. $\frac{5}{8}b^3y^2$. | 7. $\frac{a}{8}xy^4$. | 8. a^3c . |
| 9. a^2cy . | 10. $8x^2y$. | 11. $\frac{7}{10}ab^3x$. | 12. $\frac{1}{8}x^2y^4$. |

If $a=2$, $b=3$, $c=1$, $p=0$, $q=4$, $r=6$, find the value of

- | | | | |
|-------------------------------|----------------------------|---------------------------|----------------------------|
| 13. $\frac{3a^2r}{8b}$. | 14. $\frac{8ab^2}{9q^2}$. | 15. $\frac{6a^3c}{b^2}$. | 16. $\frac{4cr^2}{9a^3}$. |
| 17. $3a^2b^c$. | 18. $\frac{5}{8}ba^r$. | 19. $\frac{8b^q}{9a^r}$. | 20. $5a^bcr$. |
| 21. $\frac{2a^3p}{7r}$. | 22. $3a^2b$. | 23. 2^ra^b . | 24. c^bb^a . |
| 25. $\frac{5a^rb^q}{64r^a}$. | 26. $\frac{27a^q}{32}$. | 27. $\frac{64}{q^r}$. | 28. $\frac{b^r}{r^b}$. |

12. DEFINITION. The **square root** of any proposed expression is that quantity whose square, or second power, is equal to the given expression. Thus the square root of 81 is 9, because $9^2=81$.

The square root of a is denoted by $\sqrt[2]{a}$, or more simply $\sqrt{a}$.

Similarly the **cube**, **fourth**, **fifth**, etc., **root** of any expression is that quantity whose third, fourth, fifth, etc., power is equal to the given expression.

The roots are denoted by the symbols $\sqrt[3]{}$, $\sqrt[4]{}$, $\sqrt[5]{}$, etc.

Examples. $\sqrt[3]{27}=3$; because $3^3=27$.

$\sqrt[4]{32}=2$; because $2^4=32$.

The symbol $\sqrt{}$ is sometimes called the **radical sign**.

Example 1. Find the value of $5\sqrt{(6a^3b^4c)}$, when $a=3$, $b=1$, $c=8$.

$$\begin{aligned} 5\sqrt{(6a^3b^4c)} &= 5 \times \sqrt{(6 \times 3^3 \times 1^4 \times 8)} = 5 \times \sqrt{(6 \times 27 \times 8)} \\ &= 5 \times \sqrt{1296} = 5 \times 36 = 180. \end{aligned}$$

Example 2. Find the value of $\sqrt[3]{\left(\frac{ab^4}{8x^3}\right)}$, when $a=9$, $b=3$, $x=5$.

$$\begin{aligned} \sqrt[3]{\left(\frac{ab^4}{8x^3}\right)} &= \sqrt[3]{\left(\frac{9 \times 3^4}{8 \times 5^3}\right)} = \sqrt[3]{\left(\frac{9 \times 81}{8 \times 125}\right)} \\ &= \sqrt[3]{\left(\frac{9 \times 9 \times 9}{1000}\right)} = \frac{9}{10}. \end{aligned}$$

EXAMPLES I. c.

If $a=8$, $c=0$, $k=9$, $x=4$, $y=1$, find the value of

- | | | |
|--|--|---|
| 1. $\sqrt{(2a)}$. | 2. $\sqrt{(kx)}$. | 3. $\sqrt{(2ax)}$. |
| 4. $\sqrt{(2ak^2)}$. | 5. $\sqrt[3]{(3k)}$. | 6. $\sqrt[3]{(ax^2)}$. |
| 7. $\sqrt[3]{(8x^2y^3)}$. | 8. $\sqrt[3]{(cy^5)}$. | 9. $2x\sqrt{(2ay)}$. |
| 10. $5y\sqrt{(4kx)}$. | 11. $3c\sqrt{(kx)}$. | 12. $2xy\sqrt{(4y^5)}$. |
| 13. $\sqrt{\left(\frac{8x^3}{ak}\right)}$. | 14. $\sqrt{\left(\frac{25a}{2k}\right)}$. | 15. $\sqrt{\left(\frac{16x}{49y^3}\right)}$. |
| 16. $\sqrt{\left(\frac{ca^2}{16k}\right)}$. | 17. $\sqrt[3]{\left(\frac{3a}{k^2}\right)}$. | 18. $\sqrt[3]{\left(\frac{ax^3}{27y^3}\right)}$. |
| 19. $\sqrt[3]{\left(\frac{ca}{3k}\right)}$. | 20. $\sqrt[3]{\left(\frac{a^2k^2}{3x^3}\right)}$. | 21. $\sqrt{\left(\frac{kax^2}{18y^3}\right)}$. |

If $a=4$, $b=1$, $c=2$, $d=9$, $x=5$, $y=8$, find the value of

- | | | |
|--|--|--------------------------------------|
| 22. $6\sqrt{(4b^3)}$. | 23. $7\sqrt{(5dx)}$. | 24. $\sqrt{(c^5y)}$. |
| 25. $\sqrt{\left(\frac{1}{5xy^2}\right)}$. | 26. $\sqrt{\left(\frac{1}{8acd}\right)}$. | 27. $\frac{1}{\sqrt{(9a^3c^2)}}$. |
| 28. $\frac{1}{\sqrt{(5c^4x)}}$. | 29. $\sqrt[3]{\left(\frac{1}{4cx^3}\right)}$. | 30. $\frac{1}{\sqrt[3]{(8ab^2c)}}$. |
| 31. $\sqrt[3]{\left(\frac{3x^3}{d^2y}\right)}$. | 32. $\sqrt[3]{\left(\frac{5b^3}{x^4y}\right)}$. | 33. $\sqrt{d^c}$. |
| 34. $\sqrt{y^a}$. | 35. $\sqrt{b^d}$. | 36. $\sqrt{d^b}$. |

13. We now proceed to find the numerical value of expressions which contain more than one term. In these each term can be dealt with singly by the rules already given, and by combining the terms the numerical value of the whole expression is obtained.

The following example further illustrates the difference between coefficient and index.

Example. When $c=5$, find the value of $c^4 - 4c + 2c^3 - 3c^2$.

Here

$$c^4 = 5^4 = 5 \times 5 \times 5 \times 5 = 625;$$

$$4c = 4 \times 5 = 20;$$

$$2c^3 = 2 \times 5^3 = 2 \times 5 \times 5 \times 5 = 250;$$

$$3c^2 = 3 \times 5^2 = 3 \times 5 \times 5 = 75.$$

Hence the value of the expression

$$= 625 - 20 + 250 - 75 = 780.$$

14. By Art. 11 any term which contains a *zero factor* is itself zero, and may be called a *zero term*.

Example 1. If $a=2$, $b=0$, $x=5$, $y=3$, find the value of

$$5a^3 - ab^2 + 2x^2y + 3bxy$$

$$\begin{aligned}\text{The expression} &= (5 \times 2^3) - 0 + (2 \times 5^2 \times 3) + 0 \\ &= 40 + 150 = 190.\end{aligned}$$

Note. The two zero terms do not affect the result.

Example 2. Find the value of $\frac{3}{5}x^2 - a^2y + 7abx - \frac{5}{2}y^3$, when
 $a=5$, $b=0$, $x=7$, $y=1$.

$$\begin{aligned}\frac{3}{5}x^2 - a^2y + 7abx - \frac{5}{2}y^3 &= \frac{3}{5} \cdot 7^2 - 5^2 \cdot 1 + 0 - \frac{5}{2} \cdot 1^3 \\ &= 29\frac{2}{5} - 25 - 2\frac{1}{2} \\ &= 1\frac{9}{10}.\end{aligned}$$

Example 3. When $p=9$, $r=6$, $k=4$, find the value of

$$\begin{aligned}&\frac{1}{3}\sqrt[3]{\left(\frac{pr}{k^2}\right)} + \sqrt{(5p+3r+1)} - \frac{2r^2}{9k} \\ &\frac{1}{3}\sqrt[3]{\left(\frac{pr}{k^2}\right)} + \sqrt{(5p+3r+1)} - \frac{2r^2}{9k} = \frac{1}{3}\sqrt[3]{\left(\frac{54}{16}\right)} + \sqrt{(45+18+1)} - \frac{2 \times 36}{9 \times 4} \\ &= \frac{1}{3}\sqrt[3]{\frac{27}{8}} + \sqrt{(64)} - 2 \\ &= \frac{1}{3} \times \frac{3}{2} + 8 - 2 \\ &= 6\frac{1}{2}.\end{aligned}$$

15. In working examples the student should pay attention to the following hints.

1. Too much importance cannot be attached to neatness of style and arrangement. The beginner should remember that neatness is in itself conducive to accuracy.

2. The sign $=$ should never be used except to connect quantities which are equal. Beginners should be particularly careful not to employ the sign of equality in any vague and inexact sense.

3. Unless the expressions are very short the signs of equality in the steps of the work should be placed one under the other.

4. It should be clearly brought out how each step follows from the one before it; for this purpose it will sometimes be advisable to add short verbal explanations; the importance of this will be seen later.

EXAMPLES I. d.

If $a=2$, $b=3$, $c=1$, $d=0$, find the numerical value of

1. $6a+5b-8c+9d$.
2. $3a-4b+6c+5d$.
3. $5a+3c-2b+6d$.
4. $ab+bc+ca-da$.
5. $6ab-3cd+2da-5cb+2db$.
6. $abc+bcd+cda+dab$.
7. $3abc-2bcd+2cda-4dab$.
8. $2bc+3cd-4da+5ab$.
9. $3bcd+5cda-7dab+abc$.
10. $a^2+b^2+c^2+d^2$.
11. $2a^2+3b^3-4c^4$.
12. $a^4+b^4-c^4$.

If $a=1$, $b=2$, $c=3$, $d=0$, find the numerical value of

13. $a^3+b^3+c^3+d^3$.
14. $\frac{1}{2}bc^3-a^3-b^3-\frac{3}{4}ab^3c$.
15. $3abc-b^2c-6a^2$.
16. $2a^2+2b^2+2c^2+2d^2-2bc-2cd-2da-2ab$.
17. $c^3+\frac{4}{5}ad^4-3a^3+b^2d$.
18. $a^3+2b^3+2c^2+d^2+2ab+2bc+\frac{2}{7}cd$.
19. $2c^3+2a^2+2b^2-4cb+6abcd$.
20. $13a^2+\frac{11}{6}c^4+20ab-16ac-16bc$.
21. $6ab-\frac{4}{3}ac^2-2a+\frac{1}{8}b^4-3d+\frac{4}{5}c^3$.
22. $a^2-c^2+b^2-d^2+2ab-2cd$.
23. $2ab-\frac{3}{4}b^3+3ac-2c-d+\frac{4}{15}ad$.
24. $125b^4c-9d^5+3abc^2d$.

If $a=8$, $b=6$, $c=1$, $x=9$, $y=4$, find the value of

25. $\frac{5}{3}a-\frac{1}{6}b^3+\frac{7}{8}y^2$.
26. $\frac{5}{27}ax-\frac{32}{y^2}-\frac{6a}{cxy}$.
27. $\frac{3a^2b}{cxy^2}-\frac{5y}{a}$.
28. $\sqrt[3]{\left(\frac{6cy^4}{x^2}\right)}+2\sqrt{\left(\frac{3a^3}{4b^3}\right)}$.
29. $\frac{2}{5}ab^2-\sqrt{\left(\frac{x}{c^3}\right)}-bc^3y$.
30. $\frac{2}{b^2}(bxy)-\frac{1}{8}b^2+\frac{8x^2}{b^2y}$.
31. $\frac{3}{4}ac-\sqrt{\left(\frac{b^2}{9y}\right)}-\sqrt[3]{\left(\frac{by}{x^2}\right)}$.
32. $\frac{5b^2y^3}{12a^2x}-\sqrt{\left(\frac{x^4a}{b^2y^3}\right)}+\sqrt{\left(\frac{ab^3}{3x}\right)}$.

NEGA

16. In
tomed to
signs +
as $1\frac{1}{2}+7$
which the
sign - is
which no
The same
sion $7a+$
represent

17. In
greater th
were the
In Algebr
terms exc
stand alon

Hence a
quantities
expressed
irrespective

This ide
trations.

(i) Sup
total gain
loses £100

The corr

and the ne
debt, that is

CHAPTER II.

NEGATIVE QUANTITIES. ADDITION OF LIKE TERMS.

16. In his arithmetical work the student has been accustomed to deal with numerical quantities connected by the signs + and -; and in finding the value of an expression such as $1\frac{1}{2} + 7\frac{2}{3} - 3\frac{1}{3} + 6 - 4\frac{1}{3}$ he understands that the quantities to which the sign + is prefixed are *additive*, and those to which the sign - is prefixed are *subtractive*, while the first quantity, $1\frac{1}{2}$, to which no sign is prefixed, is counted among the additive terms. The same notions prevail in Algebra; thus in using the expression $7a + 3b - 4c - 2d$ we understand the symbols $7a$ and $3b$ to represent additive quantities, while $4c$ and $2d$ are subtractive.

17. In Arithmetic the sum of the additive terms is always greater than the sum of the subtractive terms; if the reverse were the case the result would have no arithmetical meaning. In Algebra, however, not only may the sum of the subtractive terms exceed that of the additive, but a subtractive term may stand alone, and yet have a meaning quite intelligible.

Hence all algebraical quantities may be divided into **positive quantities** and **negative quantities**, according as they are expressed with the sign + or the sign -; and this is quite irrespective of any actual process of addition and subtraction.

This idea may be made clearer by one or two simple illustrations.

(i) Suppose a man were to gain £100 and then lose £70, his total *gain* would be £30. But if he first gains £70 and then loses £100 the result of his trading is a *loss* of £30.

The corresponding algebraical statements would be

$$\begin{aligned}\text{£}100 - \text{£}70 &= + \text{£}30, \\ \text{£}70 - \text{£}100 &= - \text{£}30,\end{aligned}$$

and the negative quantity in the second case is interpreted as a *debt*, that is, a sum of money opposite in character to the positive

quantity, or *gain*, in the first case; in fact it may be said to possess a subtractive quality which would produce its effect on other transactions, or perhaps wholly counterbalance a sum gained.

(ii) Suppose a man starting from a given point were to walk along a straight road 100 yards forwards and then 70 yards backwards, his distance from the starting-point would be 30 yards. But if he first walks 70 yards forwards and then 100 yards backwards his distance from the starting-point would be 30 yards, but *on the opposite side of it*. As before we have

$$\begin{aligned} 100 \text{ yards} - 70 \text{ yards} &= +30 \text{ yards,} \\ 70 \text{ yards} - 100 \text{ yards} &= -30 \text{ yards.} \end{aligned}$$

In each of these cases the man's *absolute distance* from the starting point is the same; but by taking the positive and negative signs into account, we see that -30 is a distance from the starting point *equal in magnitude but opposite in direction* to the distance represented by $+30$. Thus the negative sign may here be taken as indicating a *reversal of direction*.

Many other illustrations might be chosen; but it will be sufficient here to remind the student that a subtractive quantity is always *opposite* in character to an additive quantity of equal *absolute value*. In other words *subtraction is the reverse of addition*.

Addition of Like Terms.

18. DEFINITION. When terms do not differ, or when they differ only in their numerical coefficients, they are called **like**, otherwise they are called **unlike**. Thus $3a$, $7a$; $5a^2b$, $2a^2b$; $3a^2b^2$, $-4a^2b^2$ are pairs of like terms; and $4a$, $3b$; $7a^2$, $9a^2b$ are pairs of unlike terms. In the present chapter we shall only consider the addition of like terms.

The Rules for adding like terms are

Rule I. *The sum of a number of like terms is a like term.*

Rule II. *If all the terms are positive, add the coefficients.*

Example. Find the value of $8a + 5a$.

Here we have to increase 8 like things by 5 like things of the same kind, and the aggregate is 13 of such things;

$$\text{for instance,} \quad 8 \text{ lbs.} + 5 \text{ lbs.} = 13 \text{ lbs.}$$

$$\text{Hence also,} \quad 8a + 5a = 13a.$$

$$\text{Similarly,} \quad 8a + 5a + a + 2a + 6a = 22a.$$

Rule
numeric

Exam

Her

qu

av

sa

Thur

Rule
gether se
coefficient
results, p
of the sum

Exam
17

Exam

The

.....

The
tive

We ne
terms ma
convenient

This pr

19. W
the result
Thus 1
of $11a$, $-$

Exampl

The s

Note. T
opposite si

Rule III. *If all the terms are negative, add the coefficients numerically and prefix the minus sign to the sum.*

Example. To find the sum of $-3x$, $-5x$, $-7x$, $-x$.

Here the word *sum* indicates the aggregate of 4 subtractive quantities of like character. In other words, we have to *take away successively* 3, 5, 7, 1 like things, and the result is the same as taking away $3+5+7+1$ such things in the aggregate.

Thus the sum of $-3x$, $-5x$, $-7x$, $-x$ is $-16x$.

Rule IV. *If the terms are not all of the same sign, add together separately the coefficients of all the positive terms and the coefficients of all the negative terms; the difference of these two results, preceded by the sign of the greater, will give the coefficient of the sum required.*

Example 1. The sum of $17x$ and $-8x$ is $9x$, for the difference of 17 and 8 is 9, and the greater is positive.

Example 2. To find the sum of $8a$, $-9a$, $-a$, $3a$, $4a$, $-11a$, a .

The sum of the coefficients of the positive terms is 16.

..... negative 21.

The difference of these is 5, and the sign of the greater is negative; hence the required sum is $-5a$.

We need not, however, adhere strictly to this rule, for the terms may be added or subtracted in the order we find most convenient.

This process is called **collecting terms**.

19. When quantities are connected by the signs $+$ and $-$, the resulting expression is called their **algebraical sum**.

Thus $11a - 27a + 13a = -3a$ states that the algebraical sum of $11a$, $-27a$, $13a$ is equal to $-3a$.

Example. Find the algebraical sum of $\frac{2}{3}a$, $3a$, $-\frac{1}{6}a$, $-2a$.

$$\begin{aligned}\text{The sum} &= 3\frac{2}{3}a - 2\frac{1}{6}a \\ &= 1\frac{1}{2}a \\ &= \frac{3}{2}a.\end{aligned}$$

Note. The sum of two quantities numerically equal but with opposite signs is zero. Thus the sum of $5a$ and $-5a$ is 0.

EXAMPLES II.

Find the sum of

1. $5a, 7a, 11a, a, 23a.$
2. $4x, x, 3x, 7x, 9x.$
3. $7b, 10b, 11b, 9b, 2b.$
4. $6c, 8c, 2c, 15c, 19c, 100c, c.$
5. $-3x, -5x, -11x, -7x.$
6. $-5b, -6b, -11b, -18b.$
7. $-3y, -7y, -y, -2y, -4y.$
8. $-c, -2c, -50c, -13c.$
9. $-11b, -5b, -3b, -b.$
10. $5x, -x, -3x, 2x, -x.$
11. $20y, -11y, -15y, y, -3y, 2y.$
12. $5f, -9f, -3f, 21f, -30f.$
13. $2s, -3s, s, -s, -8s, 5s.$
14. $7y, -11y, 16y, -3y, -2y.$
15. $5x, -7x, -2x, 7x, 2x, -5x.$
16. $7ab, -3ab, -5ab, 2ab, ab.$

Find the value of

17. $-9x^2 + 11x^2 + 3x^2 - 4x^2.$
18. $3a^2x - 18a^2x + a^2x.$
19. $3a^3 - 7a^3 - 8a^3 + 2a^3 - 11a^3.$
20. $4x^3 - 5x^3 - 8x^3 - 7x^3.$
21. $4a^2b^3 - a^2b^3 - 7a^2b^3 + 5a^2b^3 - a^2b^3.$
22. $-9x^4 - 4x^4 - 12x^4 + 13x^4 - 7x^4.$
23. $7abcd - 11abcd - 41abcd + 2abcd.$
24. $\frac{1}{2}x - \frac{1}{3}x + x + \frac{2}{3}x.$
25. $\frac{3}{8}a + \frac{5}{8}a - \frac{1}{2}a.$
26. $-5b + \frac{1}{4}b - \frac{3}{2}b + 2b - \frac{1}{2}b + \frac{7}{4}b.$
27. $-\frac{5}{3}x^2 - 2x^2 - \frac{2}{3}x^2 + x^2 + \frac{1}{2}x^2 + \frac{1}{6}x^2.$
28. $-ab - \frac{1}{2}ab - \frac{1}{3}ab - \frac{1}{4}ab - \frac{1}{5}ab + ab + \frac{5}{12}ab.$
29. $\frac{2}{3}x - \frac{3}{4}x + \frac{5}{6}x - 2x + \frac{1}{6}x - \frac{1}{3}x + x.$
30. $-\frac{5}{3}x^2 - \frac{2}{4}x^2 - \frac{4}{5}x^2 - \frac{1}{4}x^2 - x^2.$

20. V
together
same in
the case

Thus a
two expr
the secon
reasoning
write the

Thus i
the equiv

To illus
sents a g
immaterial
cedes the

21. Br
within th
of bracke
only with

$8 + (13 -$
added to 8
or togethe

Thus

Similar
added to a

Thus

9x.
 19c, 100c, c.
 11b, -18b.
 c, -13c.
 2x, -x.

CHAPTER III.

SIMPLE BRACKETS. ADDITION.

20. WHEN a number of arithmetical quantities are connected together by the signs + and -, the value of the result is the same in whatever order the terms are taken. This also holds in the case of algebraical quantities.

Thus $a-b+c$ is equivalent to $a+c-b$, for in the first of the two expressions b is taken from a , and c added to the result; in the second c is added to a , and b taken from the result. Similar reasoning applies to all algebraical expressions. Hence we may write the terms of an expression in any order we please.

Thus it appears that the expression $a-b$ may be written in the equivalent form $-b+a$.

To illustrate this we may suppose, as in Art. 17, that a represents a gain of a pounds, and $-b$ a loss of b pounds: it is clearly immaterial whether the gain precedes the loss or the loss precedes the gain.

21. Brackets () are used to indicate that the terms enclosed within them are to be considered as one quantity. The full use of brackets will be considered in Chap. VII.; here we shall deal only with the simpler cases.

$8+(13+5)$ means that 13 and 5 are to be added and their sum added to 8. It is clear that 13 and 5 may be added separately or together without altering the result.

Thus $8+(13+5)=8+13+5=26$.

Similarly $a+(b+c)$ means that the sum of b and c is to be added to a .

Thus $a+(b+c)=a+b+c$.

$8 + (13 - 5)$ means that to 8 we are to add the excess of 13 over 5; now if we add 13 to 8 we have added 5 too much, and must therefore take 5 from the result.

Thus $8 + (13 - 5) = 8 + 13 - 5 = 16$.

Similarly $a + (b - c)$ means that to a we are to add b , diminished by c .

Thus $a + (b - c) = a + b - c \dots \dots \dots (1).$

In like manner,

$a + b - c + (d - e - f) = a + b - c + d - e - f \dots \dots \dots (2).$

Conversely,

$a + b - c + d - e - f = a + b - c + (d - e - f) \dots \dots \dots (3).$

Again,

$a - b + c = a + c - b,$ [Art. 20.]

= the sum of a and $c - b$,

= the sum of a and $-b + c,$ [Art. 20.]

therefore $a - b + c = a + (-b + c) \dots \dots \dots (4).$

By considering the results (1), (2), (3), (4) we are led to the following rule:

Rule. When an expression within brackets is preceded by the sign +, the brackets can be removed without making any change in the expression.

Conversely: Any part of an expression may be enclosed within brackets and the sign + prefixed, the sign of every term within the brackets remaining unaltered.

Thus the expression $a - b + c - d + e$ may be written in any of the following ways,

$$a + (-b + c - d + e),$$

$$a - b + (c - d + e),$$

$$a - b + c + (-d + e).$$

22. The expression $a - (b + c)$ means that from a we are to take the sum of b and c . The result will be the same whether b and c are subtracted separately or in one sum. Thus

$$a - (b + c) = a - b - c.$$

Again, $a - (b - c)$ means that from a we are to subtract the excess of b over c . If from a we take b we get $a - b$; but by so doing we shall have taken away c too much, and must therefore add c to $a - b$. Thus

$$a - (b - c) = a - b + c.$$

In like manner,

$$a - b - (c - d - e) = a - b - c + d + e.$$

Accordingly the following rule may be enunciated:

Rule. When a sign -, the brackets are reversed.

Conversely, when brackets are within the brackets, the sign is reversed.

Thus the following rule may be enunciated:

We have

I. Addition

Thus $a + b$

This is known as Subtraction

II. The result

Thus $a + b$

This is known as Subtraction

23. When we have seen as a single like term, it may not be collected. a and b , all the addition and subtraction.

Also by this is, the algebraic rule. It will be seen in a wider sense of Arithmetic, algebra bears that meaning. The sum of the relative numbers.

Rule. When an expression within brackets is preceded by the sign $-$, the brackets may be removed if the sign of every term within the brackets be changed.

Conversely: Any part of an expression may be enclosed within brackets and the sign $-$ prefixed, provided the sign of every term within the brackets be changed.

Thus the expression $a - b + c + d - e$ may be written in any of the following ways,

$$a - (+b - c - d + e),$$

$$a - b - (-c - d + e),$$

$$a - b + c - (-d + e).$$

We have now established the following results:

I. *Additions and subtractions may be made in any order.*

$$\begin{aligned} \text{Thus } a + b - c + d - e - f &= a - c + b + d - f - e \\ &= a - c - f + d + b - e. \end{aligned}$$

This is known as the **Commutative Law for Addition and Subtraction.**

II. *The terms of an expression may be grouped in any manner.*

$$\begin{aligned} \text{Thus } a + b - c + d - e - f &= (a + b) - c + (d - e) - f \\ &= a + (b - c) + (d - e) - f = a + b - (c - d) - (e + f). \end{aligned}$$

This is known as the **Associative Law for Addition and Subtraction.**

Addition of Unlike Terms.

23. When two or more *like* terms are to be added together we have seen that they may be collected and the result expressed as a *single* like term. If, however, the terms are *unlike* they cannot be collected; thus in finding the sum of two unlike quantities a and b , all that can be done is to connect them by the sign of addition and leave the result in the form $a + b$.

Also by the rules for removing brackets, $a + (-b) = a - b$; that is, the algebraic sum of a and $-b$ is written in the form $a - b$.

It will be observed that in Algebra the word *sum* is used in a wider sense than in Arithmetic. Thus, in the language of Arithmetic, $a - b$ signifies that b is to be subtracted from a , and bears that meaning only; but in Algebra it is also taken to mean the sum of the two quantities a and $-b$ without any regard to the relative magnitudes of a and b .

Example 1. Find the sum of $3a - 5b + 2c$; $2a + 3b - d$; $-4a + 2b$.

$$\begin{aligned}\text{The sum} &= (3a - 5b + 2c) + (2a + 3b - d) + (-4a + 2b) \\ &= 3a - 5b + 2c + 2a + 3b - d - 4a + 2b \\ &= 3a + 2a - 4a - 5b + 3b + 2b + 2c - d \\ &= a + 2c - d,\end{aligned}$$

by collecting like terms.

The addition is, however, more conveniently effected by the following rule:

Rule. Arrange the expressions in lines so that the like terms may be in the same vertical columns: then add each column beginning with that on the left.

$$\begin{array}{r} 3a - 5b + 2c \\ 2a + 3b \quad - d \\ - 4a + 2b \\ \hline a \quad + 2c - d \end{array}$$

The algebraical sum of the terms in the first column is a , that of the terms in the second column is zero. The single terms in the third and fourth columns are brought down without change.

Example 2. Add together $-5ab + 6bc - 7ac$; $8ab + 3ac - 2ad$; $-2ab + 4ac + 5ad$; $bc - 3ab + 4ad$.

$$\begin{array}{r} -5ab + 6bc - 7ac \\ 8ab \quad + 3ac - 2ad \\ -2ab \quad + 4ac + 5ad \\ -3ab + bc \quad + 4ad \\ -2ab + 7bc \quad + 7ad \\ \hline \end{array}$$

Here we first rearrange the expressions so that like terms are in the same vertical columns, and then add up each column separately.

EXAMPLES III. a.

Find the sum of

1. $a + 2b - 3c$; $-3a + b + 2c$; $2a - 3b + c$.
2. $3a + 2b - c$; $-a + 3b + 2c$; $2a - b + 3c$.
3. $-3x + 2y + z$; $x - 3y + 2z$; $2x + y - 3z$.
4. $-x + 2y + 3z$; $3x - y + 2z$; $2x + 3y - z$.
5. $4a + 3b + 5c$; $-2a + 3b - 8c$; $a - b + c$.
6. $-15a - 19b - 18c$; $14a + 15b + 8c$; $a + 5b + 9c$.
7. $25a - 15b + c$; $13a - 10b + 4c$; $a + 20b - c$.
8. $-16a - 10b + 5c$; $10a + 5b + c$; $6a + 5b - c$.
9. $5ax - 7by + cz$; $ax + 2by - cz$; $-3ax + 2by + 3cz$.
10. $20p + q - r$; $p - 20q + r$; $p + q - 20r$.

Add t

11. $-5a$

12. $15ab$

13. $5ab$

14. $pq +$

15. $x + y$

16. $2a -$

17. $4xy -$

18. $17ab$

19. $47x -$

20. $-17b$

24. Ea
sion of the
the degree
three dimen
five dimen

A numer
are each of

The deg
highest dim
expression o
the fifth deg
dimensions
letters it inv
is said to be

A compou
its terms are
homogeneous

25. Diffe
thus the res
pressed by a

Similarly
 $5a^2b^2 - 3ab^3 -$
cannot be ab

E. A.

$-d$; $-4a+2b$.
2b)

ected by the

the like terms
column begin-

e terms in the
e terms in the
e single terms
columns are
ge.

$b+3ac-2ad$;

ange the ex-
terms are in
ns, and then
arately.

Add together the following expressions :

11. $-5ab+6bc-7ca$; $8ab-4bc+3ca$; $-2ab-2bc+4ca$.
12. $15ab-27bc-6ca$; $14ab-18bc+10ca$; $45bc-3ca-49ab$.
13. $5ab+bc-3ca$; $ab-bc+ca$; $-ab+2ca+bc$.
14. $pq+qr-rp$; $-pq+qr+rp$; $pq-qr+rp$.
15. $x+y+z$; $2x+3y-2z$; $3x-4y+z$.
16. $2a-3b+c$; $15a-21b-8c$; $24b+7c+3a$.
17. $4xy-9yz+2zx$; $-25xy+24yz-zx$; $23xy-15yz+zx$.
18. $17ab-13bc+8ca$; $-5ab+9bc-7ca$; $-7bc-ca+2ab$.
19. $47x-63y+z$; $-25x+15y-3z$; $-22x+15z+48y$.
20. $-17b-2c+23a$; $-9a+15b+7c$; $-13a+3b-4c$.

Dimension and Degree.

Ascending and Descending Powers.

24. Each of the letters composing a term is called a **dimension** of the term, and the number of letters involved is called the **degree** of the term. Thus the product abc is said to be of *three dimensions*, or of the *third degree*; and ax^4 is said to be of *five dimensions*, or of the *fifth degree*.

A numerical coefficient is not counted. Thus $8a^2b^5$ and a^2b^5 are each of *seven dimensions*.

The **degree of an expression** is the degree of the term of highest dimensions contained in it; thus a^4-8a^3+3a-5 is an *expression of the fourth degree*, and $a^2x-7b^2x^3$ is an *expression of the fifth degree*. But it is sometimes useful to speak of the dimensions of an expression with regard to some one of the letters it involves. For instance the expression ax^3-bx^2+cx-d is said to be of *three dimensions in x*.

A compound expression is said to be **homogeneous** when all its terms are of the same dimensions. Thus $8a^6-a^4b^2+9ab^5$ is a *homogeneous expression of six dimensions*.

25. Different powers of the same letter are unlike terms; thus the result of adding together $2x^3$ and $3x^2$ cannot be expressed by a single term, but must be left in the form $2x^3+3x^2$.

Similarly the algebraical sum of $5a^2b^2$, $-3ab^3$, and $-b^4$ is $5a^2b^2-3ab^3-b^4$. This expression is in its simplest form and cannot be abridged.

In adding together several algebraical expressions containing terms with different powers of the same letter, it will be found convenient to arrange all expressions in *descending* or *ascending* powers of that letter. This will be made clear by the following examples.

Example 1. Add together $3x^3 + 7 + 6x - 5x^2$; $2x^2 - 8 - 9x$; $4x - 2x^3 + 3x^2$; $3x^3 - 9x - x^2$; $x - x^2 - x^3 + 4$.

$$\begin{array}{r}
 3x^3 - 5x^2 + 6x + 7 \\
 2x^2 - 9x - 8 \\
 -2x^3 + 3x^2 + 4x \\
 3x^3 - x^2 - 9x \\
 -x^3 - x^2 + x + 4 \\
 \hline
 3x^3 - 2x^2 - 7x + 3
 \end{array}$$

In writing down the first expression we put in the first term the highest power of x , in the second term the next highest power, and so on till the last term, in which x does not appear. The other expressions are arranged in the same way, so that in each column we have like powers of the same letter.

Example 2. Add together $3ab^2 - 2b^3 + a^3$; $5a^2b - ab^2 - 3a^3$; $8a^3 + 5b^3$; $9a^2b - 2a^3 + ab^2$.

$$\begin{array}{r}
 -2b^3 + 3ab^2 + a^3 \\
 -ab^2 + 5a^2b - 3a^3 \\
 5b^3 + 8a^3 \\
 ab^2 + 9a^2b - 2a^3 \\
 \hline
 3b^3 + 3ab^2 + 14a^2b + 4a^3
 \end{array}$$

Here each expression contains powers of two letters, and is arranged according to *descending* powers of b , and *ascending* powers of a .

EXAMPLES III. b.

Find the sum of

- $2ab + 3ca + 6abc$; $-5ab + 2bc - 5abc$; $3ab - 2bc - 3ca$.
- $2x^2 - 2xy + 3y^2$; $4y^2 + 5xy - 2x^2$; $x^2 - 2xy - 6y^2$.
- $3a^2 - 7ab - 4b^2$; $-6a^2 + 9ab - 3b^2$; $4a^2 + ab + 5b^2$.
- $x^3 + xy - y^2$; $-x^2 + yz + y^2$; $-x^2 + xz + z^2$.
- $-x^2 - 3xy + 3y^2$; $3x^2 + 4xy - 5y^2$; $x^2 + xy + y^2$.
- $x^3 - x^2 + x - 1$; $2x^2 - 2x + 2$; $-3x^3 + 5x + 1$.
- $2x^3 - x^2 - x$; $4x^3 + 8x^2 + 7x$; $-6x^3 - 6x^2 + x$.
- $9x^2 - 7x + 5$; $-14x^2 + 15x - 6$; $20x^2 - 40x - 17$.
- $10x^3 + 5x + 8$; $3x^3 - 4x^2 - 6$; $2x^3 - 2x - 3$.
- $a^3 - ab + bc$; $ab + b^3 - ca$; $ca - bc + c^3$.
- $5a^3 - 3c^3 + d^3$; $b^3 - 2a^3 + 3d^3$; $4c^3 - 2a^3 - 3d^3$.

Find th

- $6x^3$
- $a^3 - a$
- $x^2 + y^2$
- $x^3 - 2y$
- $x^3 + 3a$
- $a^3 + 5a$
- $x^5 - 4x$
- $a^3 - 4a$
- $x^3 - 4x$

Add toge

- $\frac{1}{2}a - \frac{1}{3}b$
- $-\frac{1}{3}a -$
- $-2a +$
- $-\frac{1}{8}a$
- $\frac{2}{3}x^2 + \frac{1}{5}x$
- $3a^2 - \frac{2}{5}a$
- $\frac{5}{8}x^2 - \frac{1}{5}x$
- $-\frac{3}{4}x^3 +$
- $\frac{3}{8}x^2 - \frac{5}{3}x$
- $\frac{1}{2}a^3 - 2a$

Find the sum of

12. $6x^3 - 2x + 1$; $2x^3 + x + 6$; $x^2 - 7x^3 + 2x - 4$.
13. $a^3 - a^2 + 3a$; $3a^3 + 4a^2 + 8a$; $5a^3 - 6a^2 - 11a$.
14. $x^2 + y^2 - 2xy$; $2x^2 - 3y^2 - 4yz$; $2x^2 - 2x^2 - 3xz$.
15. $x^3 - 2y^3 + x$; $y^3 - 2x^3 + y$; $x^2 + 2y^2 - x + y^3$.
16. $x^3 + 3x^2y + 3xy^2$; $-3x^2y - 6xy^2 - x^3$; $3x^2y + 4xy^2$.
17. $a^3 + 5ab^2 + b^3$; $b^3 - 10ab^2 - a^3$; $5ab^2 - 2b^3 + 2^2b$.
18. $x^5 - 4x^4y - 5x^3y^2$; $3x^4y + 2x^3y^2 - 6xy^4$; $3x^3y^3 + 6xy^4 - y^5$.
19. $a^3 - 4a^2b + 6abc$; $a^2b - 10abc + c^3$; $b^3 + 3a^2b + abc$.
20. $x^3 - 4x^2y + 6xy^2$; $2x^2y - 3xy^2 + 2y^3$; $y^3 + 3x^2y + 4xy^2$.

Add together the following expressions :

21. $\frac{1}{2}a - \frac{1}{3}b$; $-a + \frac{2}{3}b$; $\frac{3}{4}a - b$.
22. $-\frac{1}{3}a - \frac{1}{4}b$; $-\frac{2}{3}a + \frac{3}{4}b$; $-2a - b$.
23. $-2a + \frac{5}{6}c$; $-\frac{1}{3}a - 2b$; $\frac{8}{3}b - 3c$.
24. $-\frac{1}{8}a - \frac{1}{4}c$; $2a - 3b$; $\frac{1}{5}b - c$.
25. $\frac{2}{3}x^2 + \frac{1}{5}xy - \frac{1}{4}y^2$; $-x^2 - \frac{2}{3}xy + 2y^2$; $\frac{2}{3}x^2 - xy - \frac{5}{4}y^2$.
26. $3a^2 - \frac{2}{5}ab - \frac{1}{2}b^2$; $-\frac{3}{2}a^2 + 2ab - \frac{2}{3}b^2$; $-\frac{2}{3}a^2 - ab + b^2$.
27. $\frac{5}{8}x^2 - \frac{1}{5}xy + \frac{3}{10}y^2$; $-\frac{3}{4}x^2 + \frac{1}{5}xy - y^2$; $\frac{1}{2}x^2 - xy + \frac{1}{5}y^2$.
28. $-\frac{3}{4}x^3 + 5ax^2 - \frac{5}{8}a^2x$; $x^3 - \frac{3}{8}ax^2 + \frac{1}{2}a^2x$; $-\frac{1}{2}x^3 + \frac{3}{4}a^2x$.
29. $\frac{3}{8}x^2 - \frac{5}{3}xy - 7y^2$; $\frac{2}{3}xy + \frac{1}{6}y^2$; $-\frac{5}{8}x^2 + 4y^2$.
30. $\frac{1}{2}a^2 - 2a^2b - \frac{3}{2}b^3$; $\frac{3}{2}a^2b - \frac{3}{4}ab^2 + 2b^3$; $-\frac{3}{2}a^3 + ab^2 + \frac{1}{2}b^2$.

CHAPTER IV.

SUBTRACTION.

26. THE simplest cases of Subtraction have already come under the head of addition of *like* terms, of which some are negative. [Art. 18.]

Thus

$$\begin{aligned} 5a - 3a &= 2a, \\ 3a - 7a &= -4a, \\ -3a - 6a &= -9a. \end{aligned}$$

Also, by the rule for removing brackets [Art. 22],

$$\begin{aligned} 3a - (-8a) &= 3a + 8a \\ &= 11a, \end{aligned}$$

and

$$\begin{aligned} -3a - (-8a) &= -3a + 8a \\ &= 5a. \end{aligned}$$

Subtraction of Unlike Terms.

27. The method is shewn in the following example.

Example. Subtract $3a - 2b - c$ from $4a - 3b + 5c$.

The difference

$$\begin{aligned} &= 4a - 3b + 5c - (3a - 2b - c) \\ &= 4a - 3b + 5c - 3a + 2b + c \\ &= 4a - 3a - 3b + 2b + 5c + c \\ &= a - b + 6c. \end{aligned}$$

The expression to be subtracted is first enclosed in brackets with a minus sign prefixed, then on removal of the brackets the like terms are combined by the rules already explained in Art. 18.

It is, however, more convenient to arrange the work as follows, the signs of all the terms in the lower line being changed.

$$\begin{array}{r} 4a - 3b + 5c \\ - 3a + 2b + c \\ \hline a - b + 6c \end{array}$$

by addition,

The like terms are written in the same vertical column, and each column is treated separately.

Rule. Change the sign of every term in the expression to be subtracted, and add to the other expression.

Note. It is not necessary that in the expression to be subtracted the signs should be *actually* changed; the operation of changing signs ought to be performed mentally.

CHAP. IV.]

Example 1.

$$\begin{array}{r} 5x^2 + xy \\ 2x^2 + 8xy - 7y^2 \\ \hline 3x^2 - 7xy + 7y^2 \end{array}$$

Example 2.

Terms contained must stand

$$\begin{array}{r} -x^3 + \dots \\ 3x^2 - 2x \\ \hline -x^3 - 3x^2 + 2x + \dots \end{array}$$

The re-arranging is convenient, ing powers of x .

Subtract

1. $4a - 3b + c$
2. $a - 3b + 5c$
3. $2x - 8y + z$
4. $15a - 27b +$
5. $-10x - 14y$
6. $-11ab + 6c$
7. $4a - 3b + 15$
8. $-16x - 18y$
9. $ab + cd - ac$
10. $-ab + cd -$

From

11. $3ab + 5cd -$
12. $yz - zx + xy$
13. $-2x^3 - x^2 -$
14. $-8x^2y + 15a$
15. $\frac{1}{2}a - b + \frac{1}{3}c$
16. $\frac{3}{4}x + y - z$ tak
17. $-a - 3b$ tak
18. $\frac{1}{2}x - \frac{2}{3}y + \frac{1}{4}$
19. $-\frac{2}{3}x - \frac{3}{4}y -$
20. $-\frac{1}{2}x + \frac{2}{3}y -$

Example 1. From $5x^2 + xy$ take $2x^2 + 8xy - 7y^2$.

$$\begin{array}{r} 5x^2 + xy \\ 2x^2 + 8xy - 7y^2 \\ 3x^2 - 7xy + 7y^2 \end{array}$$

In the first column we combine mentally $5x^2$ and $-2x^2$, the algebraic sum of which is $3x^2$. In the last column the sign of the term $-7y^2$ has to be changed before it is put down in the result.

Example 2. Subtract $3x^2 - 2x$ from $1 - x^3$.

Terms containing different powers of the same letter being *unlike* must stand in different columns.

$$\begin{array}{r} -x^3 \qquad +1 \\ \quad 3x^2 - 2x \\ -x^3 - 3x^2 + 2x + 1 \end{array}$$

In the first and last columns, as there is nothing to be subtracted, the terms are put down without change of sign. In the second and third columns each sign has to be changed.

The re-arrangement of terms in the first line is not *necessary*, but it is convenient, because it gives the result of subtraction in descending powers of x .

EXAMPLES IV. a.

Subtract

- $4a - 3b + c$ from $2a - 3b - c$.
- $a - 3b + 5c$ from $4a - 8b + c$.
- $2x - 8y + z$ from $15x + 10y - 18z$.
- $15a - 27b + 8c$ from $10a + 3b + 4c$.
- $-10x - 14y + 15z$ from $x - y - z$.
- $-11ab + 6cd$ from $-10bc + ab - 4cd$.
- $4a - 3b + 15c$ from $25a - 16b - 18c$.
- $-16x - 18y - 15z$ from $-5x + 8y + 7z$.
- $ab + cd - ac - bd$ from $ab + cd + ac + bd$.
- $-ab + cd - ac + bd$ from $ab - cd + ac - bd$.

From

- $3ab + 5cd - 4ac - 6bd$ take $3ab + 6cd - 3ac - 5bd$.
- $yz - zx + xy$ take $-xy + yz - zx$.
- $-2x^3 - x^2 - 3x + 2$ take $x^3 - x + 1$.
- $-8x^2y + 15xy^2 + 10xyz$ take $4x^2y - 6xy^2 - 5xyz$.
- $\frac{1}{2}a - b + \frac{1}{3}c$ take $\frac{1}{3}a + \frac{1}{2}b - \frac{1}{2}c$.
- $\frac{3}{4}x + y - z$ take $\frac{1}{2}x - \frac{1}{2}y - \frac{1}{3}z$.
- $-a - 3b$ take $\frac{3}{2}a + \frac{1}{3}b - \frac{1}{2}c$.
- $\frac{1}{2}x - \frac{3}{4}y + \frac{1}{10}z$ take $-\frac{1}{2}x + \frac{4}{7}y - \frac{1}{10}z$.
- $-\frac{2}{3}x - \frac{3}{8}y - 5z$ take $\frac{2}{3}x - \frac{3}{8}y - \frac{11}{3}z$.
- $-\frac{1}{2}x + \frac{2}{3}y - \frac{1}{6}$ take $\frac{1}{3}x - \frac{3}{2}y - \frac{1}{6}$.

EXAMPLES IV. b.

From

1. $3xy - 5yz + 8zx$ take $-4xy + 2yz - 10zx$.
2. $-8x^2y^2 + 15x^3y + 13xy^3$ take $4x^2y^2 + 7x^3y - 8xy^3$.
3. $-8 + 6ab + a^2b^2$ take $4 - 3ab - 5a^2b^2$.
4. $a^2bc + b^2ca + c^2ab$ take $3a^2bc - 5b^2ca - 4c^2ab$.
5. $-7a^2b + 8ab^2 + cd$ take $5a^2b - 7ab^2 + 6cd$.
6. $-8x^2y + 5xy^2 - x^2y^2$ take $8x^2y - 5xy^2 + x^2y^2$.
7. $10a^2b^2 + 15ab^2 + 8a^2b$ take $-10a^2b^2 + 15ab^2 - 8a^2b$.
8. $4x^2 - 3x + 2$ take $-5x^2 + 6x - 7$.
9. $x^3 + 11x^2 + 4$ take $8x^2 - 5x - 3$.
10. $-8a^2x^2 + 5x^2 + 15$ take $9a^2x^2 - 8x^2 - 5$.

Subtract

11. $x^3 - x^2 + x + 1$ from $x^3 + x^2 - x + 1$.
12. $3xy^2 - 3x^2y + x^3 - y^3$ from $x^3 + 3x^2y + 3xy^2 + y^3$.
13. $b^3 + c^3 - 2abc$ from $a^3 + b^3 - 3abc$.
14. $7xy^2 - y^3 - 3x^2y + 5x^3$ from $8x^3 + 7x^2y - 3xy^2 - y^3$.
15. $x^4 + 5 + x - 3x^3$ from $5x^4 - 8x^3 - 2x^2 + 7$.
16. $a^3 + b^3 + c^3 - 3abc$ from $7abc - 3a^3 + 5b^3 - c^3$.
17. $1 - x + x^5 - x^4 - x^3$ from $x^4 - 1 + x - x^2$.
18. $7a^4 - 8a^3 + 3a^2 + a$ from $a^2 - 5a^3 - 7 + 7a^5$.
19. $10a^2b + 8ab^2 - 8a^3b^3 - b^4$ from $5a^2b - 6ab^2 - 7a^3b^3$.
20. $a^3 - b^3 + 8ab^2 - 7a^2b$ from $-8ab^2 + 15a^2b + b^3$.

From

21. $\frac{1}{2}x^2 - \frac{1}{3}xy - \frac{3}{2}y^2$ take $-\frac{3}{2}x^2 + xy - y^2$.
22. $\frac{2}{3}a^2 - \frac{5}{2}a - 1$ take $-\frac{2}{3}a^2 + a - \frac{1}{2}$.
23. $\frac{1}{3}x^2 - \frac{1}{2}x + \frac{1}{8}$ take $\frac{1}{3}x - 1 + \frac{1}{2}x^2$.
24. $\frac{3}{8}x^2 - \frac{2}{3}ax$ take $\frac{1}{3} - \frac{1}{4}x^2 - \frac{5}{8}ax$.
25. $\frac{3}{4}x^3 - \frac{1}{3}xy^2 - y^2$ take $\frac{1}{2}x^2y - \frac{5}{6}y^2 - \frac{1}{3}xy^3$.
26. $\frac{1}{8}a^3 - 2ax^2 - \frac{1}{3}a^2x$ take $\frac{1}{3}a^2x + \frac{1}{4}a^3 - \frac{3}{2}ax^2$.

1. Sim

2. To
 $a - 4c + 7b$

3. Whe

4. Defin
 $x^2 + 7x$, find

5. From

6. Subtr

7. Distin
terms in the8. Write
together x,

9. Subtr

10. If th
what will -11. Writ
 $2a$ by the su12. When
 $-2x^2z, 3y^4$.13. Add
when $1 - 2y^2$

14. Expla

15. If $x =$

16. Simpl

MISCELLANEOUS EXAMPLES I.

1. Simplify (1) $4x - 2x^2 - (2x - 3x^2)$;
(2) $3a - 4b - (3b + a) - (5a - 8b)$.
 2. To the sum of $2a - 3b - 2c$ and $2b - a + 7c$ add the sum of $a - 4c + 7b$ and $c - 6b$.
 3. When $x=3$, $y=2$, $z=0$, find the value of
(1) $x^2 + \frac{3}{2}y^3 - xyz^3$; (2) $\frac{1}{4}x^3y^4 + \frac{5z^2}{6}$.
 4. Define *index, coefficient*. In the expressions $4x^2 + 3x$, $2x^3 + x^2$, $x^2 + 7x$, find (1) the sum of the indices, (2) the sum of the coefficients.
 5. From $5x^3 + 3x - 1$ take the sum of
 $2x - 5 + 7x^2$ and $3x^2 + 4 - 2x^3 + x$.
 6. Subtract $3a - 7a^3 + 5a^2$ from the sum of
 $2 + 8a^2 - a^3$ and $2a^3 - 3a^2 + a - 2$.
-
7. Distinguish between *like* and *unlike* terms. Pick out the like terms in the expression
 $a^3 - 3ab + b^2 - 2a^3 - a^2 + 3b^2 + 5ab + 7a^2$.
 8. Write down in as many ways as possible the result of adding together x , y , and z .
 9. Subtract $5x^2 + 3x - 1$ from $2x^3$, and add the result to
 $3x^2 + 3x - 1$.
 10. If the number of pounds I possess is represented by $+a$, what will $-a$ denote?
 11. Write down in algebraical symbols the result of diminishing $2a$ by the sum of $3b$ and $5c$.
 12. When $x=1$, $y=2$, $z=3$, find the value of the sum of $5x^2$, $-2x^2z$, $3y^2$. Also find the value of $2xz^2 - 3y^2$.
-
13. Add the sum of $2y - 3y^2$ and $1 - 5y^3$ to the remainder left when $1 - 2y^2 + y$ is subtracted from $5y^3$.
 14. Explain clearly why $x - (y - z) = x - y + z$.
 15. If $x=4$, $y=3$, $z=2$, $a=0$, find the value of
 $3x^2 - 2yz - ax + 5ax^2y$.
 16. Simplify $2a - b - (3a - 2b) + (2a - 3b) - (a - 2b)$.

17. Find the algebraical sum of the like terms in the expression
 $5a^3 - 4a^2b + b^3 + 6a^3b + 7ab^2 - 3a^2b + 4ab^2 + 8a^2b$.
18. A boy works $x + y$ sums, of which only $y - 2z$ are right; how many are wrong?
-
19. In the expression $3a^3 - 7a^2b + b^4$, point out the highest power, the lowest power, the positive terms, and the coefficient of a^2 .
20. Take $x^2 - y^2$ from $3xy - 4y^2$, and add the remainder to the sum of $4xy - x^2 - 3y^2$ and $2x^2 + 6y^2$.
21. If $x=1$, $y=3$, $z=5$, $w=0$, find the value of
 $\sqrt{(3xy)} + \sqrt{(5xz)} + \sqrt{(3yw)}$.
22. What is the *degree* of a term in an algebraical expression? In the expression $4x^6 - 3x^5a^2 + a^3$, what is the degree of the negative term?
23. Find the sum of $5a - 7b + c$ and $3b - 9a$, and subtract the result from $c - 4b$.
24. If $x=3$, $y=4$, $p=8$, $q=10$, find the value of

$$xyp + \frac{2y}{p-y} + 2q.$$
-
25. If x represents the date 10 A.D. what will $-3x$ stand for?
26. Add together $3x^2 - 7x + 5$ and $2x^3 + 5x - 3$, and diminish the result by $3x^2 + 2$.
27. In the expression
 $4a^2b^3 - b^4 + 3a^3b^2 + 5b^5 - ab^3x + 2x^2ab + abx^4 - a^2b^3,$
 point out which terms are *like*, and which are homogeneous. What is the degree of the expression?
28. Express in algebraical symbols the excess of the sum of a and b over c diminished by d .
29. A man walks $2a - b$ miles due North from a fixed point C, and then walks a distance $3a + 2b$ miles due South; what is his final position with regard to O?
30. What expression must be added to $5x^2 - 7x + 2$ to produce $7x^2 - 1$?

28. Mu
 addition.
 Thus

Here the
 times we ta
 Again

the number
 Also $3 \times$
 whole num

29. Wh
 positive wh
 operation p
 unity produ
 perform on
 gives $\frac{3}{4}$; tha
 three of the
 result of ta

Hence

Also, by t

CHAPTER V.

MULTIPLICATION.

28. MULTIPLICATION in its primary sense signifies repeated addition.

$$\begin{aligned}\text{Thus} \quad 3 \times 4 &= 3 \text{ taken 4 times} \\ &= 3+3+3+3.\end{aligned}$$

Here the multiplier contains four units, and the number of times we take 3 is the same as the number of units in 4.

$$\begin{aligned}\text{Again} \quad a \times b &= a \text{ taken } b \text{ times} \\ &= a+a+a+\dots,\end{aligned}$$

the number of terms being b .

Also $3 \times 4 = 4 \times 3$; and so long as a and b denote positive whole numbers, it is easy to show that $a \times b = b \times a$.

29. When the quantities to be multiplied together are not positive whole numbers, we may define multiplication as *an operation performed on one quantity which when performed on unity produces the other*. For example, to multiply $\frac{4}{5}$ by $\frac{3}{7}$, we perform on $\frac{4}{5}$ that operation which when performed on unity gives $\frac{3}{7}$; that is, we must divide $\frac{4}{5}$ into seven equal parts and take three of them. Now each part will be equal to $\frac{4}{5 \times 7}$, and the result of taking three of such parts is expressed by $\frac{4 \times 3}{5 \times 7}$.

$$\text{Hence} \quad \frac{4}{5} \times \frac{3}{7} = \frac{4 \times 3}{5 \times 7}.$$

Also, by the last article,

$$\begin{aligned}\frac{4 \times 3}{5 \times 7} &= \frac{3 \times 4}{7 \times 5} = \frac{3}{7} \times \frac{4}{5} \\ \therefore \frac{4}{5} \times \frac{3}{7} &= \frac{3}{7} \times \frac{4}{5}.\end{aligned}$$

The reasoning is clearly general, and we may now say that $a \times b = b \times a$, where a and b are any positive quantities, integral or fractional.

In the same way it easily follows that

$$\begin{aligned} abc &= a \times b \times c \\ &= (a \times b) \times c = (b \times a) \times c = bac \\ &= b \times (a \times c) = b \times c \times a = bca; \end{aligned}$$

that is, *the factors of a product may be taken in any order.* This is the **Commutative Law for Multiplication.**

Example. $2a \times 3b \times c = 2 \times 3 \times a \times b \times c = 6abc.$

30. Again, *the factors of a product may be grouped in any way we please.*

$$\begin{aligned} \text{Thus } abcd &= a \times b \times c \times d \\ &= (ab) \times (cd) = a \times (bc) \times d = a \times (bcd). \end{aligned}$$

This is the **Associative Law for Multiplication.**

31. Since, by definition, $a^3 = aaa$, and $a^5 = aaaaa$,

$$\therefore a^2 \times a^5 = aaa \times aaaaa = aaaaaaa = a^7 = a^{2+5};$$

that is, *the index of a letter in a product is the sum of its indices in the factors of the product.* This is the **Index Law for Multiplication.**

Again, $5a^2 = 5aa$, and $7a^3 = 7aaa$.

$$\therefore 5a^2 \times 7a^3 = 5 \times 7 \times aaaa = 35a^5.$$

When the expressions to be multiplied together contain powers of different letters, a similar method is used.

$$\begin{aligned} \text{Example. } 5a^3b^2 \times 8a^2bx^3 &= 5aaabb \times 8aabbxx \\ &= 40a^5b^3x^3. \end{aligned}$$

Note. The beginner must be careful to observe that in this process of multiplication the indices of one letter cannot combine in any way with those of another. Thus the expression $40a^5b^3x^3$ admits of no further simplification.

32. **Rule.** To multiply two simple expressions together, multiply the coefficients together and prefix their product to the product of the different letters, giving to each letter an index equal to the sum of the indices that letter has in the separate factors.

The rule may be extended to cases where more than two expressions are to be multiplied together.

v.]

Exam

The

The p
tinued

Exam

The

Multi

33.

together

Also

diminish

Simila

Thus i
a single
term of t
as the D**Note.**
positive v

Exam

Find t

1. $5x^2 \times$ 4. $6xy^2$ 7. $2a^3b^2$ 10. $5a^4b^3$ 13. $3a^5b^4$ 16. $5x^3y$

Example 1. Find the product of x^2 , x^3 , and x^8 .

The product $= x^2 \times x^3 \times x^8 = x^{2+3+8} = x^{13}$.

The product of three or more expressions is called **the continued product**.

Example 2. Find the continued product of $5x^2y^3$, $8y^2z^5$, and $3xz^4$.

The product $= 5x^2y^3 \times 8y^2z^5 \times 3xz^4 = 120x^3y^5z^9$.

Multiplication of a Compound Expression by a Simple Expression.

33. By definition,

$(a+b)m = m + m + m + \dots$ taken $a+b$ times

$= (m + m + m + \dots$ taken a times),

together with $(m + m + m + \dots$ taken b times)

$= am + bm \dots \dots \dots (1).$

Also $(a-b)m = m + m + m + \dots$ taken $a-b$ times

$= (m + m + m + \dots$ taken a times),

diminished by $(m + m + m + \dots$ taken b times)

$= am - bm \dots \dots \dots (2).$

Similarly $(a-b+c)m = am - bm + cm.$

Thus it appears that *the product of a compound expression by a single factor is the algebraic sum of the partial products of each term of the compound expression by that factor.* This is known as the **Distributive Law for Multiplication**.

Note. It should be observed that for the present a, b, c denote positive whole numbers, and that a is supposed greater than b .

Examples. $3(2a+3b-4c) = 6a+9b-12c.$

$(4x^2-7y-8z^3) \times 3xy^2 = 12x^3y^2 - 21xy^3 - 24xy^2z^3.$

EXAMPLES V. a.

Find the value of

- | | | |
|---------------------------------|------------------------------|------------------------------------|
| 1. $5x^2 \times 7x^5.$ | 2. $4a^3 \times 5a^8.$ | 3. $7ab \times 8a^3b^3.$ |
| 4. $6xy^2 \times 5x^3.$ | 5. $8a^3b \times b^5.$ | 6. $2abc \times 3ac^2.$ |
| 7. $2a^3b^3 \times 2a^2b^3.$ | 8. $5a^2b \times 2a.$ | 9. $4a^2b^3 \times 7a^5.$ |
| 10. $5a^4b^3 \times x^2y^2.$ | 11. $x^3y^3 \times 6a^2x^4.$ | 12. $abc \times xyz.$ |
| 13. $3a^4b^7x^5 \times 5a^3bx.$ | 14. $4a^3bx \times 7b^2x^4.$ | 15. $5a^2x \times 8cx.$ |
| 16. $5x^2y^3 \times 6a^2x^3.$ | 17. $2x^2y \times x^5y^7.$ | 18. $3a^3x^4y^7 \times a^2x^5y^9.$ |

Multiply together :

19. $ab + bc$ and a^3b .

21. $5x + 3y$ and $2x^2$.

23. $bc + ca - ab$ and abc .

25. $5x^2y + xy^2 - 7x^2y^2$ and $3x^3$.

27. $6a^3bc - 7ab^2c^2$ and a^2b^2 .

20. $5ab - 7bx$ and $4a^2bx^3$.

22. $a^2 + b^2 - c^2$ and a^3b .

24. $5a^2 + 3b^3 - 2c^2$ and $4a^2bc^3$.

26. $6x^3 - 5x^2y + 7xy^2$ and $8x^2y^3$.

Multiplication of Compound Expressions.

34. If in Art. 33 we write $c + d$ for m in (1), we have

$$\begin{aligned}(a+b)(c+d) &= a(c+d) + b(c+d) \\ &= (c+d)a + (c+d)b \quad [\text{Art. 29.}] \\ &= ac + ad + bc + bd \dots\dots\dots(3).\end{aligned}$$

Again, from (2)

$$\begin{aligned}(a-b)(c+d) &= a(c+d) - b(c+d) \\ &= (c+d)a - (c+d)b \\ &= ac + ad - (bc + bd) \\ &= ac + ad - bc - bd \dots\dots\dots(4).\end{aligned}$$

Similarly, by writing $c - d$ for m in (1),

$$\begin{aligned}(a+b)(c-d) &= a(c-d) + b(c-d) \\ &= (c-d)a + (c-d)b \\ &= ac - ad + bc - bd \dots\dots\dots(5).\end{aligned}$$

Also, from (2)

$$\begin{aligned}(a-b)(c-d) &= a(c-d) - b(c-d) \\ &= (c-d)a - (c-d)b \\ &= ac - ad - (bc - bd) \\ &= ac - ad - bc + bd \dots\dots\dots(6).\end{aligned}$$

If we consider each term on the right-hand side of (6), and the way in which it arises, we find that

$$(+a) \times (+c) = +ac.$$

$$(-b) \times (-d) = +bd.$$

$$(-b) \times (+c) = -bc.$$

$$(+a) \times (-d) = -ad.$$

These results enable us to state what is known as the **Rule of Signs** in multiplication.

Rule of Signs. *The product of two terms with like signs is positive; the product of two terms with unlike signs is negative.*

v.]

35. multipli
Perhaps
trating
by a neg
To mu
to obtain
the resul
to be tak
But 3

Similarly
and the
second +

Thus

Hence, m
to proceea
the sign o

Not

36. A
deals sole
ligible.
enabled to

Symbol
case, and
symbols a
longer be
33 and 3
require th
 $a > b$ and
assume th
are remov
thereby.

Hencefo
and the R
used. [Se

37. To
just expla
some of th

35. The rule of signs, and especially the use of the negative multiplier, will probably present some difficulty to the beginner. Perhaps the following numerical instances may be useful in illustrating the interpretation that may be given to multiplication by a negative quantity.

To multiply 3 by -4 we must do to 3 what is done to unity to obtain -4 . Now -4 means that unity is taken 4 times and the result made negative; therefore $3 \times (-4)$ implies that 3 is to be taken 4 times and the product made negative.

But 3 taken 4 times gives $+12$;

$$\therefore 3 \times (-4) = -12.$$

Similarly -3×-4 indicates that -3 is to be taken 4 times, and the sign changed; the first operation gives -12 , and the second $+12$.

Thus

$$(-3) \times (-4) = +12.$$

Hence, *multiplication by a negative quantity indicates that we are to proceed just as if the multiplier were positive, and then change the sign of the product.*

Note on Arithmetical and Symbolical Algebra.

36. Arithmetical Algebra is that part of the science which deals solely with symbols and operations arithmetically intelligible. Starting from purely arithmetical definitions, we are enabled to prove certain fundamental laws.

Symbolical Algebra assumes these laws to be true in every case, and thence finds what meaning must be attached to symbols and operations which under unrestricted conditions no longer bear an arithmetical meaning. Thus the results of Arts. 33 and 34 were proved from arithmetical definitions which require the symbols to be positive whole numbers, such that $a > b$ and $c > d$. By the principles of symbolical Algebra we assume these results to be universally true when all restrictions are removed, and accept the interpretation to which we are led thereby.

Henceforth we are able to apply the Law of Distribution and the Rule of Signs without any restriction as to the symbols used. [See Art. 33, Note.]

37. To familiarize the beginner with the principles we have just explained we add a few examples in substitutions where some of the symbols denote negative quantities.

Example 1. If $a = -4$, find the value of a^3 .

Here $a^3 = (-4)^3 = (-4) \times (-4) \times (-4) = -64$.

By repeated applications of the rule of signs it may easily be shewn that any *odd* power of a negative quantity is *negative*, and any *even* power of a negative quantity is *positive*.

Example 2. If $a = -1$, $b = 3$, $c = -2$, find the value of $-3a^4bc^3$.

Here $-3a^4bc^3 = -3 \times (-1)^4 \times 3 \times (-2)^3$
 $= -3 \times (+1) \times 3 \times (-8)$
 $= 72$.

We write down at once $(-1)^4 = +1$, and $(-2)^3 = -8$.

EXAMPLES V. b.

If $a = -2$, $b = 3$, $c = -1$, $x = -5$, $y = 4$, find the value of

- | | | | |
|---------------------|------------------|------------------|-----------------|
| 1. $3a^2b$. | 2. $8abc^2$. | 3. $-5c^3$. | 4. $6a^2c^2$. |
| 5. $4c^3y$. | 6. $3a^3c$. | 7. $-b^2c^2$. | 8. $3a^3c^2$. |
| 9. $-7a^3bc$. | 10. $-2a^4bx$. | 11. $-4a^2c^4$. | 12. $3c^3x^3$. |
| 13. $5a^2x^2$. | 14. $-7c^4xy$. | 15. $-8ax^3$. | 16. $4c^5x^3$. |
| 17. $-5a^2b^2c^2$. | 18. $-7a^3c^3$. | 19. $8c^4x^3$. | 20. $7a^5c^4$. |

If $a = -4$, $b = -3$, $c = -1$, $f = 0$, $x = 4$, $y = 1$, find the value of

- | | |
|--|---|
| 21. $3a^2 + bx - 4cy$. | 22. $2ab^2 - 3bc^2 + 2fx$. |
| 23. $fa^2 - 2b^3 - cx^3$. | 24. $3a^2y^3 - 5b^2x - 2c^3$. |
| 25. $2a^3 - 3b^3 + 7cy^4$. | 26. $3b^2y^4 - 4b^2f - 6c^4x$. |
| 27. $2\sqrt{(ac)} - 3\sqrt{(xy)} + \sqrt{(b^2c^4)}$. | 28. $3\sqrt{(acx)} - 2\sqrt{(b^2y)} - 6\sqrt{(c^2y)}$. |
| 29. $7\sqrt{(a^2x)} - 3\sqrt{(b^4c^2)} + 5\sqrt{(f^2x)}$. | |
| 30. $3c\sqrt{(3bc)} - 5\sqrt{(4c^2y^3)} - 2cy\sqrt{(3bc^5)}$. | |

38. The following examples further illustrate the rule of signs and the law of indices.

Example 1. Multiply $4a$ by $-3b$.

By the rule of signs the product is negative; also $4a \times 3b = 12ab$;

$$\therefore 4a \times (-3b) = -12ab.$$

Example 2. Multiply $-5ab^3x$ by $-ab^3x$.

Here the absolute value of the product is $5a^2b^6x^2$, and by the rule of signs the product is positive;

$$\therefore (-5ab^3x) \times (-ab^3x) = 5a^2b^6x^2.$$

Example 3. Find the continued product of $3a^2b$, $-2a^3b^2$, $-ab^4$.

$$3a^2b \times (-2a^3b^2) = -6a^5b^3;$$

$$(-6a^5b^3) \times (-ab^4) = +6a^6b^7.$$

Thus the complete product is $6a^6b^7$.

This result, however, may be written down at once; for

$$3a^2b \times 2a^3b^2 \times ab^4 = 6a^6b^7,$$

and by the rule of signs the required product is positive.

v.]

Example

The

for

(

Multip

1. ax 3. a^2b 5. $-abc$ 7. $3xy$ 9. $-x$ 11. $-ab$ 13. $5x^2y$ 15. $-5xy$ 17. $-13x$ 19. abc

Find th

21. $2a - 3$ 23. $\frac{2}{3}a - 1$ 25. $-\frac{5}{3}a^2 - 2$ 27. $-\frac{3}{2}x^2$

39. TH

both of th

or more te

replacing

We may
any two co

Rule. 1
of the secon
prefix to t
algebraical
product. 7

Example 4. Multiply $6a^3 - \frac{5}{3}a^2b - \frac{4}{5}ab^2$ by $-\frac{3}{4}ab^2$.

The product is the algebraical sum of the partial products formed according to the rule enunciated in Art. 37; thus

$$(6a^3 - \frac{5}{3}a^2b - \frac{4}{5}ab^2) \times (-\frac{3}{4}ab^2) = -\frac{9}{2}a^4b^2 + \frac{5}{4}a^3b^3 + \frac{3}{5}a^2b^4.$$

EXAMPLES V. c.

Multiply together :

- | | |
|---|--|
| 1. ax and $-3ax$. | 2. $-2abx$ and $-7abx$. |
| 3. a^2b and $-ab^2$. | 4. $6x^2y$ and $-10xy$. |
| 5. $-abcd$ and $-3a^2b^3c^4d^5$. | 6. xyz and $-5x^2y^3z$. |
| 7. $3xy + 4yz$ and $-12xyz$. | 8. $ab - bc$ and a^2bc^3 . |
| 9. $-x - y - z$ and $-3x$. | 10. $a^2 - b^2 + c^2$ and abc . |
| 11. $-ab + bc - ca$ and $-abc$. | 12. $-2a^2b - 4ab^2$ and $-7a^2b^2$. |
| 13. $5x^2y - 6xy^2 + 8x^2y^2$ and $3xy$. | 14. $-7x^3y - 5xy^3$ and $-8x^2y^3$. |
| 15. $-5xy^2z + 3xyz^2 - 8x^2yz$ and xyz . | 16. $4x^2yz^2 - 8xyz$ and $-12x^3yz^2$. |
| 17. $-13xy^2 - 15x^2y$ and $-7x^3y^3$. | 18. $8xyz - 10x^2yz^2$ and $-xyz$. |
| 19. $abc - a^2bc - ab^2c$ and $-abc$. | 20. $-a^2bc + b^2ca - c^2ab$ and $-ab$. |

Find the product of

- | | |
|--|--|
| 21. $2a - 3b + 4c$ and $-\frac{3}{2}a$. | 22. $3x - 2y - 4$ and $-\frac{5}{8}x$. |
| 23. $\frac{2}{3}a - \frac{1}{6}b - c$ and $-\frac{2}{3}ax$. | 24. $\frac{6}{7}a^2x^2 - \frac{3}{2}ax^3$ and $-\frac{7}{3}a^3x$. |
| 25. $-\frac{5}{3}a^2x^2$ and $-\frac{3}{2}a^2 + ax - \frac{3}{5}x^2$. | 26. $-\frac{7}{2}xy$ and $-3x^2 + \frac{2}{7}xy$. |
| 27. $-\frac{3}{2}x^3y^2$ and $-\frac{1}{3}x^2 + 2y^2$. | 28. $-\frac{4}{7}x^5y^3$ and $\frac{7}{4}x^3 - \frac{4}{7}y^3$. |

39. The results of Art. 33 may be extended to the case where both of the expressions to be multiplied together contain two or more terms. For instance

$$(a - b + c)m = am - bm + cm;$$

replacing m by $x - y$, we have

$$\begin{aligned} (a - b + c)(x - y) &= a(x - y) - b(x - y) + c(x - y) \\ &= (ax - ay) - (bx - by) + (cx - cy) \\ &= ax - ay - bx + by + cx - cy. \end{aligned}$$

We may now state the general rule for multiplying together any two compound expressions.

Rule. Multiply each term of the first expression by each term of the second. When the terms multiplied together have like signs, prefix to the product the sign +, when unlike prefix -; the algebraical sum of the partial products so formed gives the complete product. This process is called **Distributing the Product**.

40. It should be noticed that the product of $a+b$ and $x-y$ is briefly expressed by $(a+b)(x-y)$, in which the brackets indicate that the expression $a+b$ taken as a whole is to be multiplied by the expression $x-y$ taken as a whole. By the above rule, the value of the product is the algebraical sum of the partial products $+ax$, $+bx$, $-ay$, $-by$; the sign of each product being determined by the rule of signs.

Example 1. Multiply $x+8$ by $x+7$.

$$\begin{aligned}\text{The product} &= (x+8)(x+7) \\ &= x^2 + 8x + 7x + 56 \\ &= x^2 + 15x + 56.\end{aligned}$$

The operation is more conveniently arranged as follows :

$$\begin{array}{r}x + 8 \\ x + 7 \\ \hline x^2 + 8x \\ + 7x + 56 \\ \hline\end{array}$$

by addition, $x^2 + 15x + 56$

We begin on the left and work to the right, placing the second result one place to the right, so that like terms may stand in the same vertical column.

Example 2. Multiply $2x-3y$ by $4x-7y$.

$$\begin{array}{r}2x - 3y \\ 4x - 7y \\ \hline 8x^2 - 12xy \\ - 14xy + 21y^2 \\ \hline\end{array}$$

by addition, $8x^2 - 26xy + 21y^2$.

EXAMPLES V. d.

Find the product of

- | | |
|---------------------------|--------------------------|
| 1. $x+5$ and $x+10$. | 2. $x+5$ and $x-5$. |
| 3. $x-7$ and $x-10$. | 4. $x-7$ and $x+10$. |
| 5. $x+7$ and $x-10$. | 6. $x+7$ and $x+10$. |
| 7. $x+6$ and $x-6$. | 8. $x+8$ and $x-4$. |
| 9. $x-12$ and $x-1$. | 10. $x+12$ and $x-1$. |
| 11. $x-15$ and $x+15$. | 12. $x-15$ and $-x+3$. |
| 13. $-x-2$ and $-x-3$. | 14. $-x+7$ and $x-7$. |
| 15. $-x+5$ and $-x-5$. | 16. $x-13$ and $x+14$. |
| 17. $x-17$ and $x+18$. | 18. $x+19$ and $x-20$. |
| 19. $-x-16$ and $-x+16$. | 20. $-x+21$ and $x-21$. |
| 21. $2x-3$ and $x+8$. | 22. $2x+3$ and $x-8$. |

v.]

Find th

23. $x-5$

25. $3x-6$

27. $5x-6$

29. $3x-5$

31. $a-2b$

33. $3a-6$

35. $x+a$

37. $x-2a$

39. $xy-ai$

41. We

Example

$3x^2 - 2x - 1$

$2x - 5$

$6x^3 - 4x^2 - 1$

$-15x^2 + 1$

$6x^3 - 19x^3$

Example 2

42. When
process of M
by the rules

Example, 1

$-b$ and $x-y$
the brackets
ole is to be
le. By the
ical sum of
ign of each

ows:

the left and
placing the
place to the
terms may
the vertical

Find the product of

- | | |
|---------------------------|-----------------------------|
| 23. $x-5$ and $2x-1$. | 24. $2x-5$ and $x-1$. |
| 25. $3x-5$ and $2x+7$. | 26. $3x+5$ and $2x-7$. |
| 27. $5x-6$ and $2x+3$. | 28. $5x+6$ and $2x-3$. |
| 29. $3x-5y$ and $3x+5y$. | 30. $3x-5y$ and $3x-5y$. |
| 31. $a-2b$ and $a+3b$. | 32. $a-7b$ and $a+8b$. |
| 33. $3a-6b$ and $a-8b$. | 34. $a-9b$ and $a+5b$. |
| 35. $x+a$ and $x-b$. | 36. $x-a$ and $x+b$. |
| 37. $x-2a$ and $x+3b$. | 38. $ax-by$ and $ax+by$. |
| 39. $xy-ab$ and $xy+ab$. | 40. $2pq-3r$ and $2pq+3r$. |

41. We shall now give a few examples of greater difficulty.

Example 1. Find the product of $3x^2-2x-5$ and $2x-5$.

$$\begin{array}{r} 3x^2 - 2x - 5 \\ 2x - 5 \\ \hline 6x^3 - 4x^2 - 10x \\ - 15x^2 + 10x + 25 \\ \hline 6x^3 - 19x^2 + 25 \end{array}$$

Each term of the first expression is multiplied by $2x$, the first term of the second expression; then each term of the first expression is multiplied by -5 ; like terms are placed in the same columns and the results added.

Example 2. Multiply $a-b+3c$ by $a+2b$.

$$\begin{array}{r} a - b + 3c \\ a + 2b \\ \hline a^2 - ab + 3ac \\ 2ab - 2b^2 + 6bc \\ \hline a^2 + ab + 3ac - 2b^2 + 6bc \end{array}$$

42. When the coefficients are fractional we use the ordinary process of Multiplication, combining the fractional coefficients by the rules of Arithmetic.

Example. Multiply $\frac{1}{3}a^2 - \frac{1}{2}ab + \frac{2}{3}b^2$ by $\frac{1}{2}a + \frac{1}{3}b$.

$$\begin{array}{r} \frac{1}{3}a^2 - \frac{1}{2}ab + \frac{2}{3}b^2 \\ \frac{1}{2}a + \frac{1}{3}b \\ \hline \frac{1}{6}a^3 - \frac{1}{4}a^2b + \frac{1}{3}ab^2 \\ + \frac{1}{6}a^2b - \frac{1}{6}ab^2 + \frac{2}{9}b^3 \\ \hline \frac{1}{6}a^3 - \frac{5}{36}a^2b + \frac{1}{6}ab^2 + \frac{2}{9}b^3 \end{array}$$

43. If the expressions are not arranged according to powers, ascending or descending, of some common letter, a rearrangement will be found convenient.

Example 1. Find the product of $2a^2 + 4b^2 - 3ab$ and $3ab - 5a^2 + 4b^2$.

$$\begin{array}{r}
 2a^2 - 3ab + 4b^2 \\
 - 5a^2 + 3ab + 4b^2 \\
 \hline
 -10a^4 + 15a^3b - 20a^2b^2 \\
 + 6a^3b - 9a^2b^2 + 12ab^3 \\
 \hline
 8a^2b^2 - 12ab^3 + 16b^4 \\
 -10a^4 + 21a^3b - 21a^2b^2 + 16b^4
 \end{array}$$

The rearrangement is not necessary, but convenient, because it makes the collection of like terms more easy.

Example 2. Multiply $2xz - z^2 + 2x^2 - 3yz + xy$ by $x - y + 2z$.

$$\begin{array}{r}
 2x^2 + xy + 2xz - 3yz - z^2 \\
 x - y + 2z \\
 \hline
 2x^3 + x^2y + 2x^2z - 3xyz - xz^2 \\
 - 2x^2y \quad - 2xyz \quad - xy^2 + 3y^2z + yz^2 \\
 \hline
 4x^2z + 2xyz + 4xz^2 \quad - 6yz^2 - 2z^3 \\
 2x^3 - x^2y + 6x^2z - 3xyz + 3xz^2 - xy^2 + 3y^2z - 5yz^2 - 2z^3
 \end{array}$$

EXAMPLES V. e.

Multiply together

- $a + b + c, a + b - c.$
- $a^2 - ab + b^2, a^2 + ab + b^2.$
- $x^3 - 2x^2 + 8, x + 2.$
- $x^2 + xy + y^2, x - y.$
- $16a^2 + 12ab + 9b^2, 4a - 3b.$
- $x^2 + x - 2, x^2 + x - 6.$
- $-a^5 + a^4b - a^3b^2, -a - b.$
- $a^3 + 2a^2b + 2ab^2, a^2 - 2ab + 2b^2.$
- $x^2 - 3xy - y^2, -x^2 + xy + y^2.$
- $x^2 - 2xy + y^2, x^2 + 2xy + y^2.$
- $ab + cd + ac + bd, ab + cd - ac - bd.$
- $-3a^2b^2 + 4ab^3 + 15a^3b, 5a^2b^2 + ab^3 - 3b^4.$
- $27x^3 - 36ax^2 + 48a^2x - 64a^3, 3x + 4a.$
- $a^2 - 5ab - b^2, a^2 + 5ab + b^2.$
- $x^2 - xy + x + y^2 + y + 1, x + y - 1.$
- $a - 2b + c, a + 2b - c.$
- $x^2 + 3y^2, x + 4y.$
- $x^4 - x^2y^2 + y^4, x^2 + y^2.$
- $a^2 - 2ax + 4x^2, a^2 + 2ax + 4x^2.$
- $a^3x - ax^2 + x^3 - a^3, x + a.$
- $2x^3 - 3x^2 + 2x, 2x^2 + 3x + 2.$
- $x^3 - 7x + 5, x^2 - 2x + 3.$
- $4x^2 + 6xy + 9y^2, 2x - 3y.$
- $b^3 - a^2b^2 + a^3, a^3 + a^2b^2 + b^3.$

v.]

Multiply

- $a^2 + b^2$
- $-x^3y$
- $x^{12} - x$
- $3a^2 + 2$
- $-ax^2 +$
- $-2x^3y$
- $\frac{1}{2}a^2 + \frac{1}{3}$
- $\frac{2}{3}x^2 + x$
- $\frac{1}{2}x^2 - \frac{2}{3}$
- $\frac{2}{3}ax + \frac{2}{3}$

44. Pro

result of m
 $x + 8$ and x
 explained, it
 soon learn to
 This is do
 the terms in
 from the cor
 binomials wh

In each of t

- The pro
- The first binomial expe
- The third two binomial e
- The mid numerical qua second terms of

Multiply together

$$25. a^2 + b^2 + c^2 - bc - ca - ab, a + b + c.$$

$$26. -x^2y + y^4 + x^2y^2 + x^4 - xy^3, x + y.$$

$$27. x^{12} - x^2y^2 + x^6y^4 - x^3y^6 + y^8, x^3 + y^2.$$

$$28. 3a^2 + 2a + 2a^3 + 1 + a^4, a^2 - 2a + 1.$$

$$29. -ax^2 + 3axy^2 - 9ay^4, -ax - 3ay^2.$$

$$30. -2x^3y + y^4 + 3x^2y^2 + x^4 - 2xy^3, x^2 + 2xy + y^2.$$

$$31. \frac{1}{2}a^2 + \frac{1}{3}a + \frac{1}{4}, \frac{1}{2}a - \frac{1}{3}.$$

$$32. \frac{1}{2}x^2 - 2x + \frac{3}{2}, \frac{1}{2}x + \frac{1}{3}.$$

$$33. \frac{2}{3}x^2 + xy + \frac{3}{2}y^2, \frac{1}{3}x - \frac{1}{2}y.$$

$$34. \frac{3}{2}x^2 - ax - \frac{2}{3}a^2, \frac{3}{4}x^2 - \frac{1}{2}ax + \frac{1}{3}a^2.$$

$$35. \frac{1}{2}x^2 - \frac{2}{3}x - \frac{3}{4}, \frac{1}{2}x^2 + \frac{2}{3}x - \frac{3}{4}.$$

$$36. \frac{2}{3}ax + \frac{2}{3}x^2 + \frac{1}{3}a^2, \frac{3}{4}a^2 + \frac{3}{2}x^2 - \frac{3}{2}ax.$$

44. **Products written down by inspection.** Although the result of multiplying together two binomial factors, such as $x+8$ and $x-7$, can always be obtained by the methods already explained, it is of the utmost importance that the student should soon learn to write down the product rapidly *by inspection*.

This is done by observing in what way the coefficients of the terms in the product arise, and noticing that they result from the combination of the numerical coefficients in the two binomials which are multiplied together; thus

$$(x+8)(x+7) = x^2 + 8x + 7x + 56 \\ = x^2 + 15x + 56.$$

$$(x-8)(x-7) = x^2 - 8x - 7x + 56 \\ = x^2 - 15x + 56.$$

$$(x+8)(x-7) = x^2 + 8x - 7x - 56 \\ = x^2 + x - 56.$$

$$(x-8)(x+7) = x^2 - 8x + 7x - 56 \\ = x^2 - x - 56.$$

In each of these results we notice that :

1. The product consists of three terms.
2. The first term is the product of the first terms of the two binomial expressions.
3. The third term is the product of the second terms of the two binomial expressions.
4. The middle term has for its coefficient the sum of the numerical quantities (taken with their proper signs) in the second terms of the two binomial expressions.

The intermediate step in the work may be omitted, and the products written down at once, as in the following examples :

$$(x+2)(x+3)=x^2+5x+6.$$

$$(x-3)(x+4)=x^2+x-12.$$

$$(x+6)(x-9)=x^2-3x-54.$$

$$(x-4y)(x-10y)=x^2-14xy+40y^2.$$

$$(x-6y)(x+4y)=x^2-2xy-24y^2.$$

By an easy extension of these principles we may write down the product of *any* two binomials.

$$\text{Thus} \quad (2x+3y)(x-y) = 2x^2+3xy-2xy-3y^2 \\ = 2x^2+xy-3y^2.$$

$$(3x-4y)(2x+y) = 6x^2-8xy+3xy-4y^2 \\ = 6x^2-5xy-4y^2.$$

$$(x+4)(x-4) = x^2+4x-4x-16 \\ = x^2-16.$$

$$(2x+5y)(2x-5y) = 4x^2+10xy-10xy-25y^2 \\ = 4x^2-25y^2.$$

EXAMPLES V. f.

Write down the values of the following products :

- | | | |
|-----------------------|-----------------------|-----------------------|
| 1. $(x+8)(x-5).$ | 2. $(x+6)(x-1).$ | 3. $(x-3)(x+10).$ |
| 4. $(x-1)(x+5).$ | 5. $(x+7)(x-9).$ | 6. $(x-10)(x-8).$ |
| 7. $(x-4)(x+11).$ | 8. $(x-2)(x+4).$ | 9. $(x+2)(x-2).$ |
| 10. $(a-1)(a+1).$ | 11. $(a+9)(a-5).$ | 12. $(a-3)(a+12).$ |
| 13. $(a-8)(a+4).$ | 14. $(a-8)(a+8).$ | 15. $(a-6)(a+13).$ |
| 16. $(a+3)(a+3).$ | 17. $(a-11)(a+11).$ | 18. $(a-8)(a-8).$ |
| 19. $(x-3a)(x+2a).$ | 20. $(x+6a)(x-5a).$ | 21. $(x+3a)(x-3a).$ |
| 22. $(x+4y)(x-2y).$ | 23. $(x+7y)(x-7y).$ | 24. $(x-3y)(x-3y).$ |
| 25. $(a+3b)(a+3b).$ | 26. $(a-5b)(a+10b).$ | 27. $(a-9b)(a-8b).$ |
| 28. $(2x-5)(x+2).$ | 29. $(2x-5)(x-2).$ | 30. $(2x+3)(x-3).$ |
| 31. $(3x-1)(x+1).$ | 32. $(2x+5)(2x-1).$ | 33. $(3x+7)(2x-3).$ |
| 34. $(4x-3)(2x+3).$ | 35. $(3x+8)(3x-8).$ | 36. $(2x-5)(2x-5).$ |
| 37. $(3x-2y)(3x+y).$ | 38. $(3x+2y)(3x+2y).$ | 39. $(2x+7y)(2x-5y).$ |
| 40. $(5x+3a)(5x-3a).$ | 41. $(2x-5a)(x+5a).$ | 42. $(2x+a)(2x+a).$ |

45. W
one letter
by using
coefficients
way, and t
the end of
must be ar
of the com
represent t

Example.

$$\begin{array}{r} 2-4+0- \\ 3+4-2 \\ 6-12+0- \\ 8-16+ \\ -4+ \\ \hline 6-4-20- \end{array}$$

Thus the

The meth
multiply tw
and contain

Example.

$$\begin{array}{r} 3+2+0+4+2 \\ 2+0-1 \\ 6+4+0+8+4 \\ -3-2-0 \\ \hline 6+4-3+6+4 \end{array}$$

The law o
the com

66

Note. Begin
coefficients unt
of multiplicatio

ted, and the
examples :

write down

The Method of Detached Coefficients.

45. When two compound expressions contain powers of one letter or y , the labour of multiplication may be lessened by using **detached coefficients**, that is, by writing down the coefficients only, multiplying them together in the ordinary way, and then inserting the successive powers of the letter at the end of the operation. In using this method the expressions must be arranged according to ascending or descending powers of the common letter, and zero coefficients must be used to represent terms corresponding to missing powers of that letter.

Example. Multiply $2x^3 - 4x^2 - 5$ by $3x^2 + 4x - 2$.

$$\begin{array}{r}
 2-4+0-5 \\
 3+4-2 \\
 \hline
 6-12+0-15 \\
 8-16+0-20 \\
 -4+8-0+10 \\
 \hline
 6-4-20-7-20+10
 \end{array}$$

Here we insert a zero coefficient to represent the power of x which is absent in the multiplicand. In the product the highest power of x is clearly x^5 , and the others follow in descending order.

Thus the product is

$$6x^5 - 4x^4 - 20x^3 - 7x^2 - 20x + 10.$$

The method of detached coefficients may also be used to multiply two compound expressions which are homogeneous and contain powers of two letters.

Example. Multiply $3a^4 + 2a^3b + 4ab^3 + 2b^4$ by $2a^2 - b^2$.

$$\begin{array}{r}
 3+2+0+4+2 \\
 2+0-1 \\
 \hline
 6+4+0+8+4 \\
 -3-2-0-4-2 \\
 \hline
 6+4-3+6+4-4-2
 \end{array}$$

We write a zero coefficient to represent the term containing a^2b^2 which is absent in the first expression. Similarly, the term containing ab is represented by a zero coefficient in the second expression.

The law of formation of the literal factors is easily seen, and the complete product is

$$6a^6 + 4a^5b - 3a^4b^2 + 6a^3b^3 + 4a^2b^4 - 4ab^5 - 2b^6.$$

Note. Beginners should on no account attempt to use detached coefficients until they are well practised in the ordinary full process of multiplication.

CHAPTER VI.

DIVISION.

46. WHEN a quantity a is divided by the quantity b , the **quotient** is defined to be that which when multiplied by b produces a . This operation of division is denoted by $a \div b$, $\frac{a}{b}$, or a/b ; in each of these modes of expression a is called the **dividend**, and b the **divisor**.

Division is thus the inverse of multiplication, and

$$(a \div b) \times b = a.$$

This statement may also be expressed verbally as follows :
quotient $\times$ divisor = dividend.

Since Division is the inverse of Multiplication, it follows that the Laws of Commutation, Association, and Distribution, which have been established for Multiplication, hold for Division.

47. *The Rule of Signs holds for division.*

Thus

$$ab \div a = \frac{ab}{a} = \frac{a \times b}{a} = b.$$

$$-ab \div a = \frac{-ab}{a} = \frac{a \times (-b)}{a} = -b.$$

$$ab \div (-a) = \frac{ab}{-a} = \frac{(-a) \times (-b)}{-a} = -b.$$

$$-ab \div (-a) = \frac{-ab}{-a} = \frac{(-a) \times b}{-a} = b.$$

Hence in division as well as multiplication

like signs produce +,

unlike signs produce -.

48. T

Example
who
or o

Example

The quotie

Theref

Example

We see,
is the differ
divisor. T

The rule

Rule. 7

*subtracting
the dividend*

*To the re
of the coeffic*

Example 4

Example 5

Note. If v
same power o

Thus, by

but al

This result
full significan
Indices.

Division of Simple Expressions.

48. The method is shewn in the following examples :

Example 1. Since the product of 4 and x is $4x$, it follows that when $4x$ is divided by x the quotient is 4, or otherwise,

$$4x \div x = 4.$$

Example 2. Divide $27a^5$ by $9a^3$.

$$\begin{array}{l} \text{The quotient} = \frac{27a^5}{9a^3} = \frac{27aaaa}{9aaa} \\ \qquad \qquad \qquad = 3aa = 3a^2. \end{array}$$

We remove from the divisor and dividend the factors common to both, just as in arithmetic.

Therefore

$$27a^5 \div 9a^3 = 3a^2.$$

Example 3. Divide $35a^3b^2c^3$ by $7ab^2c^2$.

$$\text{The quotient} = \frac{35aaa \cdot bb \cdot ccc}{7a \cdot bb \cdot cc} = 5aa \cdot c = 5a^2c.$$

We see, in each case, that the index of any letter in the quotient is the difference of the indices of that letter in the dividend and divisor. This is called the **Index Law for Division**.

The rule may now be stated :

Rule. The index of each letter in the quotient is obtained by subtracting the index of that letter in the divisor from that in the dividend.

To the result so obtained prefix with its proper sign the quotient of the coefficient of the dividend by that of the divisor.

Example 4. Divide $45a^6b^2x^4$ by $-9a^3bx^2$.

$$\begin{aligned} \text{The quotient} &= (-5) \times a^{6-3}b^{2-1}x^{4-2} \\ &= -5a^3bx^2. \end{aligned}$$

Example 5. $-21a^2b^3 \div (-7a^2b^3) = 3b$.

Note. If we apply the rule to divide any power of a letter by the same power of the letter we are led to a curious conclusion.

Thus, by the rule $a^3 \div a^3 = a^{3-3} = a^0$;

but also $a^3 \div a^3 = \frac{a^3}{a^3} = 1$.

$$\therefore a^0 = 1.$$

This result will appear somewhat strange to the beginner, but its full significance will be explained in the chapter on the Theory of Indices.

Division of a Compound Expression by a Simple Expression.

49. Rule. To divide a compound expression by a single factor, divide each term separately by that factor, and take the algebraic sum of the partial quotients so obtained.

This follows at once from Art. 33.

- Examples.* (1) $(9x - 12y + 3z) \div -3 = -3x + 4y - z$.
 (2) $(36a^3b^2 - 24a^2b^5 - 20a^4b^2) \div 4a^2b = 9ab - 6b^4 - 5a^2b$.
 (3) $(2x^2 - 5xy + \frac{3}{2}x^2y^3) \div -\frac{1}{2}x = -4x + 10y - 3xy^3$.

EXAMPLES VI. a.

Divide

- | | |
|---|---|
| 1. $3x^3$ by x^2 . | 2. $27x^4$ by $-9x^3$. |
| 3. $-35x^6$ by $7x^3$. | 4. abx^2 by $-ax$. |
| 5. x^3y^3 by x^2y . | 6. a^4x^3 by $-a^2x^3$. |
| 7. $4a^2b^2c^3$ by ab^2c^2 . | 8. $12a^6b^4c^6$ by $-3a^4b^2c$. |
| 9. $-a^5c^9$ by $-ac^3$. | 10. $15x^5y^7z^4$ by $5x^2y^2z^2$. |
| 11. $-16x^3y^2$ by $-4xy^2$. | 12. $-48a^9$ by $-8a^3$. |
| 13. $35a^{11}$ by $7a^7$. | 14. $63a^7b^3c^3$ by $9a^5b^5c^3$. |
| 15. $7a^2bc$ by $-7a^2bc$. | 16. $28a^4b^3$ by $-4a^3b$. |
| 17. $16b^2yx^2$ by $-2xy$. | 18. $-50y^3x^3$ by $-5x^3y$. |
| 19. $x^2 - 2xy$ by x . | 20. $x^3 - 3x^2 + x$ by x . |
| 21. $x^6 - 7x^5 + 4x^4$ by x^2 . | 22. $10x^7 - 8x^6 + 3x^4$ by x^3 . |
| 23. $15x^5 - 25x^4$ by $-5x^3$. | 24. $27x^6 - 36x^5$ by $9x^5$. |
| 25. $-24x^6 - 32x^4$ by $-8x^3$. | 26. $34x^2y^2 - 51x^2y^3$ by $17xy$. |
| 27. $a^3 - ab - ac$ by $-a$. | 28. $a^3 - a^2b - a^2b^2$ by a^2 . |
| 29. $3x^3 - 9x^2y - 12xy^2$ by $-3x$. | 30. $4x^4y^4 - 8x^3y^2 + 6xy^3$ by $-2xy$. |
| 31. $-3a^2 + \frac{9}{2}ab - 6ac$ by $-\frac{3}{2}a$. | 32. $\frac{1}{2}x^2y^2 - 3x^3y^4$ by $-\frac{3}{2}x^3y^2$. |
| 33. $-\frac{5}{2}x^2 + \frac{5}{3}xy + \frac{1}{3}0x$ by $-\frac{5}{6}x$. | 34. $-2a^5x^3 + \frac{7}{2}a^4x^4$ by $\frac{7}{3}a^3x$. |
| 35. $\frac{1}{4}a^2x - \frac{1}{16}abx - \frac{3}{8}acx$ by $\frac{3}{8}ax$. | |

50. Rule.

descending

2. Divide the left of

3. Multiply product u

4. Subtract as may be

Repeat brought d

Example

Arrang

divide x^2 ,
divisor; th
put the pro

by su

On rep
term

The en

The reas
into as ma
quotient is
Thus $x^2 + 11$
namely $x^2 +$
thus we obt

Example 2

Simple

Division of Compound Expressions.

50. To divide one compound expression by another.

Rule. 1. Arrange divisor and dividend in ascending or descending powers of some common letter.

2. Divide the term on the left of the dividend by the term on the left of the divisor, and put the result in the quotient.

3. Multiply the **WHOLE** divisor by this quotient, and put the product under the dividend.

4. Subtract and bring down from the dividend as many terms as may be necessary.

Repeat these operations till all the terms from the dividend are brought down.

Example 1. Divide $x^2 + 11x + 30$ by $x + 6$.

Arrange the work thus :

$$(x + 6) x^2 + 11x + 30$$

divide x^2 , the first term of the dividend, by x , the first term of the divisor; the quotient is x . Multiply the *whole* divisor by x , and put the product $x^2 + 6x$ under the dividend. We then have

$$\begin{array}{r} (x + 6) x^2 + 11x + 30 \\ x^2 + 6x \end{array}$$

by subtraction

$$5x + 30.$$

On repeating the process above explained we find that the next term in the quotient is $+5$.

The entire operation is more compactly written as follows :

$$\begin{array}{r} (x + 6) x^2 + 11x + 30 \\ x^2 + 6x \quad 5x + 30 \\ \hline 5x + 30 \end{array}$$

The reason for the rule is thus: the dividend may be divided into as many parts as may be convenient, and the complete quotient is found by taking the sum of all the partial quotients. Thus $x^2 + 11x + 30$ is divided by the above process into two parts, namely $x^2 + 6x$, and $5x + 30$, and each of these is divided by $x + 6$; thus we obtain the complete quotient $x + 5$.

Example 2. Divide $24x^2 - 65xy + 21y^2$ by $8x - 3y$.

$$\begin{array}{r} (8x - 3y) 24x^2 - 65xy + 21y^2 \\ 24x^2 - 9xy \quad -56xy + 21y^2 \\ \hline -56xy + 21y^2 \end{array}$$

EXAMPLES VI. b.

Divide

1. x^2+3x+2 by $x+1$.
2. $x^2-7x+12$ by $x-3$.
3. $a^2-11a+30$ by $a-5$.
4. $a^2-49a+600$ by $a-25$.
5. $3x^2+10x+3$ by $x+3$.
6. $2x^2+11x+5$ by $2x+1$.
7. $5x^2+11x+2$ by $x+2$.
8. $2x^2+17x+21$ by $2x+3$.
9. $5x^2+16x+3$ by $x+3$.
10. $3x^2+34x+11$ by $3x+1$.
11. $4x^2+23x+15$ by $4x+3$.
12. $6x^2-7x-3$ by $2x-3$.
13. $3x^2+x-14$ by $x-2$.
14. $3x^2-x-14$ by $x+2$.
15. $6x^2-31x+35$ by $2x-7$.
16. $4x^2+x-14$ by $x+2$.
17. $12a^2-7ax-12x^2$ by $3a-4x$.
18. $15a^2+17ax-4x^2$ by $3a+4x$.
19. $12a^2-11ac-36c^2$ by $4a-9c$.
20. $9a^2+6ac-35c^2$ by $3a+7c$.
21. $60x^2-4xy-45y^2$ by $10x-9y$.
22. $96x^2-15y^2-4xy$ by $12x-5y$.
23. $7x^3+96x^2-28x$ by $7x-2$.
24. $100x^3-3x-13x^2$ by $3+25x$.
25. $27x^3+9x^2-3x-10$ by $3x-2$.
26. $16a^3-46a^2+39a-9$ by $8a-3$.
27. $15+3a-7a^2-4a^3$ by $5-4a$.
28. $16-96x+216x^2-216x^3+81x^4$ by $2-3x$.

51. The process of Art. 50 is applicable to cases in which the divisor consists of more than two terms.

Example. Divide $6x^5-x^4+4x^3-5x^2-x-15$ by $2x^2-x+3$.

$$\begin{array}{r} 2x^2-x+3 \overline{) 6x^5-x^4+4x^3-5x^2-x-15} \\ 6x^5-3x^4+9x^3 \\ \hline \end{array}$$

$$2x^4-5x^3-5x^2$$

$$2x^4-x^3+3x^2$$

$$-4x^3-8x^2-x$$

$$-4x^3+2x^2-6x$$

$$-10x^2+5x-15$$

$$-10x^2+5x-15$$

52. Sometimes it will be found convenient to arrange the expressions in *ascending* powers of some common letter.

Example. Divide $2a^3+10-16a-39a^2+15a^4$ by $2-4a-5a^2$.

$$\begin{array}{r} 2-4a-5a^2 \overline{) 10-16a-39a^2+2a^3+15a^4} \\ 10-20a-25a^2 \\ \hline \end{array}$$

$$4a-14a^2+2a^3$$

$$4a-8a^2-10a^3$$

$$-6a^2+12a^3+15a^4$$

$$-6a^2+12a^3+15a^4$$

53. V

*Example**Example*

a

Note. In
mainders ar
The resul

54. Wh
may still be

Example.

In the ex
contained in
work shoul
dimension

53. We add a few harder cases worked out in full.

Example 1. Divide $x^4 + 4a^4$ by $x^2 + 2xa + 2a^2$.

$$\begin{array}{r}
 x^4 + 4a^4 \\
 x^2 + 2xa + 2a^2 \overline{) x^4 + 4a^4} \quad (x^2 - 2xa + 2a^2 \\
 \underline{x^4 + 2x^3a + 2x^2a^2} \\
 - 2x^3a - 2x^2a^2 \\
 \underline{- 2x^3a - 4x^2a^2 - 4xa^3} \\
 2x^2a^2 + 4xa^3 + 4a^4 \\
 \underline{2x^2a^2 + 4xa^3 + 4a^4} \\
 0
 \end{array}$$

Example 2. Divide $a^3 + b^3 + c^3 - 3abc$ by $a + b + c$.

$$\begin{array}{r}
 a + b + c \overline{) a^3 + b^3 + c^3 - 3abc} \\
 \underline{a^3 + a^2b + a^2c} \\
 - a^2b - a^2c - 3abc \\
 \underline{- a^2b - ab^2 - abc} \\
 - a^2c + ab^2 - 2abc \\
 \underline{- a^2c - abc - ac^2} \\
 ab^2 - abc + ac^2 + b^3 \\
 \underline{ab^2 + b^3 + b^2c} \\
 - abc + ac^2 - b^2c \\
 \underline{- abc - b^2c - bc^2} \\
 ac^2 + bc^2 + c^3 \\
 \underline{ac^2 + bc^2 + c^3} \\
 0
 \end{array}$$

Note. In the above example the dividend and successive remainders are arranged in *descending* powers of a .

The result of this important division will be referred to later.

54. When the coefficients are fractional the ordinary process may still be employed.

Example. Divide $\frac{1}{4}x^3 + \frac{1}{7}xy^2 + \frac{1}{12}y^3$ by $\frac{1}{2}x + \frac{1}{3}y$.

$$\begin{array}{r}
 \frac{1}{2}x + \frac{1}{3}y \overline{) \frac{1}{4}x^3 + \frac{1}{7}xy^2 + \frac{1}{12}y^3} \quad (\frac{1}{2}x^2 - \frac{1}{3}xy + \frac{1}{4}y^2 \\
 \underline{\frac{1}{4}x^3 + \frac{1}{8}x^2y} \phantom{+ \frac{1}{7}xy^2 + \frac{1}{12}y^3} \\
 - \frac{1}{8}x^2y + \frac{1}{7}xy^2 \phantom{+ \frac{1}{12}y^3} \\
 \underline{- \frac{1}{8}x^2y - \frac{1}{9}xy^2} \phantom{+ \frac{1}{12}y^3} \\
 \frac{1}{8}xy^2 + \frac{1}{12}y^3 \\
 \underline{\frac{1}{8}xy^2 + \frac{1}{12}y^3} \\
 0
 \end{array}$$

In the examples given hitherto the divisor has been exactly contained in the dividend. When the division is not exact the work should be carried on until the remainder is of lower dimensions [Art. 24] than the divisor.

EXAMPLES VI. c.

Divide

1. $x^3 - x^2 - 9x - 12$ by $x^2 + 3x + 3$.
2. $2y^3 - 3y^2 - 6y - 1$ by $2y^2 - 5y - 1$.
3. $6m^3 - m^2 - 14m + 3$ by $3m^2 + 4m - 1$.
4. $6a^5 - 13a^4 + 4a^3 + 3a^2$ by $3a^3 - 2a^2 - a$.
5. $x^4 + x^3 + 7x^2 - 6x + 8$ by $x^2 + 2x + 8$.
6. $a^4 - a^3 - 8a^2 + 12a - 9$ by $a^2 + 2a - 3$.
7. $a^4 + 6a^3 + 13a^2 + 12a + 4$ by $a^2 + 3a + 2$.
8. $2x^4 - x^3 + 4x^2 + 7x + 1$ by $x^2 - x + 3$.
9. $x^5 - 5x^4 + 9x^3 - 6x^2 - x + 2$ by $x^3 - 3x + 2$.
10. $x^5 - 4x^4 + 3x^3 + 3x^2 - 3x + 2$ by $x^2 - x - 2$.
11. $30x^4 + 11x^3 - 82x^2 - 5x + 3$ by $2x - 4 + 3x^2$.
12. $30y + 9 - 71y^3 + 28y^4 - 35y^2$ by $4y^2 - 13y + 6$.
13. $6k^5 - 15k^4 + 4k^3 + 7k^2 - 7k + 2$ by $3k^3 - k + 1$.
14. $15 + 2m^4 - 31m + 9m^2 + 4m^3 + m^5$ by $3 - 2m - m^2$.
15. $2x^3 - 8x + x^4 + 12 - 7x^2$ by $x^2 + 2 - 3x$.
16. $x^5 - 2x^4 - 4x^3 + 19x^2$ by $x^3 - 7x + 5$.
17. $192 - x^4 + 128x + 4x^2 - 8x^3$ by $16 - x^2$.
18. $14x^4 + 45x^3y + 78x^2y^2 + 45xy^3 + 14y^4$ by $2x^2 + 5xy + 7y^2$.
19. $x^5 - x^4y + x^3y^2 - x^2 + x^2 - y^3$ by $x^3 - x - y$.
20. $x^5 + x^4y - x^3y^2 + x^3 - 2xy^2 + y^3$ by $x^2 + xy - y^2$.
21. $a^6 - b^6$ by $a^3 - b^3$.
22. $x^9 - y^9$ by $x^2 + xy + y^2$.
23. $x^7 - 2y^{14} - 7x^5y^4 - 7xy^{12} + 14x^3y^8$ by $x - 2y^2$.
24. $a^3 + 3a^2b + b^3 - 1 + 3ab^2$ by $a + b - 1$.
25. $x^3 - y^3$ by $x^3 + x^2y + xy^2 + y^3$.
26. $a^{12} - b^{12}$ by $a^3 - b^3$.
27. $a^{12} + 2a^6b^6 + b^{12}$ by $a^4 + 2a^2b^2 + b^4$.
28. $1 - a^3 - 8x^3 - 6ax$ by $1 - a - 2x$.

Find the quotient of

29. $\frac{1}{8}a^3 - \frac{9}{4}a^2x + \frac{2}{3}ax^2 - 27x^3$ by $\frac{1}{2}a - 3x$.
30. $\frac{1}{27}a^3 - \frac{1}{12}a^2 + \frac{1}{16}a - \frac{1}{64}$ by $\frac{1}{3}a - \frac{1}{4}$.
31. $\frac{3}{4}a^2c^3 + \frac{1}{12}a^5$ by $\frac{1}{5}a^2 + \frac{1}{2}ac$.
32. $\frac{9}{16}a^4 - \frac{3}{4}a^3 - \frac{7}{4}a^2 + \frac{4}{3}a + \frac{1}{6}$ by $\frac{3}{2}a^2 - \frac{8}{3} - a$.
33. $36x^2 + \frac{1}{6}y^2 + \frac{1}{4} - 4xy - 6x + \frac{1}{3}y$ by $6x - \frac{1}{3}y - \frac{1}{3}$.
34. $\frac{8}{27}a^5 - \frac{2}{512}ax^4$ by $\frac{2}{3}a - \frac{3}{4}x$.

[Examples 6-18, 20 will furnish practice in the use of Detached Coefficients as explained in Art. 45.]

55. they are

and so on
all positive

and so on
alternately
always odd

and so on
alternately
always even

the index
by $x + y$

All the
follows:

- (1) x^n
 - (2) x^n
 - (3) x^n
 - (4) x^n
- even whole

55. The following examples in division may be easily verified; they are of great importance, and should be carefully noticed.

$$\text{I. } \begin{cases} \frac{x^2-y^2}{x-y} = x+y, \\ \frac{x^3-y^3}{x-y} = x^2+xy+y^2, \\ \frac{x^4-y^4}{x-y} = x^3+x^2y+xy^2+y^3, \end{cases}$$

and so on; the divisor being $x-y$, the terms in the quotient *all positive*, and the index in the dividend *either odd or even*.

$$\text{II. } \begin{cases} \frac{x^3+y^3}{x+y} = x^2-xy+y^2, \\ \frac{x^5+y^5}{x+y} = x^4-x^3y+x^2y^2-xy^3+y^4, \\ \frac{x^7+y^7}{x+y} = x^6-x^5y+x^4y^2-x^3y^3+x^2y^4-xy^5+y^6, \end{cases}$$

and so on; the divisor being $x+y$, the terms in the quotient *alternately positive and negative*, and the index in the dividend *always odd*.

$$\begin{cases} \frac{x^2-y^2}{x+y} = x-y, \\ \frac{x^4-y^4}{x+y} = x^3-x^2y+xy^2-y^3, \\ \frac{x^6-y^6}{x+y} = x^5-x^4y+x^3y^2-x^2y^3+xy^4-y^5, \end{cases}$$

and so on; the divisor being $x+y$, the terms in the quotient *alternately positive and negative*, and the index in the dividend *always even*.

IV. The expressions x^2+y^2 , x^4+y^4 , x^6+y^6 , ... (where the index is *even*, and the terms *both positive*) are *never* divisible by $x+y$ or $x-y$.

All these different cases may be more concisely stated as follows:

- (1) x^n-y^n is divisible by $x-y$ if n be any whole number.
- (2) x^n+y^n is divisible by $x+y$ if n be any *odd* whole number.
- (3) x^n-y^n is divisible by $x+y$ if n be any *even* whole number.
- (4) x^n+y^n is never divisible by $x+y$ or $x-y$, when n is an *even* whole number.

CHAPTER VII.

REMOVAL AND INSERTION OF BRACKETS.

56. QUANTITIES are enclosed within brackets to indicate that they must all be operated upon in the same way. Thus in the expression $2a - 3b - (4a - 2b)$ the brackets indicate that the expression $4a - 2b$ treated as a whole has to be subtracted from $2a - 3b$. When we wish to enclose within brackets part of an expression already enclosed within brackets it is usual to employ brackets of different forms. The brackets in common use are $()$, $\{\}$, $[\]$. Sometimes a line called a **vinculum** is drawn over the symbols to be connected; thus $a - \overline{b+c}$ is used with the same meaning as $a - (b+c)$, and hence $a - \overline{b+c} = a - b - c$.

Removal of Brackets.

57. To remove brackets it is usually best to begin with the inside pair, and in dealing with each pair in succession we apply the rules already given in Arts. 21, 22.

Example 1. Simplify, by removing brackets, the expression
 $a - 2b - [4a - 6b - \{3a - c + (5a - 2b - \overline{3a - c + 2b})\}]$.

Removing the brackets one by one, we have

$$\begin{aligned} & a - 2b - [4a - 6b - \{3a - c + (5a - 2b - 3a + c - 2b)\}] \\ &= a - 2b - [4a - 6b - \{3a - c + 5a - 2b - 3a + c - 2b\}] \\ &= a - 2b - [4a - 6b - 3a + c - 5a + 2b + 3a - c + 2b] \\ &= a - 2b - 4a + 6b + 3a - c + 5a - 2b - 3a + c - 2b \\ &= 2a, \text{ by collecting like terms.} \end{aligned}$$

Example 2. Simplify the expression

$$-[-2x - \{3y - (2x - 3y) + (3x - 2y)\} + 2x].$$

$$\begin{aligned} \text{The expression} &= -[-2x - \{3y - 2x + 3y + 3x - 2y\} + 2x] \\ &= -[-2x - 3y + 2x - 3y - 3x + 2y + 2x] \\ &= 2x + 3y - 2x + 3y + 3x - 2y - 2x \\ &= x + 4y. \end{aligned}$$

Simpli

1. $a -$
2. $a -$
3. $a -$
4. $\{a -$
5. $2a -$
6. $- \{ -$
7. $- [a -$
8. $- (-$
9. $- [-$
10. $- 5x$
11. $- (-$
12. $3 -$
13. $- 2a$
14. $3x -$
15. $- [5x$
16. $- [15$
17. $8x - \{$
18. $- [x -$
19. $- [a +$
20. $- [a -$

58. A
every term
plied by th

Note. T

fraction is

Again, an
the line ab
root of the

Thus

whereas

EXAMPLES VII. a.

Simplify by removing brackets

1. $a - (b - c) \div a + (b - c) + b - (c + a)$.
2. $a - [b + \{a - (b + a)\}]$.
3. $a - [2a - \{3b - (4c - 2a)\}]$.
4. $\{a - (b - c)\} + \{b - (c - a)\} - \{c - (a - b)\}$.
5. $2a - (5b + [3c - a]) - (5a - [b + c])$.
6. $-\{-[-(a - \overline{b - c})]\}$.
7. $-[a - \{b - (c - a)\}] - [b - \{c - (a - b)\}]$.
8. $-(-(-(-x))) - (-(-y))$.
9. $-[-\{-(b + c - a)\}] + [-\{-(c + a - b)\}]$.
10. $-5x - [3y - \{2x - (2y - x)\}]$.
11. $-(-(-a)) - (-(-(-x)))$.
12. $3a - [a + b - \{a + b + c - (a + b + c + d)\}]$.
13. $-2a - [3x + \{3c - (4y + 3x + 2a)\}]$.
14. $3x - [5y - \{6z - (4x - 7y)\}]$.
15. $-[5x - (11y - 3x)] - [5y - (3x - 6y)]$.
16. $-[15x - \{14y - (15z + 12y) - (10x - 15z)\}]$.
17. $8x - \{16y - [3x - (12y - x) - 8y] + x\}$.
18. $-[x - \{z + (x - z) - (z - x) - z\} - x]$.
19. $-[a + \{a - (a - x) - (a + x) - a\} - a]$.
20. $-[a - \{a + (x - a) - (x - a) - a\} - 2a]$.

58. A coefficient placed before any bracket indicates that every term of the expression within the bracket is to be multiplied by that coefficient.

Note. The line between the numerator and denominator of a fraction is a kind of vinculum. Thus $\frac{x-5}{3}$ is equivalent to $\frac{1}{3}(x+5)$.

Again, an expression of the form $\sqrt{(x+y)}$ is often written $\sqrt{x+y}$, the line above being regarded as a vinculum indicating the square root of the compound expression $x+y$ taken as a whole.

Thus $\sqrt{25+144} = \sqrt{169} = 13$,
 whereas $\sqrt{25} + \sqrt{144} = 5 + 12 = 17$.

59. Sometimes it is advisable to simplify in the course of the work.

Example. Find the value of

$$84 - 7[-11x - 4\{-17x + 3(8 - 9 - 5x)\}].$$

$$\begin{aligned} \text{The expression} &= 84 - 7[-11x - 4\{-17x + 3(8 - 9 + 5x)\}] \\ &= 84 - 7[-11x - 4\{-17x + 3(5x - 1)\}] \\ &= 84 - 7[-11x - 4\{-17x + 15x - 3\}] \\ &= 84 - 7[-11x - 4\{-2x - 3\}] \\ &= 84 - 7[-11x + 8x + 12] \\ &= 84 - 7[-3x + 12] \\ &= 84 + 21x - 84 \\ &= 21x. \end{aligned}$$

When the beginner has had a little practice the number of steps may be considerably diminished.

EXAMPLES VII. b.

Simplify by removing brackets

1. $a - [2b + \{3c - 3a - (a + b)\} + 2a - (b + 3c)].$
2. $a + b - (c + a - [b + c - (a + b - \{c + a - (b + c - a)\})]).$
3. $a - (b - c) - [a - b - c - 2\{b + c - 3(c - a) - d\}].$
4. $2x - (3y - 4z) - \{2x - (3y + 4z)\} - \{3y - (4z + 2x)\}.$
5. $b + c - (a + b - [c + a - (b + c - \{a + b - (c + a - b)\})]).$
6. $3b - \{5a - [6a + 2(10a - b)]\}.$
7. $a - (b - c) - [a - b - c - 2\{b + c\}].$
8. $3a^2 - [6a^2 - \{8b^2 - (9c^2 - 2a^2)\}].$
9. $b - (c - a) - [b - a - c - 2\{c + a - 3(a - b) - d\}].$
10. $-20(a - d) + 3(b - c) - 2[b + c + d - 3\{c + d - 4(d - a)\}].$
11. $-4(a + d) + 24(b - c) - 2[c + d + a - 3\{d + a - 4(b + c)\}].$
12. $-10(a + b) - [c + a - 3\{a + 2b - (c + a - b)\}] + 4c.$
13. $a - 2(b - c) - [-\{ - (4a - b - c - 2\{a + b + c\})\}].$
14. $8(b - c) - [-\{a - b - 3(c - b + a)\}].$
15. $2(3b - 5a) - 7[a - 6\{2 - 5(a - b)\}].$
16. $6\{a - 2[b - 3(c + d)]\} - 4\{a - 3[b - 4(c - d)]\}.$
17. $5\{a - 2[a - 2(a + x)]\} - 4\{a - 2[a - 2(a + x)]\}.$
18. $-10\{a - 6[a - (b - c)]\} + 60\{b - (c + a)\}.$

19. $-3\{-$
20. $-2\{-$
21. $\frac{1}{4}(a -$
22. $35\left[\frac{3a}{2}\right]$
23. $\frac{3}{8}\left\{\frac{4}{3}(a\right.$
24. $\frac{1}{2}x - \frac{1}{2}$

60. The important Arts. 21, 22

(1) Any and the sign brackets removed

(2) Any brackets and term within

Examples.

61. The two ways.

Example. The may be written

or
or

62. Whenever a bracket, it may be of the expression

E.A.

19. $-3\{-2[-4(-a)]\} + 5\{-2[-2(-a)]\}.$
 20. $-2\{-[-(x-y)]\} + \{-2[-(x-y)]\}.$
 21. $\frac{1}{4}\{a-5(b-a)\} - \frac{3}{2}\left\{\frac{1}{3}\left(b-\frac{a}{3}\right) - \frac{2}{9}\left[a-\frac{3}{4}\left(b-\frac{4a}{5}\right)\right]\right\}.$
 22. $35\left[\frac{3x-4y}{5} - \frac{1}{10}\left\{3x-\frac{5}{7}(7x-4y)\right\}\right] + 8(y-2x).$
 23. $\frac{3}{8}\left\{\frac{4}{3}(a-b)-8(b-c)\right\} - \left\{\frac{b-c}{2} - \frac{c-a}{3}\right\} - \frac{1}{2}\left\{c-a-\frac{2}{3}(a-b)\right\}.$
 24. $\frac{1}{2}x - \frac{1}{2}\left(\frac{2}{3}y - \frac{1}{2}z\right) - \left[x - \left\{\frac{1}{2}x - \left(\frac{1}{3}y - \frac{1}{4}z\right)\right\} - \left(\frac{2}{3}y - \frac{1}{2}z\right)\right].$

Insertion of Brackets.

60. The converse operation of inserting brackets is important. The rules for doing this have been enunciated in Arts. 21, 22; for convenience we repeat them.

(1) *Any part of an expression may be enclosed within brackets and the sign + prefixed, the sign of every term within the brackets remaining unaltered.*

(2) *Any part of an expression may be enclosed within brackets and the sign - prefixed, provided the sign of every term within the brackets be changed.*

Examples. $a-b+c-d-e=a-b+(c-d-e).$

$$a-b+c-d-e=a-(b-c)-(d+e).$$

$$x^2-ax+bx-ab=(x^2-ax)+(bx-ab).$$

$$xy-ax-by+ab=(xy-by)-(ax-ab).$$

61. The terms of an expression can be bracketed in various ways.

Example. The expression $ax-bx+cx-ay+by-cy$ may be written
 (ax-bx)+(cx-ay)+(by-cy),
 or
 (ax-bx+cx)-(ay-by+cy),
 or
 (ax-ay)-(bx-by)+(cx-cy).

62. Whenever a factor is common to every term within a bracket, it may be removed and placed outside as a multiplier of the expression within the bracket.

E.A.

D

Example 1. In the expression

$$ax^3 - cx + 7 - dx^2 + bx - c - dx^3 + bx^2 - 2x$$

bracket together the powers of x so as to have the sign + before each bracket.

$$\begin{aligned}\text{The expression} &= (ax^3 - dx^3) + (bx^2 - dx^2) + (bx - cx - 2x) + (7 - c) \\ &= x^3(a - d) + x^2(b - d) + x(b - c - 2) + (7 - c) \\ &= (a - d)x^3 + (b - d)x^2 + (b - c - 2)x + 7 - c.\end{aligned}$$

In this last result the compound expressions $a - d$, $b - d$, $b - c - 2$ are regarded as the coefficients of x^3 , x^2 , and x respectively.

Example 2. In the expression $-a^2x - 7a + a^2y + 3 - 2x - ab$ bracket together the powers of a so as to have the sign - before each bracket.

$$\begin{aligned}\text{The expression} &= -(a^2x - a^2y) - (7a + ab) - (2x - 3) \\ &= -a^2(x - y) - a(7 + b) - (2x - 3) \\ &= -(x - y)a^2 - (7 + b)a - (2x - 3).\end{aligned}$$

EXAMPLES VII. c.

In the following expressions bracket the powers of x so that the signs before all the brackets shall be positive :

1. $ax^4 + bx^2 + 5 + 2bx - 5x^2 + 2x^4 - 3x$.
2. $3bx^2 - 7 - 2x + ab + 5ax^3 + cx - 4x^2 - bx^3$.
3. $2 - 7x^3 + 5ax^2 - 2cx + 9ax^3 + 7x - 3x^2$.
4. $2cx^5 - 3abx + 4dx - 3bx^4 - a^2x^5 + x^4$.

In the following expressions bracket the powers of x so that the signs before all the brackets shall be negative :

5. $ax^2 + 5x^3 - a^2x^4 - 2bx^3 - 3x^2 - bx^4$.
6. $7x^3 - 3c^2x - abx^5 + 5ax + 7x^5 - abcx^3$.
7. $ax^2 + a^2x^3 - bx^2 - 5x^3 - cx^3$.
8. $3b^2x^4 - bx - ax^4 - cx^4 - 5c^2x - 7x^4$.

Simplify the following expressions, and in each result re-group the terms according to powers of x :

9. $ax^3 - 2cx - [bx^2 - \{cx - dx - (bx^3 + 3cx^2)\} - (cx^2 - bx)]$.
10. $5ax^3 - 7(bx - cx^2) - \{6bx^2 - (3ax^2 + 2ax) - 4cx^3\}$.
11. $ax^2 - 3\{-ax^3 + 3bx - 4[\frac{1}{8}cx^3 - \frac{2}{3}(ax - bx^2)]\}$.
12. $x^5 - 4bx^4 - \frac{1}{6}\left[12ax - 4\left\{3bx^4 - 9\left(\frac{cx}{2} - bx^5\right) - \frac{3}{2}ax^4\right\}\right]$.
13. $x\{x - b - x(a - bx)\} + ax - x\{x - x(ax - b)\}$.

pressions w
more conver
powers of so

Example 1

The sum

Example 2.

The prod

Add togeth
arrange the r

1. $ax^3 - 2cx$,
2. $x^2 - x - 1$,
3. $a^2x^3 - 5x$,
4. $ax^2 + bx -$
5. $px^2 - qx$,

Multiply to
arrange the re

6. $ax^2 + bx +$
8. $ax^2 - bx - c$
10. $ax^2 - 2bx +$
12. $x^3 + ax^2 - b$
13. $ax^3 - x^2 + 3$
14. $x^4 - ax^3 - b$

63. In certain cases of addition, multiplication, etc., of expressions which involve literal coefficients, the results may be more conveniently written by grouping the terms according to powers of some common letter.

Example 1. Add together $ax^3 - 2bx^2 + 3$, $bx - cx^3 - x^2$ and $x^3 - ax^2 + cx$.

$$\begin{aligned}\text{The sum} &= ax^3 - 2bx^2 + 3 + bx - cx^3 - x^2 + x^3 - ax^2 + cx \\ &= ax^3 - cx^3 + x^3 - ax^2 - 2bx^2 - x^2 + bx + cx + 3 \\ &= (a - c + 1)x^3 - (a + 2b + 1)x^2 + (b + c)x + 3.\end{aligned}$$

Example 2. Multiply $ax^2 - 2bx + 3c$ by $px - q$.

$$\begin{aligned}\text{The product} &= (ax^2 - 2bx + 3c)(px - q) \\ &= apx^3 - 2bpx^2 + 3cp x - aqx^2 + 2bqx - 3cq \\ &= apx^3 - (2bp + aq)x^2 + (3cp + 2bq)x - 3cq.\end{aligned}$$

EXAMPLES VII. d.

Add together the following expressions, and in each case arrange the result according to powers of x :

1. $ax^3 - 2cx$, $bx^2 - cx^3$, $cx^3 - x$.
2. $x^3 - x - 1$, $ax^2 - bx^3$, $bx + x^3$.
3. $a^2x^3 - 5x$, $2ax^2 - 5ax^3$, $2x^3 - bx^2 - ax$.
4. $ax^2 + bx - c$, $qx - r - px^2$, $x^2 + 2x + 3$.
5. $px^3 - qx$, $qx^2 - px$, $q - x^3$, $px^2 + qx^3$.

Multiply together the following expressions, and in each case arrange the result according to powers of x :

- | | |
|---|------------------------------------|
| 6. $ax^2 + bx + 1$ and $cx + 2$. | 7. $cx^2 - 2x + 3$ and $ax - b$. |
| 8. $ax^2 - bx - c$ and $px + q$. | 9. $2x^2 - 3x - 1$ and $bx + c$. |
| 10. $ax^3 - 2bx + 3c$ and $x - 1$. | 11. $px^3 - 2x - q$ and $ax - 3$. |
| 12. $x^3 + ax^2 - bx - c$ and $x^3 - ax^2 - bx + c$. | |
| 13. $ax^3 - x^2 + 3x - b$ and $ax^3 + x^2 + 3x + b$. | |
| 14. $x^4 - ax^3 - bx^2 + cx + d$ and $x^4 + ax^3 - bx^2 - cx + d$. | |

CHAPTER VIII.

SIMPLE EQUATIONS.

64. AN **equation** asserts that two expressions are equal, but we do not usually employ the word equation in so wide a sense.

Thus the statement $x+3+x+4=2x+7$, which is *always* true whatever value x may have, is called an **identical equation**, or briefly an **identity**.

The parts of an equation to the right and left of the sign of equality are called **members** or **sides** of the equation, and are distinguished as the *right side* and *left side*.

65. Certain equations are only true for particular values of the symbols employed. Thus $3x=6$ is only true when $x=2$, and is called an **equation of condition**, or more usually an equation. Consequently an *identity* is an equation which is always true whatever be the values of the symbols involved; whereas an *equation* (in the ordinary use of the word) is only true for particular values of the symbols. In the above example $3x=6$, the value 2 is said to **satisfy** the equation. The object of the present chapter is to explain how to treat an equation of the simplest kind in order to discover the value which satisfies it.

66. The letter whose value it is required to find is called the **unknown quantity**. The process of finding its value is called **solving the equation**. The value so found is called the **root** or the **solution** of the equation.

67. The solution of equations, and the operations subsidiary to it, form an extremely important part of Mathematics. All sorts of mathematical problems consist in the indirect determination of some quantity by means of its relations to other quantities which are known, and these relations are all expressed by means of equations. The operation in general of solving a problem in Mathematics, other than a transformation, is firstly to express the conditions of the problem by means of one or more equations, and secondly to solve these equations. For example, the problem which is expressed by the equation above given is the very simple question, "What is the number such that if multiplied by 3, the product is 6?" In the present chapter, it is the second of these two operations, the solution of an equation, that is considered.

CHAP. VIII.]

68. An
the first de
denote the

69. The
upon the fol

1. If to
2. If from
3. If equ
4. If equ

70. Cons
It is requ
consistent w
Dividing b

Similarly,
multiplying

Again, in t
terms, we ha

71. To sol
This case o
quantity occu
transpose an
changing its si
Subtract x

Adding 8 to

Thus we see
appears as $-x$
one side and a

It is evident
Hence we may

Rule. Any
equation to the

68. An equation which involves the unknown quantity in the first degree is called a **simple equation**. It is usual to denote the unknown quantity by the letter x .

69. The process of solving a simple equation depends only upon the following **axioms**:

1. If to equals we add equals the sums are equal.
2. If from equals we take equals the remainders are equal.
3. If equals are multiplied by equals the products are equal.
4. If equals are divided by equals the quotients are equal.

70. Consider the equation $7x=14$.

It is required to find what numerical value x must have consistent with this statement.

Dividing both sides by 7 we get

$$x=2.$$

[Axiom 4.]

Similarly, if

$$\frac{x}{2} = -6,$$

multiplying both sides by 2, we get

$$x = -12.$$

[Axiom 3.]

Again, in the equation $7x-2x-x=23+15-10$, by collecting terms, we have

$$4x=28.$$

$$\therefore x=7.$$

71. To solve $3x-8=x+12$.

This case differs from the preceding in that the unknown quantity occurs on both sides of the equation. We can, however, **transpose** any term from one side to the other by simply *changing its sign*. This we proceed to shew.

Subtract x from both sides of the equation, and we get

$$3x-x-8=12.$$

[Axiom 2.]

Adding 8 to both sides, we have

$$3x-x=12+8.$$

[Axiom 1.]

Thus we see that $+x$ has been removed from one side, and appears as $-x$ on the other; and -8 has been removed from one side and appears as $+8$ on the other.

It is evident that similar steps may be employed in all cases. Hence we may enunciate the following rule:

Rule. Any term may be transposed from one side of the equation to the other by changing its sign.

It appears from this that *we may change the sign of every term in an equation*; for this is equivalent to transposing all the terms, and then making the right- and left-hand members change places.

Example. Take the equation $-3x - 12 = x - 24$.

$$\begin{array}{l} \text{Transposing,} \\ \text{or} \end{array} \quad \begin{array}{l} -x + 24 = 3x + 12, \\ 3x + 12 = -x + 24, \end{array}$$

which is the original equation with the sign of every term changed.

72. To solve $\frac{x}{2} - 3 = \frac{x}{4} + \frac{x}{5}$.

Here it will be convenient to begin by clearing the equation of fractional coefficients. This can always be done by multiplying both sides of the equation by the least common multiple of the denominators. [Axiom 3.]

Thus, multiplying by 20,

$$\begin{array}{l} \text{transposing,} \end{array} \quad \begin{array}{l} 10x - 60 = 5x + 4x; \\ 10x - 5x - 4x = 60; \\ \therefore x = 60. \end{array}$$

73. We can now give a general rule for solving any simple equation with one unknown quantity.

Rule. *First, if necessary, clear of fractions; then transpose all the terms containing the unknown quantity to one side of the equation, and the known quantities to the other. Collect the terms on each side; divide both sides by the coefficient of the unknown quantity, and the value required is obtained.*

Example 1. Solve $5(x - 3) - 7(6 - x) + 3 = 24 - 3(8 - x)$.

$$\begin{array}{l} \text{Removing brackets,} \\ \text{transposing,} \end{array} \quad \begin{array}{l} 5x - 15 - 42 + 7x + 3 = 24 - 24 + 3x; \\ 5x + 7x - 3x = 24 - 24 + 15 + 42 - 3; \\ \therefore 9x = 54; \\ \therefore x = 6. \end{array}$$

Example 2. Solve $5x - (4x - 7)(3x - 5) = 6 - 3(4x - 9)(x - 1)$.

Simplifying, we have

$$5x - (12x^2 - 41x + 35) = 6 - 3(4x^2 - 13x + 9).$$

And by removing brackets

$$5x - 12x^2 + 41x - 35 = 6 - 12x^2 + 39x - 27.$$

Erase the term $-12x^2$ on each side and transpose; thus

$$\begin{array}{l} 5x + 41x - 39x = 6 - 27 + 35; \\ \therefore 7x = 14; \\ \therefore x = 2. \end{array}$$

Note. Since it, in the first, until we have

Example 3.
Removing

transposing,

74. It is habit of ver- The habit of self-reliant a

In the cas when we su equation we

Example.

When $x = 2$

The right s

Thus, since equation.

Solve the f

1. $3x + 15 =$
3. $3x + 4 =$
5. $8(x - 1) +$
6. $15(x - 1)$

Note. Since the - sign before a bracket affects every term within it, in the first line of work of Ex. 2, we do not remove the brackets until we have formed the products.

Example 3. Solve $7x - 5[x - \{7 - 6(x - 3)\}] = 3x + 1$.

Removing brackets, we have

$$7x - 5[x - \{7 - 6x + 18\}] = 3x + 1,$$

$$7x - 5[x - 25 + 6x] = 3x + 1,$$

$$7x - 5x + 125 - 30x = 3x + 1;$$

transposing,

$$7x - 5x - 30x - 3x = 1 - 125;$$

$$\therefore -31x = -124;$$

$$\therefore x = 4.$$

74. It is extremely useful for the beginner to acquire the habit of **verifying**, that is, proving the truth of his results. The habit of applying such tests tends to make the student self-reliant and confident in his own accuracy.

In the case of simple equations we have only to shew that when we substitute the value of x in the two sides of the equation we obtain the same result.

Example. To shew that $x=2$ satisfies the equation

$$5x - (4x - 7)(3x - 5) = 6 - 3(4x - 9)(x - 1).$$

When $x=2$, the left side $5x - (4x - 7)(3x - 5)$

$$= 10 - (8 - 7)(6 - 5)$$

$$= 10 - 1 = 9.$$

The right side $6 - 3(4x - 9)(x - 1)$

$$= 6 - 3(8 - 9)(2 - 1)$$

$$= 6 - 3(-1) = 9.$$

Thus, since these two results are the same, $x=2$ satisfies the equation.

EXAMPLES VIII. a.

Solve the following equations :

1. $3x + 15 = x + 25.$

2. $2x - 3 = 3x - 7.$

3. $3x + 4 = 5(x - 2).$

4. $2x + 3 = 16 - (2x - 3).$

5. $8(x - 1) + 17(x - 3) = 4(4x - 9) + 4.$

6. $15(x - 1) + 4(x + 3) = 2(7 + x).$

Solve the following equations :

7. $5x - 6(x - 5) = 2(x + 5) + 5(x - 4).$
8. $8(x - 3) - (6 - 2x) = 2(x + 2) - 5(5 - x).$
9. $7(25 - x) - 2x = 2(3x - 25).$
10. $3(169 - x) - (78 + x) = 29x.$
11. $5x - 17 + 3x - 5 = 6x - 7 - 8x + 115.$
12. $7x - 39 - 10x + 15 = 100 - 33x + 26.$
13. $118 - 65x - 123 = 15x + 35 - 120x.$
14. $157 - 21(x + 3) = 163 - 15(2x - 5).$
15. $179 - 18(x - 10) = 158 - 3(x - 17).$
16. $97 - 5(x + 20) = 111 - 8(x + 3)$
17. $x - [3 + \{x - (3 + x)\}] = 5.$
18. $5x - (3x - 7) - \{4 - 2x - (6x - 3)\} = 10.$
19. $14x - (5x - 9) - \{4 - 3x - (2x - 3)\} = 30.$
20. $25x - 19 - [3 - \{4x - 5\}] = 3x - (6x - 5).$
21. $(x + 1)(2x + 1) = (x + 3)(2x + 3) - 14.$
22. $(x + 1)^2 - (x^2 - 1) = x(2x + 1) - 2(x + 2)(x + 1) + 20.$
23. $2(x + 1)(x + 3) + 8 = (2x + 1)(x + 5).$
24. $6(x^2 - 3x + 2) - 2(x^2 - 1) = 4(x + 1)(x + 2) - 24.$
25. $2(x - 4) - (x^2 + x - 20) = 4x^2 - (5x + 3)(x - 4) - 64.$
26. $(x + 15)(x - 3) - (x^2 - 6x + 9) = 30 - 15(x - 1).$
27. $2x - 5\{3x - 7(4x - 9)\} = 66.$
28. $20(2 - x) + 3(x - 7) - 2[x + 9 - 3\{9 - 4(2 - x)\}] = 22.$
29. $x + 2 - [x - 8 - 2\{8 - 3(5 - x) - x\}] = 0.$
30. $3(5 - 6x) - 5[x - 5\{1 - 3(x - 5)\}] = 23.$
31. $(x + 1)(2x + 3) = 2(x + 1)^2 + 8.$
32. $3(x - 1)^2 - 3(x^2 - 1) = x - 15.$
33. $(3x + 1)(2x - 7) = 6(x - 3)^2 + 7.$
34. $x^2 - 8x + 25 = x(x - 4) - 25(x - 5) - 16.$
35. $x(x + 1) + (x + 1)(x + 2) = (x + 2)(x + 3) + x(x + 4) - 0.$
36. $2(x + 2)(x - 4) = x(2x + 1) - 21.$
37. $(x + 1)^2 + 2(x + 3)^2 = 3x(x + 2) + 35.$
38. $4(x + 5)^2 - (2x + 1)^2 = 3(x - 5) + 180.$
39. $84 + (x + 4)(x - 3)(x + 5) = (x + 1)(x + 2)(x + 3)$
40. $(x + 1)(x + 2)(x + 6) = x^3 + 9x^2 + 4(7x - 1).$

75. The following examples illustrate the most useful methods of solving fractional equations.

viii.]

Example

Multiply
thus
removing br
transposing,
collecting te

Note. He

line between
as a bracket.

76. In c
multiply th
clear of frac

Example 2

Multiplying

transposing,

Now clear
thus

77. To so
express the d
but it is after

Example 1.

Expressing

clearing of fra
transposing,

Example 1. Solve $4 - \frac{x-9}{8} = \frac{x}{22} - \frac{1}{2}$.

Multiply by 88, the least common multiple of the denominators;

thus $352 - 11(x-9) = 4x - 44$;
 removing brackets, $352 - 11x + 99 = 4x - 44$;
 transposing, $-11x - 4x = -44 - 352 - 90$;
 collecting terms and changing signs, $15x = 495$;

$$\therefore x = 33.$$

Note. Here $-\frac{x-9}{8}$ is equivalent to $-\frac{1}{8}(x-9)$, the *vinculum* or line between the numerator and denominator having the same effect as a bracket. [Art. 58.]

76. In certain cases it will be found more convenient not to multiply throughout by the L.C.M. of the denominator, but to clear of fractions in two or more steps.

Example 2. Solve $\frac{x-4}{3} + \frac{2x-3}{35} = \frac{5x-32}{9} - \frac{x+9}{28}$.

Multiplying throughout by 9, we have

$$3x - 12 + \frac{18x-27}{35} = 5x - 32 - \frac{9x+81}{28};$$

transposing, $\frac{18x-27}{35} + \frac{9x+81}{28} = 2x - 20$.

Now clear of fractions by multiplying by $5 \times 7 \times 4$ or 140;

thus $72x - 108 + 45x + 405 = 280x - 2800$;
 $\therefore 2800 - 108 + 405 = 280x - 72x - 45x$;
 $\therefore 3097 = 163x$;
 $\therefore x = 19$.

77. To solve equations whose coefficients are decimals, we may express the decimals as vulgar fractions, and proceed as before; but it is often found more simple to work entirely in decimals.

Example 1. Solve $\cdot 6x + \cdot 25 - \frac{1}{9}x = 1\cdot 8 - \cdot 75x - \frac{1}{3}$.

Expressing the decimals as vulgar fractions, we have

clearing of fractions, $\frac{2}{3}x + \frac{1}{4} - \frac{1}{9}x = 1\frac{8}{9} - \frac{3}{4}x - \frac{1}{3}$;
 $24x + 9 - 4x = 68 - 27x - 12$;
 transposing, $24x - 4x + 27x = 68 - 12 - 9$,

$$47x = 47;$$

$$\therefore x = 1.$$

Example 2. Solve $\cdot 375x - 1\cdot 875 = \cdot 12x + 1\cdot 185$.
 Transposing, $\cdot 375x - \cdot 12x = 1\cdot 185 + 1\cdot 875$;
 collecting terms, $(\cdot 375 - \cdot 12)x = 3\cdot 06$,
 that is, $\cdot 255x = 3\cdot 06$;
 $\therefore x = \frac{3\cdot 06}{\cdot 255}$
 $= 12$.

EXAMPLES VIII. b.

1. $\frac{x}{4} + \frac{x-5}{3} = 10$.
2. $\frac{x-5}{10} + \frac{x+5}{5} = 5$.
3. $\frac{x-2}{2} + \frac{x+10}{9} = 5$.
4. $\frac{x+19}{5} = 3 + \frac{x}{4}$.
5. $\frac{x-4}{7} = \frac{x-10}{5}$.
6. $\frac{x-1}{8} = 1 + \frac{x+1}{18}$.
7. $\frac{4(x+2)}{5} = 7 + \frac{5x}{13}$.
8. $\frac{x+4}{14} + \frac{x-4}{6} = 2$.
9. $\frac{x+20}{9} + \frac{3x}{7} = 6$.
10. $\frac{x-8}{7} + \frac{x-3}{3} + \frac{5}{21} = 0$.
11. $\frac{x+5}{6} - \frac{x+1}{9} = \frac{x+3}{4}$.
12. $\frac{4-5x}{6} - \frac{1-2x}{3} = \frac{13}{42}$.
13. $\frac{5(x+5)}{8} - \frac{2(x-3)}{7} = 5\frac{1}{2}$.
14. $\frac{4(x+2)}{3} - \frac{6(x-7)}{7} = 12$.
15. $1 + \frac{x}{2} - \frac{2x}{3} = \frac{3x}{4} - 4\frac{1}{2}$.
16. $\frac{x}{2} + \frac{x}{3} - \frac{x}{4} + \frac{x}{5} = 7\frac{5}{8}$.
17. $\frac{3}{16}(x-1) - \frac{5}{12}(x-4) = \frac{2}{5}(x-6) + \frac{5}{48}$.
18. $x + \frac{5}{3}(x-7) - \frac{6}{7}(x-8) = 3x - 14\frac{1}{3}$.
19. $\frac{3x}{4} - \frac{6}{17}(x+10) - (x-3) = \frac{x-7}{51} - 4\frac{3}{4}$.
20. $\frac{7x}{5} - \frac{1}{14}(x-11) = \frac{3}{7}(x-25) + 34$.
21. $3 + \frac{x}{4} = \frac{1}{2}\left(4 - \frac{x}{3}\right) - \frac{5}{6} + \frac{1}{3}\left(11 - \frac{x}{2}\right)$.
22. $\frac{1}{5}(x-8) + \frac{4+x}{4} + \frac{x-1}{7} = 7 - \frac{23-x}{5}$.

23. $\frac{1}{3}\left(\frac{x}{4} - 3\right)$

24. $x - \left(3x - \frac{x}{4}\right)$

25. $\frac{x}{4} - \frac{x+5}{5}$

26. $\cdot 5x - \cdot 3$

28. $2\cdot 25x -$

30. $\cdot 6x - 7$

31. $12\{3x -$

32. $\cdot 25(x -$

33. $\frac{x + \cdot 75}{\cdot 125}$

35. $\cdot 5x - \cdot 2$

[Some of
will furnish.]

78. Before
draw attention
in solving e
down the sol

Case I. S
Multiplying

Case II. S

Multiplying

$$23. \frac{1}{3}\left(\frac{x}{4}-3\right)+\frac{5x}{6}-\frac{5x}{4}=\frac{x-12}{5}-\frac{x+3}{3}.$$

$$24. x-\left(3x-\frac{2x-5}{10}\right)=\frac{1}{6}(2x-57)-\frac{5}{3}.$$

$$25. \frac{x}{4}-\frac{x+10}{5}+\frac{4x}{4}=x-1-\frac{x-2}{3}.$$

$$26. \cdot 5x - \cdot 3x = \cdot 25x - 1.$$

$$27. 3+\frac{x}{5}=7-\frac{x}{2}.$$

$$28. 2\cdot 25x - \cdot 125 = 3x + 3\cdot 75.$$

$$29. \cdot 2x - \cdot 16x = \cdot 6 - \cdot 3.$$

$$30. \cdot 6x - \cdot 7x + \cdot 75x - \cdot 875x + 15 = 0.$$

$$31. 12\{3x - \cdot 25(x-4) - \cdot 3(5x+14)\} = 47.$$

$$32. \frac{\cdot 25(x-3) + \cdot 3(x-4)}{\cdot 125} = 5x - 19.$$

$$33. \frac{x + \cdot 75}{\cdot 125} - \frac{x - \cdot 25}{\cdot 25} = 15.$$

$$34. \frac{5}{6}x + \cdot 25x - \cdot 3x = x - 3.$$

$$35. \cdot 5x - \cdot 2x = \cdot 3x - 1\cdot 5.$$

$$36. 1\cdot 5 = \frac{\cdot 36}{\cdot 2} - \frac{\cdot 09x - \cdot 18}{\cdot 9}.$$

[Some of the examples in *Miscellaneous Examples II.*, p. 80, will furnish further practice in *Simple Equations*.]

78. Before concluding this chapter it will be worth while to draw attention to the following cases which occur so frequently in solving equations that the beginner should learn to write down the solution at sight.

Case I. Suppose

$$\frac{7x}{5} = \frac{4}{3}.$$

Multiplying both sides by 5, we have

$$\left. \begin{aligned} 7x &= \frac{4 \times 5}{3} \\ \therefore x &= \frac{4 \times 5}{3 \times 7} \end{aligned} \right\} \dots\dots\dots(1).$$

Case II. Suppose

$$\frac{5}{3x} = \frac{9}{7}.$$

Multiplying both sides by $3x$, we have

$$\left. \begin{aligned} 5 &= \frac{9 \times 3x}{7} \\ 5 \times 7 &= 9 \times 3x \\ \frac{5 \times 7}{9 \times 3} &= x \end{aligned} \right\} \dots\dots\dots(2).$$

By a careful examination of the results in (1) and (2), the truth of the following principles will be evident :

Any factor of the numerator of one side of an equation may be transferred to the denominator of the other side, and any factor of the denominator of one side may be transferred to the numerator of the other side.

The ready application of these principles will be found very useful.

Example 1. If

$$\frac{3x}{14} = \frac{9}{35},$$

then

$$x = \frac{9 \times 14}{35 \times 3} = 1\frac{1}{5}.$$

Example 2. If

$$\frac{2}{x} = -5,$$

then

$$\frac{2}{5} = -x; \quad \therefore x = -\frac{2}{5}.$$

After a little practice the arithmetic should be performed mentally, and the intermediate steps omitted.

EXAMPLES VIII. c.

Write down the values of x which satisfy the following equations :

1. $\frac{2}{x} = \frac{3}{4}$

2. $\frac{3}{7} = \frac{x}{14}$

3. $\frac{3}{5} = \frac{6}{x}$

4. $-\frac{x}{3} = 2$

5. $\frac{x}{17} = \frac{29}{51}$

6. $\frac{2x}{15} = \frac{8}{45}$

7. $\frac{3}{2x} = -\frac{1}{8}$

8. $-\frac{4x}{3} = -\frac{1}{2}$

9. $\frac{5}{3x} = \frac{25}{27}$

10. $\frac{5}{2x} = -\frac{10}{3}$

11. $\frac{13}{21} = \frac{65x}{84}$

12. $-\frac{7}{2} = \frac{1}{3x}$

13. $\frac{3}{8} = \frac{x}{4}$

14. $\frac{8}{21x} = -\frac{4}{7}$

15. $\frac{36}{35} = \frac{9}{5x}$

16. $\frac{5}{8} = \frac{15}{2x}$

17. $\frac{x}{18} = \frac{9}{42}$

18. $\frac{4}{3x} = \frac{16}{27}$

19. $\frac{49}{15} = \frac{7}{3x}$

20. $\frac{56}{15} = \frac{8x}{3}$

21. $\frac{19x}{7} = \frac{57}{49}$

79. IN the beginning means of solving will frequently question will be evident to analogous "50 by 6." the second number which is given

80. The deduction to the student find them

Example

Take a
The answer
Hence the
Similar

Example

Example

Example
4 miles an hour

In 1 hour

In a hour

Example

of each is the

d (2), the

tion may
ny factor
numerator

und very

entally,

lowing

3
5
5
7
1
x

7
9

CHAPTER IX.

SYMBOLICAL EXPRESSION.

79. In solving algebraical problems the chief difficulty of the beginner is to express the conditions of the question by means of symbols. A question proposed in algebraical symbols will frequently be found puzzling, when a similar arithmetical question would present no difficulty. Thus, the answer to the question "find a number greater than x by a " may not be self-evident to the beginner, who would of course readily answer an analogous arithmetical question, "find a number greater than 50 by 6." The process of addition which gives the answer in the second case supplies the necessary hint; and, just as the number which is greater than 50 by 6 is $50 + 6$, so the number which is greater than x by a is $x + a$.

80. The following examples will perhaps be the best introduction to the subject of this chapter. After the first we leave to the student the choice of arithmetical instances, should he find them necessary.

Example 1. By how much does x exceed 17?

Take a numerical instance; "by how much does 27 exceed 17?"

The answer obviously is 10, which is equal to $27 - 17$.

Hence the excess of x over 17 is $x - 17$.

Similarly the defect of x from 17 is $17 - x$.

Example 2. If x is one part of 45 the other part is $45 - x$.

Example 3. If x is one factor of 45 the other factor is $\frac{45}{x}$.

Example 4. How far can a man walk in a hours at the rate of 4 miles an hour?

In 1 hour he walks 4 miles,

In a hours he walks a times as far, that is, $4a$ miles.

Example 5. If £20 is divided equally among y persons, the share of each is the total sum divided by the number of persons, or $\frac{20}{y}$.

Example 6. If 17 be divided by 6 the quotient is 2, and the remainder 5,

$$\frac{17}{6} = 2 + \frac{5}{6}.$$

that is, So if N be divided by D , and the quotient be Q and the remainder R , we have

$$\frac{N}{D} = Q + \frac{R}{D},$$

or

$$N = QD + R.$$

Thus, if the divisor is x , the quotient y , and the remainder z , the dividend is $xy + z$.

Example 7. A and B are playing for money; A begins with $\pounds p$ and B with q shillings: after B has won $\pounds x$, how many shillings has each?

What B has won A has lost,

$\therefore A$ has $20(p - x)$ shillings,

B has $q + 20x$ shillings.

EXAMPLES IX. a.

1. What must be added to x to make y ?
2. By what must 3 be multiplied to make a ?
3. What dividend gives b as the quotient when 5 is the divisor?
4. What is the defect of $2c$ from $3d$?
5. By how much does $3k$ exceed k ?
6. If 100 be divided into two parts and one part be x what is the other?
7. If a be one factor of b , what is the other?
8. What number is less than 20 by c ?
9. What is the price in pence of a oranges at tenpence a dozen?
10. What is the price in pence of 100 oranges when x cost sixpence?
11. If the difference of two numbers be 11, and if the smaller be x , what is the greater?
12. If the sum of two numbers be c and one of them is 20, what is the other?
13. What is the excess of 90 over x ?
14. By how much does x exceed 30?
15. If 100 contains x five times, what is the value of x ?
16. What is the cost in pounds of 40 books at x shillings each?

17. In x ...
18. How...
19. If x take?
20. What...
21. What a score is x ?
22. How hour?
23. How...
24. In x ...
25. How an hour?
26. A tra from Bristol
27. How p miles an h
28. What travels 30 mi
29. A m has he?
30. If I s have I left?
31. Out o pence; exp
32. By h
33. What
34. If a l 3s. 4d., how
35. If I sovereigns an
36. In ho one horse eat
37. If I s out of a year
38. A booc if there are 1
39. I hav z pence in a have I left?
40. In a and the rest

17. In x years a man will be 36 years old, what is his present age?
18. How old will a man be in a years if his present age is x years?
19. If x men take 5 days to reap a field, how long will one man take?
20. What value of x will make $5x$ equal to 20?
21. What is the price in shillings of 120 apples, when the cost of a score is x pence?
22. How many hours will it take to walk x miles at 4 miles an hour?
23. How far can I walk in x hours at the rate of y miles an hour?
24. In x days a man walks y miles, what is his rate per day?
25. How many minutes will it take to walk x miles at a miles an hour?
26. A train goes x miles an hour, how long does it take to go from Bristol to London, a distance of 120 miles?
27. How many miles is it between two places, if a train travelling p miles an hour takes 5 hours to perform the journey?
28. What is the velocity in feet per second of a train which travels 30 miles in x hours?
29. A man has a crowns and b florins, how many shillings has he?
30. If I spend x shillings out of a sum of £20, how many shillings have I left?
31. Out of a purse containing $£a$ and b shillings a man spends c pence; express in pence the sum left.
32. By how much does $2x - 5$ exceed $x + 1$?
33. What number must be taken from $a - 2b$ to leave $a - 3b$?
34. If a bill is shared equally amongst x persons and each pays 3s. 4d., how many pence does the bill amount to?
35. If I give away c shillings out of a purse containing a sovereigns and b florins, how many shillings have I left?
36. In how many weeks will x horses eat 100 bushels of oats if one horse eats y bushels a week?
37. If I spend x shillings a week, how many pounds do I save out of a yearly income of $£y$?
38. A bookshelf contains x Latin, y Greek, and z English books: if there are 100 books, how many are there in other languages?
39. I have x pounds in my purse, y shillings in one pocket, and z pence in another: if I give away half-a-crown, how many pence have I left?
40. In a class of x boys, y work at Classics, z at Mathematics, and the rest are idle: what is the excess of workers over idlers?

81. We subjoin a few harder examples worked out in full.

Example 1. What is the present age of a man who x years hence will be m times as old as his son now aged y years?

In x years the son's age will be $y+x$ years; hence the father's age will be $m(y+x)$ years; therefore now the father's age is $m(y+x) - x$ years.

Example 2. Find the simple interest on $\pounds k$ in n years at f per cent.

Interest on $\pounds 100$ for 1 year is $\pounds f$,

$$\therefore \dots\dots\dots \pounds 1 \dots\dots\dots \pounds \frac{f}{100},$$

$$\therefore \dots\dots\dots \pounds k \dots\dots\dots \pounds \frac{kf}{100},$$

$$\therefore \text{Interest on } \pounds k \text{ for } n \text{ years is } \pounds \frac{nkf}{100}.$$

Example 3. A room is x yards long, y feet broad, and a feet high; find how many square yards of carpet will be required for the floor, and how many square yards of paper for the walls.

(1) The area of the floor is $3xy$ square feet;

$\therefore$ the number of square yards of carpet required is $\frac{3xy}{9} = \frac{xy}{3}$.

(2) The perimeter of the room is $2(3x+y)$ feet;

$\therefore$ the area of the walls is $2a(3x+y)$ square feet;

$\therefore$ number of square yards of paper required is $\frac{2a(3x+y)}{9}$.

Example 4. The digits of a number beginning from the left are a, b, c ; what is the number?

Here c is the digit in the units' place; b standing in the tens' place represents b tens; similarly a represents a hundreds. The number is therefore equal to a hundreds + b tens + c units

$$= 100a + 10b + c.$$

If the digits of the number are inverted, a new number is formed which is symbolically expressed by

$$100c + 10b + a.$$

Example 5. What is (1) the sum, (2) the product of three consecutive numbers of which the least is n ?

The numbers consecutive to n are $n+1, n+2$;

$$\therefore \text{the sum} = n + (n+1) + (n+2)$$

$$= 3n+3.$$

And the product $= n(n+1)(n+2)$.

We may r
by $2n$, where
sion is exactl

Similarly,
this expressio

Example 6.
c boys can mo
of n boys?

1. Write
2. Write
greatest.

3. Write c
one.

4. What i

5. What i

6. Find th

middle one is 2

7. A man

coach, b by tra

the boat carry

8. A horse

week: how ma

9. If a ma

years hence?

10. A boy i

that of his fath

11. What i

old as a child t

12. A 's age

old: find A 's a

13. What is

in full.
ears hence

the father's
age is

at f per

We may remark here that any *even* number may be denoted by $2n$, where n is any positive whole number; for this expression is exactly divisible by 2.

Similarly, any odd number may be denoted by $2n+1$; for this expression when divided by 2 leaves remainder 1.

Example 6. How many days will a men take to mow b acres if c boys can mow a acres in b days, and each man's work equals that of n boys?

Since c boys can mow a acres in b days;
 $\therefore$ 1 boy..... $\frac{bc}{c}$ days,
 $\therefore n$ boys, or 1 man, ... $\frac{bc}{n}$ days,
 $\therefore a$ men..... $\frac{bc}{an}$ days,
 $\therefore a$ men..... 1 acre... $\frac{bc}{a^2n}$ days;
 therefore a men can mow b acres in $\frac{b^2c}{a^2n}$ days.

EXAMPLES IX. b.

1. Write down four consecutive numbers of which x is the least.
2. Write down three consecutive numbers of which y is the greatest.
3. Write down five consecutive numbers of which x is the middle one.
4. What is the next even number after $2n$?
5. What is the odd number next before $2x+1$?
6. Find the sum of three consecutive odd numbers of which the middle one is $2n+1$.
7. A man makes a journey of x miles. He travels a miles by coach, b by train, and finishes the journey by boat. How far does the boat carry him?
8. A horse eats a bushels and a donkey b bushels of corn in a week: how many bushels will they together consume in n weeks?
9. If a man was x years old 5 years ago, how old will he be y years hence?
10. A boy is x years old, and 5 years hence his age will be half that of his father. How old is the father now?
11. What is the age of a man who y years ago was m times as old as a child then aged x years?
12. A 's age is double B 's, B 's is three times C 's, and C is x years old: find A 's age.
13. What is the interest on £1000 in b years at c per cent.?

14. What is the interest on $\text{£}x$ in a years at 5 per cent.?
15. What is the interest on $\text{£}50a$ in a years at a per cent.?
16. What is the interest on $\text{£}24xy$ in x months at y per cent. per annum?
17. A room is x yards in length, and y feet in breadth: how many square feet are there in the area of the floor?
18. A square room measures x feet each way: how many square yards of carpet will be required to cover it?
19. A room is x feet long and x yards in width: how many yards of carpet two feet wide will be required for the floor?
20. What is the cost in pounds of carpeting a room a yards long b feet broad with carpet costing c shillings a square yard?
21. How many yards of carpet x inches wide will be required to cover the floor of a room y feet long and z feet broad?
22. A room is a yards long, and b yards broad; in the middle there is a carpet c feet square: how many square yards of oil-cloth will be required to cover the rest of the floor?
23. How many miles can a person walk in 45 minutes if he walks a miles in x hours?
24. How long will it take a person to walk b miles if he walks 20 miles in c hours?
25. If a train travels a miles in b hours, how many feet does it move through in one second?
26. A train is running with a velocity of x feet per second: how many miles will it travel in y hours?
27. How long will x men take to mow y acres of corn, if each man mow z acres a day?
28. How many men will be required to do in x hours what y men do in zx hours?
29. What is the rate per cent. which will produce $\text{£}y$ interest from a principal of $\text{£}1000$ in r years?
30. Find in how many years a principal of $\text{£}a$ will produce $\text{£}p$ interest at r per cent. per annum.

*82. In Example 6, Art 80, we proved

$$\frac{N}{D} = Q + \frac{R}{D}$$

a result which gives in a single statement a general relation expressing the connection between a number, its divisor, and the resulting quotient and remainder.

This is an example of a very important class of algebraical statements known as *formulæ*, and it may be worth while briefly to foreshadow their use and application, not only in Algebra, but also in other branches of Mathematics and elementary Physics.

DEFINITION
ing among c
regarded as

Thus in th
quantities, w
of N . Or, a
must 96 be c
11?" Here
from the for

whence $D = 1$

*83. A fe
cases in one
algebraical f
class of resu
and easily ap
much of the
and their rea
occurrence.

It would b
allusion to oth
but on accoun
may be usefu
more element
studies.

(1) If a tr
given by the

(2) If a p
its volume

These ca
superficial
cubic inch
three quant
metric.

Example. T
each side of wh
number of cubi

From the f

DEFINITION. A *formula* is a relation established by reasoning among certain quantities, any one of which may in turn be regarded as the unknown.

Thus in the formula above mentioned, if Q , R , and D are given quantities, we have an equation to find the corresponding value of N . Or, a question may be proposed as follows: "By what must 96 be divided so as to give a quotient 5, and a remainder 11?" Here we have given $N=96$, $Q=5$, $R=11$, and therefore from the formula we obtain

$$\frac{96}{D} = 5 + \frac{11}{D},$$

whence $D=17$, the required divisor.

*83. A formula, it must be observed, includes all particular cases in one general statement; and so by the use of a single algebraical formula we are enabled briefly to express a whole class of results in a form at once simple, easily remembered, and easily applied. Experience will convince the student how much of the power and utility of Algebra lies in its formulæ, and their ready application to all kinds of problems of every day occurrence.

It would be out of place here to make more than a passing allusion to other branches of Mathematics, or to Physical Science; but on account of the interest and importance of the subject, it may be useful to draw the reader's attention to a few of the more elementary formulæ he is likely to meet with in his other studies.

(1) If a triangle on a base b , has a height h , its area (A) is given by the formula $A = \frac{1}{2}bh$.

(2) If a pyramid of height h stands on a base whose area is a^2 , its volume (V) is given by the formula

$$V = \frac{1}{3}a^2h.$$

In these cases any linear unit, inch, foot ... being chosen, the superficial and solid units will be respectively the square and cubic inch, &c.; and in each of these formulæ if two of the three quantities be given, the third is easily obtained by Arithmetic.

Example. The Great Pyramid of Egypt stands on a square base each side of which is 764 feet; and its height is 480 feet. Find the number of cubic feet of stone used in its construction.

$$\begin{aligned} \text{From the formula, } V &= \frac{1}{3} \times (764)^2 \times 480 \\ &= 160 \times 764 \times 764 \\ &= 93391360 \text{ cubic feet.} \end{aligned}$$

*84. We have in this chapter given several examples involving space, velocity, and time; and all these can be solved without difficulty by common sense reasoning. At the same time we may remark that they are only particular cases of the general formula $s=vt$, in which s denotes the space described by a body which moves with uniform velocity v for a time t .

In this formula, if t denotes the number of seconds the body has been in motion, and v the number of feet passed over in one second, then s is the space (in feet) described in t seconds.

Example. If a train has a velocity of 75 feet a second, how long will it take to cross a viaduct which is 300 yards in length?

Substituting the values of s and v (expressed in feet) in the formula, we get

$$900 = 75t,$$

$$t = \frac{900}{75}$$

$$= 12.$$

Therefore the time is 12 seconds.

*85. Another very interesting case is that of a body falling vertically under the action of gravity.

It is proved in works on Dynamics that if a body fall freely from rest, and if s denote the space (in feet) described in t seconds,

$$s = \frac{1}{2}gt^2.$$

In this formula g denotes the number of feet by which the velocity is increased in each successive second in consequence of the earth's attraction, and it is found by experiment that $g=32.2$ nearly.

Example 1. A stone dropped from the Clifton suspension bridge takes 4 seconds before it reaches the water. Find the height of the bridge above the river.

$$\begin{aligned}\text{From the above formula, } s &= \frac{1}{2} \times 32.2 \times (4)^2 \\ &= 257.6,\end{aligned}$$

and the height is therefore 257.6 feet.

Example 2. How long will it take a stone to reach the bottom of a well 144.9 feet deep?

$$\text{From the formula, } 144.9 = \frac{1}{2} \times 32.2 \times t^2;$$

$$\therefore t^2 = \frac{144.9}{16.1} = 9;$$

$$\therefore t = 3.$$

Therefore the time is 3 seconds.

PRO

86. THE
to solve vari

The metho
Represent
in symbolical
obtain a sim
already given

Example 1.
difference is 4

Let x be
Their sum
Hence

and
so that

The beginn
it satisfies th

Example 2.
greater may e
200.

Let x be t
Three tim

Eight tim

Whence t

and

CHAPTER X.

PROBLEMS LEADING TO SIMPLE EQUATIONS.

86. The principles of the last chapter may now be employed to solve various problems.

The method of procedure is as follows :

Represent the unknown quantity by a symbol x , and express in symbolical language the conditions of the question ; we thus obtain a simple equation which can be solved by the methods already given in Chapter VIII.

Example 1. Find two numbers whose sum is 28, and whose difference is 4.

Let x be the smaller number, then $x + 4$ is the greater.

Their sum is $x + (x + 4)$, which is to be equal to 28.

Hence $x + x + 4 = 28$;

$$2x = 24 ;$$

$$\therefore x = 12,$$

and $x + 4 = 16$;

so that the numbers are 12 and 16.

The beginner is advised to test his solution by finding whether it satisfies the data of the question or not.

Example 2. Divide 60 into two parts, so that three times the greater may exceed 100 by as much as 8 times the less falls short of 200.

Let x be the greater part, then $60 - x$ is the less.

Three times the greater part is $3x$, and its excess over 100 is

$$3x - 100.$$

Eight times the less is $8(60 - x)$, and its defect from 200 is

$$200 - 8(60 - x).$$

Whence the symbolical statement of the question is

$$3x - 100 = 200 - 8(60 - x) ;$$

$$3x - 100 = 200 - 480 + 8x,$$

$$480 - 100 - 200 = 8x - 3x,$$

$$5x = 180,$$

$$\therefore x = 36, \text{ the greater part,}$$

$$\text{and } 60 - x = 24, \text{ the less.}$$

Example 3. Divide £47 between A , B , C , so that A may have £10 more than B , and B £8 more than C .

Suppose that C has x pounds; then B has $x+8$ pounds, and A has $x+8+10$ pounds.

$$\begin{aligned}\text{Hence} \quad & x+(x+8)+(x+8+10)=47; \\ & x+x+8+x+8+10=47, \\ & \qquad \qquad \qquad 3x=21; \\ & \qquad \qquad \qquad \therefore x=7;\end{aligned}$$

so that C has £7, B £15, A £25.

Example 4. A person spent £28. 4s. in buying geese and ducks; if each goose cost 7s., and each duck 3s., and if the total number of birds bought was 108: how many of each did he buy?

In questions of this kind it is of essential importance to have all quantities expressed in the same denomination; in the present instance it will be convenient to express the money in shillings.

Let x be the number of geese, then $108-x$ is the number of ducks.

Since each goose costs 7 shillings, x geese cost $7x$ shillings.

And since each duck costs 3 shillings, $108-x$ ducks cost $3(108-x)$ shillings.

Therefore the amount spent is

$$7x+3(108-x) \text{ shillings};$$

but the question states that the amount is also £28. 4s., that is 564 shillings.

$$\text{Hence} \quad 7x+3(108-x)=564;$$

$$7x+324-3x=564,$$

$$4x=240,$$

$$\therefore x=60, \text{ the number of geese,}$$

$$108-x=48, \text{ the number of ducks.}$$

and

Example 5. A is twice as old as B , ten years ago he was four times as old: what are their present ages?

Let B 's age be x years, then A 's age is $2x$ years.

Ten years ago their ages were respectively, $x-10$ and $2x-10$ years; thus we have

$$2x-10=4(x-10);$$

$$2x-10=4x-40,$$

$$2x=30;$$

$$\therefore x=15,$$

so that B is 15 years old, A 30 years.

Note. In the above examples the unknown quantity x represents a number of pounds, ducks, years, etc.; and the student must be careful to avoid beginning a solution with a supposition of the kind, "let $x=A$'s share" or "let x =the ducks", or any statement so vague and inexact.

1. One number them.

2. The difference greater the result

3. Find a number 11 than its double

4. A man train, and the miles, how far

5. What 28?

6. If 288 to three times

7. Twenty 16 is above set

8. Divide shall be equal

9. Find the

10. The sum to it is five times

11. Find their difference

12. A and money is double

13. Find a product of the second.

14. The difference is 121: find

15. The difference squares is 27:

16. Divide more than A ,

17. A sum florins or shillings

18. If silk in buying 23 y per yard.

19. A father only be twice

20. A is 20 as B 's is below

EXAMPLES X. a.

1. One number exceeds another by 5, and their sum is 29; find them.
2. The difference between two numbers is 8; if 2 be added to the greater the result will be three times the smaller: find the numbers.
3. Find a number such that its excess over 50 may be greater by 11 than its defect from 89.
4. A man walks 10 miles, then travels a certain distance by train, and then twice as far by coach. If the whole journey is 70 miles; how far does he travel by train?
5. What two numbers are those whose sum is 58, and difference 28?
6. If 288 be added to a certain number, the result will be equal to three times the excess of the number over 12: find the number.
7. Twenty-three times a certain number is as much above 14 as 16 is above seven times the number: find it.
8. Divide 105 into two parts, one of which diminished by 20 shall be equal to the other diminished by 15.
9. Find three consecutive numbers whose sum shall equal 84.
10. The sum of two numbers is 8, and one of them with 22 added to it is five times the other: find the numbers.
11. Find two numbers differing by 10 whose sum is equal to twice their difference.
12. *A* and *B* begin to play each with £60. If they play till *A*'s money is double *B*'s, what does *A* win?
13. Find a number such that if 5, 15, and 35 are added to it, the product of the first and third results may be equal to the square of the second.
14. The difference between the squares of two consecutive numbers is 121: find the numbers.
15. The difference of two numbers is 3, and the difference of their squares is 27: find the numbers.
16. Divide £380 between *A*, *B*, and *C*, so that *B* may have £30 more than *A*, and *C* may have £20 more than *B*.
17. A sum of £8. 17s. is made up of 124 coins which are either florins or shillings: how many are there of each?
18. If silk costs six times as much as linen, and I spend £9. 8s. in buying 23 yards of silk and 50 yards of linen: find the cost of each per yard.
19. A father is four times as old as his son; in 24 years he will only be twice as old: find their ages.
20. *A* is 25 years older than *B*, and *A*'s age is as much above 20 as *B*'s is below 85: find their ages.

21. A 's age is six times B 's, and fifteen years hence A will be three times as old as B : find their ages.

22. A sum of £4. 5s. was paid in crowns, half-crowns, and shillings. The number of half-crowns used was four times the number of crowns and twice the number of shillings: how many were there of each?

23. The sum of the ages of A and B is 30 years, and five years hence A will be three times as old as B : find their present ages.

24. In a cricket match the byes were double of the wides, and the remainder of the score was greater by three than twelve times the number of byes. If the whole score was 138, how were the runs obtained?

25. The length of a room exceeds its breadth by 3 feet; if the length had been increased by 3 feet, and the breadth diminished by 2 feet, the area would not have been altered: find the dimensions.

26. The length of a room exceeds its breadth by 8 feet; if each had been increased by 2 feet, the area would have been increased by 60 square feet: find the original dimensions of the room.

27. We add some problems which lead to equations with fractional coefficients.

Example 1. Find two numbers which differ by 4, and such that one-half of the greater exceeds one-sixth of the less by 8.

Let x be the smaller number, then $x+4$ is the greater.

One-half of the greater is represented by $\frac{1}{2}(x+4)$, and one sixth of the less by $\frac{1}{6}x$.

Hence

$$\frac{1}{2}(x+4) - \frac{1}{6}x = 8;$$

multiplying by 6,

$$3x + 12 - x = 48;$$

$$\therefore 2x = 36;$$

and

$$\therefore x = 18, \text{ the less number,}$$

$$x+4 = 22, \text{ the greater.}$$

Example 2. A has £9, and B has 4 guineas; after B has won from A a certain sum, A has then five-sixths of what B has: how much did B win?

Suppose that B wins x shillings, A has then $180 - x$ shillings, and B has $84 + x$ shillings.

Hence

$$180 - x = \frac{5}{6}(84 + x);$$

$$1080 - 6x = 420 + 5x,$$

$$11x = 660;$$

$$\therefore x = 60.$$

Therefore B wins 60 shillings, or £3.

1. Find a may be equal

2. What together make

3. There by 3: find it.

4. Find a fifths of it by

5. The together make

6. Two exceeds one fif

7. Two other: find th

8. There a greater exceed

9. Find th by 10, 17, and

10. A and five-elevenths than half of wh

11. From a divided by 4; and the result

12. In a cel besides this it how much port

13. Two-fift B 's is equal to

14. A , B , a than five-sixths the share of eac

15. A man a it, and gained t

16. The wid width had been would have bee

17. What is when he has tw 3 per cent., and

18. I bought five-sixths of th teen for sixpenc

EXAMPLES X. b.

1. Find a number such that the sum of its sixth and ninth parts may be equal to 15.
2. What is the number whose eighth, sixth, and fourth parts together make up 13?
3. There is a number whose fifth part is less than its fourth part by 3: find it.
4. Find a number such that six-sevenths of it shall exceed four-fifths of it by 2.
5. The fifth, fifteenth, and twenty-fifth parts of a number together make up 23: find the number.
6. Two consecutive numbers are such that one-fourth of the less exceeds one fifth of the greater by 1: find the numbers.
7. Two numbers differ by 28, and one is eight-ninths of the other: find them.
8. There are two consecutive numbers such that one-fifth of the greater exceeds one-seventh of the less by 3: find them.
9. Find three consecutive numbers such that if they be divided by 10, 17, and 26 respectively, the sum of the quotients will be 10.
10. *A* and *B* begin to play with equal sums, and when *B* has lost five-elevenths of what he had to begin with, *A* has gained £6 more than half of what *B* has left: what had they at first?
11. From a certain number 3 is taken, and the remainder is divided by 4; the quotient is then increased by 4 and divided by 5 and the result is 2: find the number.
12. In a cellar one-fifth of the wine is port and one-third claret; besides this it contains 15 dozen of sherry and 30 bottles of hock: how much port and claret does it contain?
13. Two-fifths of *A*'s money is equal to *B*'s, and seven-ninths of *B*'s is equal to *C*'s; in all they have £770: what have they each?
14. *A*, *B*, and *C* have £1285 between them: *A*'s share is greater than five-sixths of *B*'s by £25, and *C*'s is four-fifteenths of *B*'s: find the share of each.
15. A man sold a horse for £35 and half as much as he gave for it, and gained thereby ten guineas: what did he pay for the horse?
16. The width of a room is two-thirds of its length. If the width had been 3 feet more, and the length 3 feet less, the room would have been square: find its dimensions.
17. What is the property of a person whose income is £430, when he has two-thirds of it invested at 4 per cent., one-fourth at 3 per cent., and the remainder at 2 per cent.?
18. I bought a certain number of apples at three a penny, and five-sixths of that number at four a penny; by selling them at sixteen for sixpence I gained $3\frac{1}{2}d.$: how many apples did I buy?

CHAPTER XI.

 HIGHEST COMMON FACTOR, LOWEST COMMON MULTIPLE
OF SIMPLE EXPRESSIONS.

Highest Common Factor.

88. DEFINITION. The **highest common factor** of two or more algebraical expressions is the expression of highest dimensions [Art. 24] which divides each of them without remainder.

The abbreviation H.C.F. is sometimes used instead of the words *highest common factor*.

89. In the case of *simple expressions* the highest common factor can be written down by inspection.

Example 1. The highest common factor of a^4, a^3, a^2, a^6 is a^2 .

Example 2. The highest common factor of a^3b^4, ab^5c^2, a^2b^7c is ab^4 ; for a is the highest power of a that will divide a^3, a, a^2 ; b^4 is the highest power of b that will divide b^4, b^5, b^7 ; and c is not a common factor.

90. If the expressions have numerical coefficients, find by Arithmetic their greatest common measure, and prefix it as a coefficient to the algebraical highest common factor.

Example. The highest common factor of $21a^4x^3y, 35a^2x^4y, 28a^3xy^4$ is $7a^2xy$; for it consists of the product of

- (1) the greatest common measure of the numerical coefficients;
- (2) the highest power of each letter which divides every one of the given expressions.

EXAMPLES XI. a.

Find the highest common factor of

1. $4ab^2, 2a^2b.$
2. $3x^2y^2, x^3y^2.$
3. $6xy^2z, 8x^2y^2z^2.$
4. $abc, 2ab^2c.$
5. $5a^3b^3, 15abc^2.$
6. $9x^2y^2z^2, 12xy^2z.$
7. $4a^2b^3c^2, 6a^3b^2c^3.$
8. $7a^2b^4c^5, 14ab^3c^3.$
9. $15x^4y^3z^2, 12x^2yz^2.$
10. $8a^2x, 6abxy, 10abx^2y^2.$
11. $49ax^2, 63ay^2, 56az^2.$
12. $17ab^2c, 34a^2bc, 51abc^2.$
13. $a^3x^2y^3, b^3xy^2, c^3x^2y.$
14. $24a^2b^3c^3, 64a^3b^3c^2, 48a^3b^3c^3.$
15. $25xy^2z, 100x^2yz, 125xy.$
16. $a^3bpxy, b^3qxy, a^3bxr^2.$
17. $15a^3b^3c^7, 60a^3b^7c^6, 25a^4b^5c^2.$
18. $35a^2c^3b, 42a^3c^2b^2, 30ac^2b^3.$

91. DEFINITION. The **lowest common multiple** of two or more algebraical expressions which are not prime to each other is the expression of lowest dimensions which is divisible by each of them without remainder. The abbreviation L.C.M. is sometimes used instead of the words *lowest common multiple*.

92. In the case of *simple expressions* the lowest common multiple can be written down by inspection.

Example 1.

Example 2. The lowest common multiple of a^3b^7 and a^2b^4 is a^3b^7 ; for a^3 is the highest power of a that will divide a^3, a^2 ; and b^7 is the highest power of b that will divide b^7, b^4 .

93. If the expressions have numerical coefficients, find by Arithmetic their least common measure, and prefix it as a coefficient to the algebraical lowest common multiple.

Example. The lowest common multiple of $28a^3xy^4$ and $420a^2x^4y$ is $840a^3x^4y^4$.

- (1) the least common measure of the numerical coefficients;
- (2) the highest power of each letter which divides every one of the given expressions.

Find the lowest common multiple of

1. $abc, 2a^2.$
2. $5a^2bc^3, 4a^3b^2c.$
3. $ac, bc, ab.$
4. $2x, 3y, 4z.$
5. $a^2bc, b^2ca, c^2ab.$
6. $2x^2y^3, 3xy^2z.$
7. $35a^2c^3b, 42a^3c^2b^2, 30ac^2b^3.$

Find both the highest common factor and the lowest common multiple of

19. $2abc, 3ca, 4ab.$
20. $13a^2bc, 39a^3b^2c^2.$
21. $3ab, 2bc, 4ac.$
22. $15p^3q^4, 20p^2q^5.$

Lowest Common Multiple.

91. DEFINITION. The **lowest common multiple** of two or more algebraical expressions is the expression of lowest dimensions which is divisible by each of them without remainder.

The abbreviation L.C.M. is sometimes used instead of the words *lowest common multiple*.

92. In the case of *simple expressions* the lowest common multiple can be written down by inspection.

Example 1. The lowest common multiple of a^4, a^3, a^2, a^6 is a^6 .

Example 2. The lowest common multiple of a^3b^4, ab^5, a^2b^7 is a^3b^7 ; for a^3 is the lowest power of a that is divisible by each of the quantities a^3, a, a^2 ; and b^7 is the lowest power of b that is divisible by each of the quantities b^4, b^5, b^7 .

93. If the expressions have numerical coefficients, find by Arithmetic their least common multiple, and prefix it as a coefficient to the algebraical lowest common multiple.

Example. The lowest common multiple of $21a^4x^2y, 35a^2xy, 28a^2xy^4$ is $420a^4x^2y^4$; for it consists of the product of

- (1) the least common multiple of the numerical coefficients;
- (2) the lowest power of each letter which is divisible by every power of that letter occurring in the given expressions.

EXAMPLES XI. b.

Find the lowest common multiple of

- | | | |
|--|--|------------------------|
| 1. $abc, 2a^2.$ | 2. $x^2y^2, xyz.$ | 3. $3x^2yz, 4x^2y^3.$ |
| 4. $5a^2bc^3, 4ab^2c.$ | 5. $3a^4b^2c^3, 5a^2b^3c^5.$ | 6. $12ab, 8xy.$ |
| 7. $ac, bc, ab.$ | 8. $a^2c, bc^3, cb^3.$ | 9. $2ab, 3bc, 4ca.$ |
| 10. $2x, 3y, 4z.$ | 11. $3x^2, 4y^2, 3z^2.$ | 12. $7a^2, 2ab, 3b^3.$ |
| 13. $a^2bc, b^3ca, c^2ab.$ | 14. $5a^2c, 6cb^3, 3bc^2.$ | |
| 15. $2x^2y^3, 3xy, 4x^3y^4.$ | 16. $7x^4y, 8xy^5, 2x^3y^3.$ | |
| 17. $35a^2c^3b, 42a^3cb^3, 30ac^2b^3.$ | 18. $66a^4b^2c^3, 44a^3b^4c^2, 24a^2b^3c^4.$ | |

Find both the highest common factor and the lowest common multiple of

- | | | |
|--------------------------------------|----------------------------------|----------------------------------|
| 19. $2abc, 3ca, 4bca.$ | 20. $2xy, 4yz, 6zxy.$ | 21. $9abc, 3b^2c, cab.$ |
| 22. $13a^2bc, 39a^3bc^2.$ | 23. $17xyz^2, 51x^2y.$ | 24. $15x^2y^2z, 25xy^3z^2.$ |
| 25. $3ab, 2bc, 5cab.$ | 26. $17m^2n^3p^2, 51m^4p^4.$ | 27. $x^3y^2, y^2z^4, z^4x^2y^5.$ |
| 28. $15p^3q^4, 20m^2p^2q^3, 30mp^3.$ | 29. $72k^2m^3n^4, 108k^3m^2n^5.$ | |

CHAPTER XII.

ELEMENTARY FRACTIONS.

94. DEFINITION. If a quantity x be divided into b equal parts, and a of these parts be taken, the result is called the fraction $\frac{a}{b}$ of x . If x be the unit, the fraction $\frac{a}{b}$ of x is called simply "the fraction $\frac{a}{b}$ "; so that the fraction $\frac{a}{b}$ represents a equal parts, b of which make up the unit.

95. In this chapter we propose to deal only with the easier kinds of fractions, where the numerator and denominator are simple expressions. Their reduction and simplification will be performed by the usual arithmetical rules. The proofs of these rules will be given in Chapters XIX. and XXI.

Rule. To reduce a fraction to its lowest terms: divide numerator and denominator by every factor which is common to them both, that is by their highest common factor.

Dividing numerator and denominator of a fraction by a common factor is called *cancelling* that factor.

Examples. (1) $\frac{6a^2c}{9ac^3} = \frac{2a}{3c^2}$
 (2) $\frac{7x^2yz}{28x^3yz^2} = \frac{1}{4xz}$
 (3) $\frac{35a^5b^3c}{7ab^2c} = \frac{5a^4b}{1} = 5a^4b$.

EXAMPLES XII. a.

Reduce to lowest terms :

- | | | | |
|-----------------------------------|---------------------------------------|-------------------------------------|---------------------------------------|
| 1. $\frac{3a}{8ab}$ | 2. $\frac{4a^2}{16ab}$ | 3. $\frac{2xy^2}{5x^2y}$ | 4. $\frac{3abc}{15a^2b^3c}$ |
| 5. $\frac{x^2yz^3}{x^3yz^2}$ | 6. $\frac{15ab}{25bc}$ | 7. $\frac{21x^2y^2}{28y^2z^2}$ | 8. $\frac{8a^2b}{12b^2c}$ |
| 9. $\frac{12mn^2p}{15m^2np^3}$ | 10. $\frac{15m^2p^3}{18n^4p}$ | 11. $\frac{abc^2}{a^3b^2c}$ | 12. $\frac{3x^2yz^3}{5xy^2z^2}$ |
| 13. $\frac{2xy^2z^4}{4x^2y^2z}$ | 14. $\frac{5a^3b^2c^4}{15ab^2c}$ | 15. $\frac{mn^4pq}{m^2n^3p^2}$ | 16. $\frac{4m^3n^2p^5}{6m^4np^2}$ |
| 17. $\frac{15ax^3y^2}{25a^2xy^2}$ | 18. $\frac{39a^2b^4c^3}{52a^3b^3c^4}$ | 19. $\frac{38k^2p^3m^4}{57k^2pm^2}$ | 20. $\frac{46x^3y^4z^5}{69x^2y^3z^4}$ |

96. Rule. *metically, multiply the numerator by the denominator, and all the denominator by the numerator, and divide the result by the product of the two denominators.*

Example 1.

by cancelling

Example 2.

all the factors

97. Rule. *Divide the numerator by the denominator, and the remainder by the denominator, and the result by the denominator.*

Example.

all the factors

Simplify the

1. $\frac{2ab}{3cd} \times \frac{c^2d^3}{ab^2}$
4. $\frac{7a^2b^3}{9ax^2y} \times \frac{18}{15}$
7. $\frac{3a^2b}{4b^3c} \times \frac{2c^2}{8a^3}$
9. $\frac{7m^2p}{17x^2y} \times \frac{51}{21}$
11. $\frac{15b^2}{40c} \times \frac{27c^3}{81d}$
13. $\frac{8ax^2}{7by} \times \frac{49c}{64d}$
15. $\frac{45a^2b^3c^4}{27x^2y^2z} \times \frac{9}{1}$
17. $\frac{m^2}{8n} \times \frac{36p^2q}{81mn}$

Multiplication and Division of Fractions.

96. Rule. To multiply algebraical fractions: as in Arithmetic, multiply together all the numerators to form a new numerator, and all the denominators to form a new denominator.

Example 1. $\frac{2a}{3b} \times \frac{5x^2}{2a^2b} \times \frac{3b^2}{2x} = \frac{2a \times 5x^2 \times 3b^2}{3b \times 2a^2b \times 2x} = \frac{5x}{2a}$

by cancelling like factors in numerator and denominator.

Example 2. $\frac{3a^2b}{5c^2} \times \frac{7bc}{3a^3} \times \frac{5ca}{7b^2} = 1$,

all the factors cancelling each other.

97. Rule. To divide one fraction by another: invert the divisor and proceed as in multiplication.

Example. $\frac{7a^3}{4x^2y^2} \times \frac{6c^2x}{5ab^2} \div \frac{28a^2c^2}{15b^2xy^2} = \frac{7a^3}{4x^2y^2} \times \frac{6c^2x}{5ab^2} \times \frac{15b^2xy^2}{28a^2c^2} = \frac{9c}{8x}$

all the other factors cancelling each other.

EXAMPLES XII. b.

Simplify the following expressions:

1. $\frac{2ab}{3cd} \times \frac{c^2d^3}{ab^2}$
2. $\frac{12a^2bc}{8ab^3} \times \frac{24ab^2}{36bc^2}$
3. $\frac{15xyz^3}{a^2bc} \times \frac{3a^3x}{5yz}$
4. $\frac{7a^2b^3}{9ax^2y} \times \frac{18x^2c}{15ac^4}$
5. $\frac{8m^2n^3}{5x^2yz} \times \frac{15xyz^3}{16mn^2}$
6. $\frac{21k^2p^3}{13mn^2} \times \frac{39n^2m^2}{28p^2k^3}$
7. $\frac{3a^2b}{4b^3c} \times \frac{2c^2}{8a^3} \div \frac{6ac}{16b^2x}$
8. $\frac{2x^2y}{3yz} \times \frac{5x^2x}{7xy^2} \div \frac{21x^2y^3z^2}{40xy^2z}$
9. $\frac{7m^2p}{17x^2y} \times \frac{51y^3z}{21p^2n} \div \frac{m^2x^2}{p^2yz}$
10. $\frac{26xk^2p^3}{58mp^4} \times \frac{2xk^2}{13pkm} \div \frac{2x^2k^4}{87m^2p^2}$
11. $\frac{15b^2}{40c} \times \frac{27c^2}{81d^2} \div \frac{abc}{14d^3}$
12. $\frac{b^3}{3c} \times \frac{4c^2}{5d^3} \div \frac{16a^2b^2c^2}{15d^3}$
13. $\frac{8ax^2}{7by} \times \frac{49cy^2}{64dx^3}$
14. $\frac{15abc}{16xyz} \times \frac{128x^3y^2z^2}{100a^2bc}$
15. $\frac{45a^2b^3c^4}{27x^2y^2z} \times \frac{243xy^2z^3}{180a^2bc^2}$
16. $\frac{104xyzk^2p}{28xy^2kp^2} \times \frac{56y^2z^3p}{26y^2z^2k}$
17. $\frac{m^2}{8n} \times \frac{36p^2q^2}{81mn} \div \frac{15mpx^5}{27n^2x^2y}$
18. $\frac{a^3}{b^3} \times \frac{xy^2}{ab} \times \frac{pb^2}{ax} \div \frac{ap}{b^2}$

Reduction to a Common Denominator.

98. In order to find the sum or difference of any fractions, we must, as in Arithmetic, first reduce them to a common denominator; and it is most convenient to take the lowest common multiple of the denominators of the given fractions.

Example. Express with lowest common denominator the fractions

$$\frac{a}{3xy}, \frac{b}{6xyz}, \frac{c}{2yz}.$$

The lowest common multiple of the denominators is $6xyz$. Multiplying the numerator of each fraction by the factor which is required to make its denominator $6xyz$, we have the equivalent fractions

$$\frac{2ax}{6xyz}, \frac{b}{6xyz}, \frac{3cx}{6xyz}.$$

Note. The same result would clearly be obtained by dividing the lowest common denominator by each of the denominators in turn, and multiplying the corresponding numerators by the respective quotients.

EXAMPLES XII. c.

Express as equivalent fractions with common denominator:

- | | | | |
|-------------------------------------|--------------------------------------|--------------------------------------|--|
| 1. $\frac{2x}{a}, \frac{y}{2a}.$ | 2. $\frac{4x}{3y}, \frac{y}{x^2}.$ | 3. $\frac{a}{2b}, \frac{b}{c}.$ | 4. $\frac{a}{b}, \frac{c}{d}, 2.$ |
| 5. $\frac{2a}{b}, \frac{b}{3c}.$ | 6. $\frac{m}{4n}, \frac{p}{5n}.$ | 7. $\frac{k}{2x}, \frac{p}{3x}.$ | 8. $\frac{m}{3x}, \frac{n}{6x}.$ |
| 9. $\frac{a}{bc}, \frac{b}{ca}.$ | 10. $\frac{a}{x}, \frac{b}{x^2}.$ | 11. $\frac{2}{x}, \frac{3}{y}.$ | 12. $\frac{x}{y}, \frac{y}{x}, 3x.$ |
| 13. $\frac{2x}{3y}, \frac{3y}{2x}.$ | 14. $\frac{4a}{5b}, \frac{3a}{10c}.$ | 15. $\frac{3a}{7b}, \frac{5b}{21c}.$ | 16. $\frac{2}{a}, \frac{b}{3}, \frac{a}{9}.$ |

Addition and Subtraction of Fractions.

99. Rule. To add or subtract fractions: express all the fractions with their lowest common denominator; form the algebraical sum of the numerators, and retain the common denominator.

Example 1. Simplify $\frac{5x}{3} + \frac{3}{4}x - \frac{7x}{6}.$

The least common denominator is 12.

$$\text{The expression} = \frac{20x + 9x - 14x}{12} = \frac{15x}{12} = \frac{5x}{4}.$$

Example 2.

The expr

Example 3.

The expr

Note. The erasing equals cancelling e reducing fractions he must numerator and

Thus in $\frac{6ax}{3}$ and not the because it on fraction is the

When no be understood

Example 4.

If a fraction before combin

Example 5.

Simplify th

$$1. \frac{x}{2} + \frac{x}{3}.$$

$$5. \frac{x}{2} + \frac{y}{5}.$$

$$9. \frac{x}{7} - \frac{y}{21}.$$

$$13. \frac{2x}{3} + \frac{4x}{5}.$$

Example 2. Simplify $\frac{3ab}{5x} - \frac{ab}{2x} - \frac{1}{10} \cdot \frac{ab}{x}$.

The expression = $\frac{6ab - 5ab - ab}{10x} = \frac{0}{10x} = 0$.

Example 3. Simplify $\frac{2x}{a^2c^2} - \frac{y}{3ca^3}$.

The expression = $\frac{6ax - cy}{3a^3c^2}$, and admits of no further simplification.

Note. The beginner must be careful to distinguish between **erasing equal terms with different signs**, as in Example 2, and **cancelling equal factors** in the course of multiplication, or in reducing fractions to lowest terms. Moreover, in simplifying fractions he must remember that a factor can only be removed from numerator and denominator when it divides each *taken as a whole*.

Thus in $\frac{6ax - cy}{3a^3c^2}$, c cannot be cancelled because it only divides cy and not the *whole* numerator. Similarly a cannot be cancelled because it only divides $6ax$ and not the whole numerator. The fraction is therefore in its simplest form.

When no denominator is expressed the denominator 1 may be understood.

Example 4. $3x - \frac{a^2}{4y} = \frac{3x}{1} - \frac{a^2}{4y} = \frac{12xy - a^2}{4y}$.

If a fraction is not in its lowest terms it should be simplified before combining it with other fractions.

Example 5. $\frac{ax}{2} - \frac{x^2y}{3xy} = \frac{ax}{2} - \frac{x}{3} = \frac{3ax - 2x}{6}$.

EXAMPLES XII. d.

Simplify the following expressions :

- | | | | |
|-------------------------------------|-------------------------------------|--------------------------------------|--------------------------------------|
| 1. $\frac{x}{2} + \frac{x}{3}$. | 2. $\frac{y}{4} - \frac{y}{5}$. | 3. $\frac{a}{3} - \frac{a}{4}$. | 4. $\frac{2x}{3} - \frac{5}{x}$. |
| 5. $\frac{x}{2} + \frac{y}{5}$. | 6. $\frac{a}{4} - \frac{b}{6}$. | 7. $\frac{m}{8} - \frac{n}{12}$. | 8. $\frac{2m}{15} - \frac{n}{5}$. |
| 9. $\frac{x}{7} - \frac{y}{21}$. | 10. $\frac{a}{13} + \frac{b}{39}$. | 11. $\frac{p}{16} - \frac{q}{48}$. | 12. $\frac{5m}{12} - \frac{n}{36}$. |
| 13. $\frac{2x}{3} + \frac{4x}{5}$. | 14. $\frac{5x}{4} - \frac{4x}{5}$. | 15. $\frac{5x}{6} - \frac{7x}{12}$. | 16. $\frac{2a}{5} - \frac{4b}{15}$. |

Simplify the following expressions :

17. $\frac{a}{2} - \frac{a}{3} + \frac{a}{5}$.

18. $\frac{x}{4} - \frac{x}{8} + \frac{x}{12}$.

19. $\frac{x}{3} + \frac{x}{6} - \frac{x}{9}$.

20. $\frac{2x}{3} - \frac{x}{6} + \frac{3x}{4}$.

21. $\frac{5x}{6} - \frac{x}{12} + \frac{x}{9}$.

22. $\frac{7x}{8} + \frac{x}{12} - \frac{x}{4}$.

23. $\frac{x}{a} - \frac{y}{b}$.

24. $\frac{3x^3}{ax^2} + \frac{2y}{3b}$.

25. $a + \frac{b}{c}$.

26. $x - \frac{y^2}{yz}$.

27. $\frac{a^3}{3a^2} - \frac{b^2}{a}$.

28. $a^2 + \frac{b^3}{a}$.

29. $\frac{3x^2}{6x} - \frac{y^2}{x^2}$.

30. $p^3 - \frac{k^5}{p^2}$.

MISCELLANEOUS EXAMPLES II.

(Chiefly on Chapters I.-VIII.)

1. What expression must be added to $4x^3 - 3x^2 + 2$ to produce $4x^3 + 7x - 6$?

2. If $A = 6x - 3y + 2z$, $B = x + y + z$, and $C = 10x + y - 7z$, find the value of $A + 4B - C$.

3. If $x = 3$, $y = 4$, $z = 1$, find the value of

$$\sqrt{2xy + 4xz} + \sqrt{9y} + \frac{2xyz}{3}.$$

4. Simplify by removing brackets

$$a^2 + 2d^2 - (2e^2 - b^2) - \{(d^2 - c^2 - e^2) + (d^2 - e^2)\}.$$

5. Multiply $x^3 + x^2 + 3x + 5$ by $x^2 - x - 2$.

6. Solve the equations :

(1) $3 - 4x = 36x - 17$;

(2) $5x - 15 = 17x + 21$.

7. Divide $x^4 - 10x^2 + 9$ by $x^2 - 2x - 3$.

8. Simplify $7a - 4b - \{5a - 3[b - 2(a - b)]\}$.

9. In an examination A has $x + y$ marks, B has $2x - 3y$, and C has twice as many as A ; how many marks have A , B , and C together?

10. Find the sum of $1 - 2x + x^2$, $3x - 2x^2$, $5x^2 - 7x - 2$, arranging the result in descending powers of x .

11. Write

(1) ()

12. Solve t

(1) ()

(2) ()

13. From t
sum of $-8ab$,

14. When a

15. From w
as to give for r

16. Multipl

17. Simplify

18. Solve th

Verify the se

19. Divide :

20. Add tog

$$a + 2b - (2$$

21. To what
make $9x^3 - 6x -$

22. What va
less than the pr

23. When a =

24. Solve the

Shew also that

E.A.

11. Write down the following products:

$$(1) (x+17)(x-3); \quad (2) (3x-8)(8x+3).$$

12. Solve the equations:

$$(1) 7x-3-(7-5x)=3-3x-(5x+8);$$

$$(2) (5x+1)(x-2)-(4x-3)(3x-1)=10-(7x+2)(x+1).$$

13. From the sum of $3ab$, $-5ab$, $2ab$, $7ab$, $-9ab$, subtract the sum of $-8ab$, $6ab$, $-9ab$, $10ab$.

14. When $a=4$, $b=3$, $c=2$, find the numerical value of

$$\frac{2a+b(2c-a)}{3b-\sqrt{2c^3}}.$$

15. From what expression must $11a^2-5ab-7bc$ be subtracted so as to give for remainder $7b(a+c)+5a^2$?

16. Multiply x^2+6x^2+8x-8 by x^2-2x+4 .

17. Simplify

$$12a-[6a-2\{3a-4(b-a)\}-(9a+8)].$$

18. Solve the equations:

$$(1) 3(2x-1)+2(3x-2)+3=4(x-5);$$

$$(2) \frac{1}{3}(x+1)+\frac{1}{4}(x+3)=\frac{1}{5}(x+4)+16.$$

Verify the solution in each case.

19. Divide $3p^5+16p^4-33p^3+14p^2$ by p^2+7p .

20. Add together

$$a+2b-(2c+d), \quad 3a-(b-2c)+2d, \quad \text{and} \quad 2a-[b-(2c-3d)].$$

21. To what expression must $7x^2-6x^2-5x$ be added so as to make $9x^2-6x-7x^2$?

22. What value of x will make the product of $x+1$ and $2x+1$ less than the product of $x+3$ and $2x+3$ by 14?

23. When $a=2$, $b=3$, $c=1$, $q=4$, $r=6$, find the value of

$$5a^3c^r-3a^2b+2r^2a^5-c^2bq.$$


24. Solve the equation:

$$x-\frac{x-13}{9}=\frac{6x+1}{5}+\frac{2}{3}\left(6-\frac{3x}{2}\right).$$


Shew also that $x=3$ does not satisfy the equation,

E.A.

F



(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

25. A horse can eat $3m+2n$ bushels of corn in a week; how many weeks will he be in eating $12m^2-7mn-10n^2$ bushels?

26. Subtract the sum of

$$2x^3-3x+4 \text{ and } -3x^2+2x-7$$

from

$$4x^3-3x^2+x-6-[2x^3-(x-6)].$$

27. Find the value of $a^3+b^3+c^3-3abc$, when $a=1$, $b=4$, $c=-5$.

28. Solve the equations:

$$(1) \frac{2x}{15} + \frac{x-6}{12} = \frac{3}{10} \left(\frac{x}{2} - 5 \right);$$

$$(2) \frac{2(x-1)}{5} + \frac{15}{2} \left(1 - \frac{x}{3} \right) + \frac{19}{10} = \frac{9}{5} \left(\frac{x}{6} - \frac{1}{3} \right).$$

29. Divide $3y^5-37y^4+35y^3+7y^2+2$ by $y(y-1)(y+4)-2$.

30. Divide £1120 between A and B so that for every half-crown that A receives B may receive a shilling.

31. Find the value of

$$(a+b)^2 + (b+c)^2 + (c+a)^2$$

when $a=-1$, $b=-2$, $c=3$.

32. Multiply $(2m^2+8)(m+2)$ by $3m-6$.

33. Divide the product of

$$x-2, x+3, \text{ and } 2x-7$$

by the sum of $3(x^2-2x-2)$ and $5x-x^2-15$.

34. A man walks at the rate of a miles an hour for p hours; he then rides for q hours at the rate of b miles an hour. How far has he travelled, and how long would it have taken to ride the same distance at c miles an hour?

Also work out the result supposing $p=7$, $q=3$, $a=4$, $b=9$, $c=11$.

35. Solve the equations:

$$(1) \frac{3x}{2} - \frac{5}{7} = 21x - \frac{1}{3} \left(2x + 10\frac{3}{14} \right);$$

$$(2) 3x-4 - \frac{4(7x-9)}{15} = \frac{4}{5} \left(6 + \frac{x-1}{3} \right).$$

36. An egg-dealer bought a certain number of eggs at $1s. 4d.$ per score, and five times the number at $6s. 3d.$ per hundred. He sold the whole at $10d.$ per dozen, gaining £1. 7s. by the transaction. How many eggs did he buy?

100. Con-
unknown qu-

From this

that is,

From this
to x there wi
be able to f
satisfy the gi

For instan

Again, if a

But if also

as

we have from

If now we
the values of

Therefore

Multiplying

Substituting

and

Thus, if bo
of x and y , th

week; how
s?

4, $c = -5$.

2.

half-crown

ours; he
w far has
the same

9, $c = 11$.

. 4d. per
He sold
nsaction.

CHAPTER XIII.

SIMULTANEOUS EQUATIONS.

100. CONSIDER the equation $2x + 5y = 23$, which contains *two* unknown quantities.

From this we get $5y = 23 - 2x$,

that is, $y = \frac{23 - 2x}{5}$ (1).

From this it appears that for every value we choose to give to x there will be one corresponding value of y . Thus we shall be able to find as many pairs of values as we please which satisfy the given equation.

For instance, if $x = 1$, then from (1) $y = \frac{21}{5}$.

Again, if $x = -2$, then $y = \frac{27}{5}$; and so on.

But if also we have a second equation of the same kind, such as

$$3x + 4y = 24,$$

we have from this $y = \frac{24 - 3x}{4}$ (2).

If now we seek values of x and y which satisfy *both* equations, the values of y in (1) and (2) must be identical.

Therefore $\frac{23 - 2x}{5} = \frac{24 - 3x}{4}$.

Multiplying up, $92 - 8x = 120 - 15x$;

$$\therefore 7x = 28;$$

$$\therefore x = 4.$$

Substituting this value in the first equation, we have

$$8 + 5y = 23;$$

$$\therefore 5y = 15;$$

$$\therefore y = 3,$$

$$x = 4.$$

and

Thus, if both equations are to be satisfied by the *same* values of x and y , there is only one solution possible.

101. DEFINITION. When two or more equations are satisfied by the same values of the unknown quantities they are called **simultaneous equations**.

We proceed to explain the different methods for solving simultaneous equations. In the present chapter we shall confine our attention to the simpler cases in which the unknown quantities are involved in the first degree.

102. In the example already worked we have used the method of solution which best illustrates the meaning of the term *simultaneous equation*; but in practice it will be found that this is rarely the readiest mode of solution. It must be borne in mind that since the two equations are simultaneously true, *any* equation formed by combining them will be satisfied by the values of x and y which satisfy the original equations. Our object will always be to obtain an equation which involves *one only* of the unknown quantities.

103. The process by which we get rid of either of the unknown quantities is called **elimination**, and it must be effected in different ways according to the nature of the equations proposed.

Example 1. Solve

$$\begin{aligned} 3x + 7y &= 27 \dots\dots\dots(1), \\ 5x + 2y &= 16 \dots\dots\dots(2). \end{aligned}$$

To eliminate x we multiply (1) by 5 and (2) by 3, so as to make the coefficients of x in both equations equal. This gives

$$15x + 35y = 135,$$

$$15x + 6y = 48;$$

$$29y = 87;$$

$$\therefore y = 3.$$

To find x , substitute this value of y in *either* of the given equations.

Thus from (1) $3x + 21 = 27;$

$$\therefore x = 2,$$

$$y = 3.$$

and

Note. When one of the unknowns has been found, it is immaterial which of the equations we use to complete the solution. Thus, in the present example, if we substitute 3 for y in (2), we have

$$5x + 6 = 16;$$

$$\therefore x = 2, \text{ as before.}$$

Example 2.

Here it will be found that multiplying (1) by 2 and adding,

Substitute

and

Note. Add the equations, unlike in sign, and the signs will be the same.

Example 3.

Here we can eliminate y from (1). Thus

and from (1)

104. Any equation which is sufficient; but frequently the same result is obtained by subtracting (1) from (2).

Example 1.

Noticing that the coefficients of y are the same, we can multiply (1) by 2 and subtract (2) from it.

Thus

subtracting, that is, subtracting (2) from (1), and therefore find

Example 2. Solve $7x + 2y = 47$ (1),
 $5x - 4y = 1$ (2).

Here it will be more convenient to eliminate y .

Multiplying (1) by 2, $14x + 4y = 94$,
 and from (2) $5x - 4y = 1$;
 adding, $19x = 95$;
 $\therefore x = 5$.

Substitute this value in (1),

$$\therefore 35 + 2y = 47 ;$$

and

$$\therefore y = 6, \}$$

$$x = 5. \}$$

Note. *Add* when the coefficients of one unknown are equal and *unlike* in sign ; *subtract* when the coefficients are equal and *like* in sign.

Example 3. Solve $2x = 5y + 1$ (1),
 $24 - 7x = 3y$ (2).

Here we can eliminate x by substituting in (2) its value obtained from (1). Thus

$$24 - \frac{7}{2}(5y + 1) = 3y ;$$

$$\therefore 48 - 35y - 7 = 6y ;$$

$$\therefore 41 = 41y ;$$

$$\therefore y = 1, \}$$

$$x = 3. \}$$

and from (1)

104. Any one of the methods given above will be found sufficient ; but there are certain arithmetical artifices which will frequently shorten the work.

Example 1. Solve $171x - 213y = 642$ (1),
 $114x - 326y = 244$ (2).

Noticing that 171 and 114 contain a common factor 57, we shall make the coefficients of x in the two equations equal to the *least common multiple* of 171 and 114 if we multiply (1) by 2 and (2) by 3.

Thus $342x - 426y = 1284$,
 $342x - 978y = 732$;
 subtracting, $552y = 552$;
 that is, $y = 1$,
 and therefore from (1) $x = 5$.

- Example 2.* Solve $127x + 59y = 1928$ (1),
 $59x + 127y = 1792$ (2).
 By addition, $186x + 186y = 3720$;
 $\therefore x + y = 20$ (3).
 Subtracting (2) from (1), $68x - 68y = 136$;
 $\therefore x - y = 2$ (4).
 Thus, by an easy combination of (1) and (2), the problem is reduced to the solution of the equations (3) and (4). From these we obtain by addition $2x = 22$, and by subtraction $2y = 18$.
 Therefore $x = 11$, and $y = 9$.

EXAMPLES XIII. a.

Solve the equations :

- | | | |
|---|---|--|
| 1. $3x + 4y = 10$,
$4x + y = 9$. | 2. $x + 2y = 13$,
$3x + y = 14$, | 3. $4x + 7y = 29$,
$x + 3y = 11$. |
| 4. $2x - y = 9$,
$3x - 7y = 19$. | 5. $5x + 6y = 17$,
$6x + 5y = 16$. | 6. $2x + y = 10$,
$7x + 8y = 53$. |
| 7. $8x - y = 34$,
$x + 8y = 53$. | 8. $15x + 7y = 29$,
$9x + 15y = 39$, | 9. $14x - 3y = 39$,
$6x + 17y = 35$. |
| 10. $28x - 23y = 33$,
$63x - 25y = 101$. | 11. $35x + 17y = 86$,
$56x - 13y = 17$. | 12. $15x + 77y = 92$,
$55x - 33y = 22$. |
| 13. $5x - 7y = 0$,
$7x + 5y = 74$. | 14. $21x - 50y = 60$,
$28x - 27y = 199$. | 15. $39x - 8y = 99$,
$52x - 15y = 80$. |
| 16. $5x = 7y - 21$,
$21x - 9y = 75$. | 17. $6y - 5x = 18$,
$12x - 9y = 0$. | 18. $8x = 5y$,
$13x = 8y + 1$. |
| 19. $3x = 7y$,
$12y = 5x - 1$. | 20. $19x + 17y = 0$,
$2x - y = 53$. | 21. $93x + 15y = 123$,
$15x + 93y = 201$. |

105. We add a few cases in which, before proceeding to solve, it will be necessary to simplify the equations.

- Example 1.* Solve $5(x + 2y) - (3x + 11y) = 14$ (1),
 $7x - 9y - 3(x - 4y) = 38$ (2).
 From (1) $5x + 10y - 3x - 11y = 14$;
 $\therefore 2x - y = 14$ (3).
 From (2) $7x - 9y - 3x + 12y = 38$;
 $\therefore 4x + 3y = 38$ (4).
 From (3) $6x - 3y = 42$.
 By addition $10x = 80$; whence $x = 8$. From (3) we obtain $y = 2$.

*Example 2.*Clear of fr
from (1)

From (2)

Eliminatio

Eliminatio

Note. Sol
second unk
stituting the

Solve the

$$1. \frac{2x}{3} + y =$$

$$x + \frac{y}{4} =$$

$$4. x - y = 5$$

$$\frac{x}{4} - \frac{y}{5} = 2$$

$$7. \frac{2}{5}x - \frac{1}{12}y =$$

$$4x - y =$$

$$10. \frac{x}{7} + \frac{y}{5} = 1$$

$$x + \frac{y}{3} = 4$$

Example 2. Solve $3x - \frac{y-5}{7} = \frac{4x-3}{2}$ (1),

$\frac{3y+4}{5} - \frac{1}{3}(2x-5) = y$ (2).

Clear of fractions. Thus

from (1) $42x - 2y + 10 = 28x - 21$;

$\therefore 14x - 2y = -31$ (3).

From (2) $9y + 12 - 10x + 25 = 15y$;

$\therefore 10x + 6y = 37$ (4).

Eliminating y from (3) and (4), we find that

$$x = \frac{14}{13}.$$

Eliminating x from (3) and (4), we find that

$$y = \frac{207}{26}.$$

Note. Sometimes, as in the present instance, the value of the second unknown is more easily found by elimination than by substituting the value of the unknown already found.

EXAMPLES XIII. b.

Solve the equations :

1. $\frac{2x}{3} + y = 16,$

2. $\frac{x}{5} + \frac{y}{2} = 5,$

3. $\frac{5x}{6} - y = 3,$

$x + \frac{y}{4} = 14.$

$x - y = 4.$

$x - \frac{5y}{6} = 8.$

4. $x - y = 5,$

5. $\frac{x}{9} + \frac{y}{7} = 10,$

6. $x = 3y,$

$\frac{x}{4} - \frac{y}{5} = 2.$

$\frac{x}{3} + y = 50.$

$\frac{x}{3} + y = 34.$

7. $\frac{2}{5}x - \frac{1}{12}y = 3,$

8. $\frac{1}{2}x - \frac{1}{5}y = 4,$

9. $2x + y = 0,$

$4x - y = 20.$

$\frac{1}{7}x + \frac{1}{15}y = 3.$

$\frac{1}{2}y - 3x = 8.$

10. $\frac{x}{7} + \frac{y}{5} = 1\frac{3}{7},$

11. $3x - 7y = 0,$

12. $\frac{x}{5} - \frac{y}{4} = 0,$

$x + \frac{y}{3} = 4\frac{2}{3}.$

$\frac{2}{7}x + \frac{5}{3}y = 7.$

$3x + \frac{1}{2}y = 17.$

Solve the equations :

$$13. \frac{x}{3} + \frac{y}{4} = 3x - 7y - 37 = 0.$$

$$14. \frac{x+1}{10} = \frac{3y-5}{2} = \frac{x-y}{8}.$$

$$15. \frac{x+3}{5} = \frac{8-y}{4} = \frac{3(x+y)}{8}.$$

$$16. \frac{x}{13} - \frac{y}{7} = 6x - 10y - 8 = 0.$$

106. In order to solve simultaneous equations which contain two unknown quantities we have seen that we must have two equations. Similarly we find that in order to solve simultaneous equations which contain three unknown quantities we must have three equations.

Rule. *Eliminate one of the unknowns from any pair of the equations, and then eliminate the same unknown from another pair. Two equations involving two unknowns are thus obtained, which may be solved by the rules already given. The remaining unknown is then found by substituting in any one of the given equations.*

Example 1. Solve

$$\begin{aligned} 6x + 2y - 5z &= 13 & \dots\dots\dots(1), \\ 3x + 3y - 2z &= 13 & \dots\dots\dots(2), \\ 7x + 5y - 3z &= 26 & \dots\dots\dots(3). \end{aligned}$$

Choose y as the unknown to be eliminated.

Multiply (1) by 3 and (2) by 2,

$$\begin{aligned} 18x + 6y - 15z &= 39, \\ 6x + 6y - 4z &= 26; \\ \text{subtracting,} \quad 12x - 11z &= 13 & \dots\dots\dots(4). \end{aligned}$$

Again, multiply (1) by 5 and (3) by 2,

$$\begin{aligned} 30x + 10y - 25z &= 65, \\ 14x + 10y - 6z &= 52; \\ \text{subtracting,} \quad 16x - 19z &= 13 & \dots\dots\dots(5). \end{aligned}$$

Multiply (4) by 4 and (5) by 3,

$$\begin{aligned} 48x - 44z &= 52, \\ 48x - 57z &= 39; \\ \text{subtracting,} \quad 13z &= 13, \end{aligned}$$

and from (4)
from (1)

$$\begin{aligned} z &= 1, \\ x &= 2, \\ y &= 3. \end{aligned}$$

Note. After a little practice the student will find that the solution may often be considerably shortened by a suitable combination of the proposed equations. Thus, in the present instance, by adding (1) and (2) and subtracting (3) we obtain $2x - 4z = 0$, or $x = 2z$. Substituting in (1) and (2) we have two easy equations in y and z .

Some mo
with advan

Example 2

From the e
we have

Also, from
we have

And, from
we have
Eliminating

and from (1)
whence
Also by sub

Example 3.

Multiplying

or

Thus the c
equation whic
but a single e
minate. [Art

In this and
the equations
deducible from
between the u
other equation

Some modification of the foregoing rule may often be used with advantage.

Example 2. Solve $\frac{x}{2} - 1 = \frac{y}{6} + 1 = \frac{z}{7} + 2,$

$$\frac{y}{3} + \frac{z}{2} = 13.$$

From the equation

$$\frac{x}{2} - 1 = \frac{y}{6} + 1,$$

we have

$$3x - y = 12 \dots\dots\dots(1).$$

Also, from the equation

$$\frac{x}{2} - 1 = \frac{z}{7} + 2,$$

we have

$$7x - 2z = 42 \dots\dots\dots(2).$$

And, from the equation

$$\frac{y}{3} + \frac{z}{2} = 13,$$

we have

$$2y + 3z = 78 \dots\dots\dots(3).$$

Eliminating z from (2) and (3), we have

$$21x + 4y = 282;$$

and from (1)

$$12x - 4y = 48;$$

whence

$$x = 10, y = 18.$$

Also by substitution in (2) we obtain $z = 14$.

Example 3. Consider the equations

$$5x - 3y - z = 6 \dots\dots\dots(1),$$

$$13x - 7y + 3z = 14 \dots\dots\dots(2),$$

$$7x - 4y = 8 \dots\dots\dots(3).$$

Multiplying (1) by 3 and adding to (2), we have

$$28x - 16y = 32,$$

or

$$7x - 4y = 8.$$

Thus the combination of equations (1) and (2) leads us to an equation which is identical with (3), and so to find x and y we have but a single equation $7x - 4y = 8$, the solution of which is indeterminate. [Art. 100.]

In this and similar cases the anomaly arises from the fact that the equations are not *independent*; in other words, one equation is deducible from the others, and therefore contains no new relation between the unknown quantities which is not already implied in the other equations.

EXAMPLES XIII. c.

Solve the equations :

$$\begin{aligned} 1. \quad & x+2y+2z=11, \\ & 2x+y+z=7, \\ & 3x+4y+z=14. \end{aligned}$$

$$\begin{aligned} 3. \quad & x+4y+3z=17, \\ & 3x+3y+z=16, \\ & 2x+2y+z=11. \end{aligned}$$

$$\begin{aligned} 5. \quad & 2x+y+z=16, \\ & x+2y+z=9, \\ & x+y+2z=3. \end{aligned}$$

$$\begin{aligned} 7. \quad & 3x+2y-z=20, \\ & 2x+3y+6z=70, \\ & x-y+6z=41. \end{aligned}$$

$$9. \quad 3x-4y=6z-16, \quad 4x-y-z=5, \quad x=3y+2(z-1).$$

$$10. \quad 5x+2y=14, \quad y-6z=-15, \quad x+2y+z=0.$$

$$11. \quad x-\frac{y}{5}=6, \quad y-\frac{z}{7}=8, \quad z-\frac{x}{2}=10.$$

$$12. \quad \frac{y+z}{4}=\frac{z+x}{3}=\frac{x+y}{2}, \quad x+y+z=27.$$

$$13. \quad \frac{y-z}{3}=\frac{y-x}{2}=5z-4x, \quad y+z=2x+1.$$

$$14. \quad 2x+3y=5, \quad 2z-y=1, \quad 7x-9z=3.$$

$$\begin{aligned} 15. \quad & \frac{1}{2}(x+z-5)=y-z \\ & =2x-11 \\ & =9-(x+2z). \end{aligned}$$

$$\begin{aligned} 2. \quad & x+3y+4z=14, \\ & x+2y+z=7, \\ & 2x+y+2z=2. \end{aligned}$$

$$\begin{aligned} 4. \quad & 3x-2y+z=2, \\ & 2x+3y-z=5, \\ & x+y+z=6. \end{aligned}$$

$$\begin{aligned} 6. \quad & x-2y+3z=2, \\ & 2x-3y+z=1, \\ & 3x-y+2z=9. \end{aligned}$$

$$\begin{aligned} 8. \quad & 2x+3y+4z=20, \\ & 3x+4y+5z=26, \\ & 3x+5y+6z=31. \end{aligned}$$

Example 1.

Multiply (1)

adding,

multiplying u

and by substi

Example 2.

clearing of fra

from (1)

from (2)

from (3)

Multiply (4)

dividing by 7.

from (5)

from (5)

from (4)

*107. DEFINITION. If the product of two quantities be equal to unity, each is said to be the **reciprocal** of the other. Thus if $ab=1$, a and b are **reciprocals**. They are so called because $a=\frac{1}{b}$ and $b=\frac{1}{a}$; and consequently a is related to b exactly as b is related to a .

The reciprocals of x and y are $\frac{1}{x}$ and $\frac{1}{y}$ respectively, and in solving the following equations we consider $\frac{1}{x}$ and $\frac{1}{y}$ as the unknown quantities.

Example 1. Solve $\frac{8}{x} - \frac{9}{y} = 1$ (1),

$\frac{10}{x} + \frac{6}{y} = 7$ (2).

Multiply (1) by 2 and (2) by 3; thus

$$\frac{16}{x} - \frac{18}{y} = 2,$$

$$\frac{30}{x} + \frac{18}{y} = 21;$$

adding,

$$\frac{46}{x} = 23;$$

multiplying up,

$$46 = 23x,$$

$$x = 2;$$

and by substituting in (1),

$$y = 3.$$

Example 2. Solve $\frac{1}{2x} + \frac{1}{4y} - \frac{1}{3z} = \frac{1}{4}$ (1),

$\frac{1}{x} = \frac{1}{3y}$ (2),

$\frac{1}{x} - \frac{1}{5y} + \frac{4}{z} = 2\frac{2}{15}$ (3),

clearing of fractional coefficients, we obtain

from (1) $\frac{6}{x} + \frac{3}{y} - \frac{4}{z} = 3$ (4),

from (2) $\frac{3}{x} - \frac{1}{y} = 0$ (5),

from (3) $\frac{15}{x} - \frac{3}{y} + \frac{60}{z} = 32$ (6).

Multiply (4) by 15 and add the result to (6); we have

$$\frac{105}{x} + \frac{42}{y} = 77;$$

dividing by 7, $\frac{15}{x} + \frac{6}{y} = 11$ (7);

from (5) $\frac{18}{x} - \frac{6}{y} = 0;$

$$\therefore \frac{33}{x} = 11;$$

$$\therefore x = 3,$$

from (5) $y = 1,$

from (4) $z = 2.$

*EXAMPLES XIII. d.

Solve the equations :

$$1. \frac{5}{x} + \frac{6}{y} = 3,$$

$$\frac{15}{x} + \frac{3}{y} = 4.$$

$$2. \frac{6}{x} - \frac{7}{y} = 2,$$

$$\frac{2}{x} + \frac{14}{y} = 3.$$

$$3. \frac{12}{x} - \frac{4}{y} = 2,$$

$$\frac{3}{x} - \frac{2}{y} = 0.$$

$$4. \frac{5}{x} + \frac{16}{y} = 79,$$

$$\frac{16}{x} - \frac{1}{y} = 44.$$

$$5. \frac{21}{x} + \frac{12}{y} = 5,$$

$$\frac{1}{y} - \frac{1}{x} = \frac{1}{42}.$$

$$6. \frac{5}{x} + \frac{3}{y} = 30,$$

$$\frac{9}{x} - \frac{5}{y} = 2.$$

$$7. \frac{8}{x} - \frac{9}{y} = 7,$$

$$6\left(\frac{1}{x} + \frac{1}{y}\right) = 1,$$

$$8. \frac{25}{x} + \frac{24}{y} = 1,$$

$$20\left(\frac{2}{x} + \frac{3}{y}\right) = 7,$$

$$9. \frac{4}{x} + \frac{27}{y} = 42,$$

$$\frac{14}{x} - \frac{15}{y} = 1.$$

$$10. \frac{3}{x} + \frac{5}{y} = \frac{8}{15},$$

$$9y - 22x = \frac{3xy}{25}.$$

$$11. \frac{1}{4x} + \frac{1}{3y} = 2,$$

$$\frac{1}{y} - \frac{1}{2x} = 1.$$

$$12. 2y - x = 4xy,$$

$$\frac{4}{y} - \frac{3}{x} = 9,$$

$$13. \frac{1}{x} - \frac{2}{y} + 4 = 0,$$

$$\frac{1}{y} - \frac{1}{z} + 1 = 0,$$

$$\frac{2}{z} + \frac{3}{x} = 14.$$

$$14. \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 36,$$

$$\frac{1}{x} + \frac{3}{y} - \frac{1}{z} = 28,$$

$$\frac{1}{x} + \frac{1}{3y} + \frac{1}{2z} = 20.$$

$$15. \frac{9}{x} - \frac{2}{y} = \frac{5}{z} - \frac{3}{x} = \frac{7}{y} + \frac{15}{2z} = 4.$$

PROBLEMS

108. In the following problems, it is seen that if the number of unknown quantities is less than the number of problems, the problems will always be solvable. If the number of unknown quantities is equal to the number of problems, the problems will contain as many conditions as are necessary to be determined.

Example 1. A man has a certain number of pounds, the fifth of whose

Let x be the number of pounds.

Then

Also

or

By addition
The number of pounds is

Example 2. A man has a certain number of pounds, the fifth of whose is £3. 5s. 6d., and the sixth of whose is £4. 6s. 2d.; find the number of pounds.

Suppose x be the number of pounds, and

Then from the first condition,

Multiplying by 60,

Subtracting

And from the second condition, whence

$\therefore$ the number of pounds is 100, and the number of shillings is 100.

CHAPTER XIV.

PROBLEMS LEADING TO SIMULTANEOUS EQUATIONS.

108. IN the Examples discussed in the last chapter we have seen that it is essential to have as many equations as there are unknown quantities to determine. Consequently in the solution of problems which give rise to simultaneous equations, it will always be necessary that the statement of the question should contain as many independent conditions as there are quantities to be determined.

Example 1. Find two numbers whose difference is 11, and one-fifth of whose sum is 9.

Let x be the greater number, y the less ;

Then $x - y = 11$ (1),

Also $\frac{x+y}{5} = 9$,

or $x + y = 45$ (2).

By addition $2x = 56$; and by subtraction $2y = 34$.

The numbers are therefore 28 and 17.

Example 2. If 15 lbs. of tea and 17 lbs. of coffee together cost £3. 5s. 6d., and 25 lbs. of tea and 13 lbs. of coffee together cost £4. 6s. 2d. ; find the price of each per pound.

Suppose a pound of tea to cost x shillings,
and coffee y

Then from the question, we have

$$15x + 17y = 65\frac{1}{2} \text{(1),}$$

$$25x + 13y = 86\frac{1}{4} \text{(2).}$$

Multiplying (1) by 5 and (2) by 3, we have

$$75x + 85y = 327\frac{1}{2},$$

$$75x + 39y = 258\frac{1}{2}.$$

Subtracting,

$$46y = 69,$$

$$y = 1\frac{1}{2}.$$

And from (1)

$$15x + 25\frac{1}{2} = 65\frac{1}{2},$$

whence

$$15x = 40 ;$$

$$\therefore x = 2\frac{2}{3}.$$

$\therefore$ the cost of a pound of tea is $2\frac{2}{3}$ shillings, or 2s. 8d.,

and the cost of a pound of coffee is $1\frac{1}{2}$ shillings, or 1s. 6d.

Example 3. A person spent 15s. 2d. in buying oranges at the rate of 3 for twopence and apples at fivepence a dozen; if he had bought five times as many oranges and a quarter of the number of apples he would have spent £2. 4s. 2d. How many of each did he buy?

Let x be the number of oranges, and y the number of apples.

x oranges cost $\frac{2x}{3}$ pence,

y apples cost $\frac{5y}{12}$ pence,

$$\therefore \frac{2x}{3} + \frac{5y}{12} = 182 \dots\dots\dots(1).$$

Again, $5x$ oranges cost $5x \times \frac{2}{3}$, or $\frac{10x}{3}$ pence,

and $\frac{y}{4}$ apples cost $\frac{y}{4} \times \frac{5}{12}$, or $\frac{5y}{48}$ pence,

$$\therefore \frac{10x}{3} + \frac{5y}{48} = 530 \dots\dots\dots(2).$$

Multiply (1) by 5 and subtract (2) from the result;

$$\therefore \left(\frac{25}{12} - \frac{5}{48} \right) y = 380;$$

$$\therefore \frac{95y}{48} = 380;$$

$$\therefore y = 192,$$

$$x = 153.$$

and from (1)

Thus there were 153 oranges and 192 apples.

Example 4. If the numerator of a fraction is increased by 2 and the denominator by 1, it becomes equal to $\frac{5}{8}$; and, if the numerator and denominator are each diminished by 1, it becomes equal to $\frac{1}{2}$: find the fraction.

Let x be the numerator of the fraction, y the denominator; then the fraction is $\frac{x}{y}$.

From the first supposition,

$$\frac{x+2}{y+1} = \frac{5}{8} \dots\dots\dots(1),$$

from the second,

$$\frac{x-1}{y-1} = \frac{1}{2} \dots\dots\dots(2).$$

These equations give $x=8$, $y=15$.

Thus the fraction is $\frac{8}{15}$.

Example 5. is zero, and the number so

Let x be the

$y \dots\dots$

then, since the represented by

And if the represented by

or

that is,

Again, since we have

From (1) and

Hence the result

1. Find two

2. The sum find the number

3. One third their difference

4. One ninth difference is 30

5. Half the difference is 10

6. Six pounds and eleven pence Find the cost

7. Six hundred thirteen cows the value of each

8. A, B, C, C, and B has £50 less than

9. A, B, C much as C, and have £50 less has.

Example 5. The middle digit of a number between 100 and 1000 is zero, and the sum of the other digits is 11. If the digits be reversed, the number so formed exceeds the original number by 495; find it.

Let x be the digit in the units' place,

y hundreds' place;

then, since the digit in the tens' place is 0, the number will be represented by $100y + x$. [Art. 81, Ex. 4.]

And if the digits are reversed the number so formed will be represented by $100x + y$.

$$\therefore 100x + y - (100y + x) = 495,$$

or

$$100x + y - 100y - x = 495;$$

$$\therefore 99x - 99y = 495,$$

that is,

$$x - y = 5 \text{ (1),}$$

Again, since the sum of the digits is 11, and the middle one is 0, we have

$$x + y = 11 \text{ (2).}$$

From (1) and (2) we find $x = 8$, $y = 3$.

Hence the number is 308.

EXAMPLES XIV.

1. Find two numbers whose sum is 34, and whose difference is 10.
2. The sum of two numbers is 13, and their difference is 37; find the numbers.

3. One third of the sum of two numbers is 14, and one half of their difference is 4; find the numbers.

4. One nineteenth of the sum of two numbers is 4, and their difference is 30; find the numbers.

5. Half the sum of two numbers is 20, and three times their difference is 18; find the numbers.

6. Six pounds of tea and eleven pounds of sugar cost £1. 3s. 8d., and eleven pounds of tea and six pounds of sugar cost £1. 18s. 8d. Find the cost of tea and sugar per pound.

7. Six horses and seven cows can be bought for £250, and thirteen cows and eleven horses can be bought for £461. What is the value of each animal?

8. A , B , C , D have £290 between them; A has twice as much as C , and B has three times as much as D ; also C and D together have £50 less than A . Find how much each has.

9. A , B , C , D have £270 between them; A has three times as much as C , and B five times as much as D ; also A and B together have £50 less than eight times what C has. Find how much each has.

10. Four times B 's age exceeds A 's age by twenty years, and one third of A 's age is less than B 's age by two years: find their ages.
11. One eleventh of A 's age is greater by two years than one seventh of B 's, and twice B 's age is equal to what A 's age was thirteen years ago: find their ages.
12. In eight hours A walks twelve miles more than B does in seven hours; and in thirteen hours B walks seven miles more than A does in nine hours. How many miles does each walk per hour?
13. In eleven hours C walks $12\frac{1}{2}$ miles less than D does in twelve hours; and in five hours D walks $3\frac{1}{4}$ miles less than C does in seven hours. How many miles does each walk per hour?
14. Find a fraction such that if 1 be added to its denominator it reduces to $\frac{1}{2}$, and reduces to $\frac{3}{8}$ on adding 2 to its numerator.
15. Find a fraction which becomes $\frac{1}{2}$ on subtracting 1 from the numerator and adding 2 to the denominator, and reduces to $\frac{1}{3}$ on subtracting 7 from the numerator and 2 from the denominator.
16. If 1 be added to the numerator of a fraction it reduces to $\frac{1}{2}$; if 1 be taken from the denominator it reduces to $\frac{1}{7}$: required the fraction.
17. If $\frac{2}{3}$ be added to the numerator of a certain fraction the fraction will be increased by $\frac{1}{11}$, and if $\frac{1}{2}$ be taken from its denominator the fraction becomes $\frac{2}{9}$: find it.
18. The sum of a number of two digits and of the number formed by reversing the digits is 110, and the difference of the digits is 6: find the numbers.
19. The sum of the digits of a number is 13, and the difference between the number and that formed by reversing the digits is 27: find the numbers.
20. A certain number of two digits is three times the sum of its digits, and if 45 be added to it the digits will be reversed: find the number.
21. A certain number between 10 and 100 is eight times the sum of its digits, and if 45 be subtracted from it the digits will be reversed: find the number.
22. A man has a number of pounds and shillings, and he observes that if the pounds were turned into shillings and the shillings into pounds he would gain £5. 14s.; but if the pounds were turned into half-sovereigns and the shillings into half-crowns he would lose £1. 13s. 6d. What sum has he?
23. In a bag containing black and white balls, half the number of white is equal to a third of the number of black; and twice the whole number of balls exceeds three times the number of black balls by four. How many balls did the bag contain?

24. A number is diminished by 10, and the result is equal to the original number and right-hand digit. Find the number.

25. The velocity of a man together with each man and each woman.

26. A grocer has 5s. a pound quantity of each.

27. A train mile an hour time: had he $2\frac{1}{2}$ hours long.

28. A man and partly at 4, and vice same time.

29. Two people together in 9 hours if they walk in.

30. A family in food £1. 17s. afford only £1 one-half, and one adult and one child.

31. If I lend a certain time for a fourth of much do I lend.

32. A takes doubles his pace of walking.

24. A number consists of three digits, the right-hand one being zero. If the left-hand and middle digits be interchanged the number is diminished by 180; if the left-hand digit be halved and the middle and right-hand digits be interchanged the number is diminished by 454. Find the number.

25. The wages of 10 men and 8 boys amount to £1. 17s.; if 4 men together receive 1s. more than 6 boys, what are the wages of each man and boy?

26. A grocer wishes to mix spice at 8s. a pound with another sort at 5s. a pound to make 60 pounds to be sold at 6s. a pound: what quantity of each must he take?

27. A traveller walks a certain distance; had he gone half a mile an hour faster, he would have walked it in four-fifths of the time: had he gone half a mile an hour slower, he would have been $2\frac{1}{2}$ hours longer on the road. Find the distance.

28. A man walks 35 miles partly at the rate of 4 miles an hour, and partly at 5; if he had walked at 5 miles an hour when he walked at 4, and vice versa, he would have covered two miles more in the same time. Find the time he was walking.

29. Two persons, 27 miles apart, setting out at the same time are together in 9 hours if they walk in the same direction, but in 3 hours if they walk in opposite directions: find their rates of walking.

30. A family, consisting of three adults and five children, spends in food £1. 17s. 6d. a week. Distress, however, comes when they can afford only £1 per week, and the food of each adult is diminished by one-half, and of each child by one-third. Find the cost per week of an adult and of a child.

31. If I lend a sum of money at 6 per cent., the interest for a certain time exceeds the loan by £100; but if I lend it at 3 per cent., for a fourth of the time, the loan exceeds its interest by £425. How much do I lend?

32. A takes 3 hours longer than B to walk 30 miles; but if he doubles his pace he takes 2 hours less time than B: find their rates of walking.

It will be
nator are op

CHAPTER XV.

INVOLUTION.

109. DEFINITION. **Involution** is the general name for multiplying an expression by itself so as to find its second, third, fourth, or any other power.

Involution may always be effected by actual multiplication. Here, however, we shall give some rules for writing down at once

- (1) any power of a simple expression ;
- (2) the square and cube of any binomial ;
- (3) the square of any multinomial.

110. It is evident from the Rule of Signs that

- (1) no *even* power of *any* quantity can be *negative* ;
- (2) any *odd* power of a quantity will have *the same sign* as the quantity itself.

Note. It is especially worthy of notice that the *square* of every expression, whether positive or negative, is *positive*.

111. From definition we have, by the rules of multiplication,

$$\begin{aligned} (a^2)^3 &= a^2 \cdot a^2 \cdot a^2 = a^{2+2+2} = a^6. \\ (-x^3)^2 &= (-x^3)(-x^3) = x^{3+3} = x^6. \\ (-a^5)^3 &= (-a^5)(-a^5)(-a^5) = -a^{5+5+5} = -a^{15}. \\ (-3a^3)^4 &= (-3)^4(a^3)^4 = 81a^{12}. \end{aligned}$$

Hence we obtain a rule for raising a simple expression to any proposed power.

Rule. (1) *Raise the coefficient to the required power by Arithmetic, and prefix the proper sign found by the Rule of Signs.*

(2) *Multiply the index of every factor of the expression by the exponent of the power required.*

Write do

1. $3ab^3$.

5. $4a^4b^5x^2$.

9. $4xyz^3$.

13. $-\frac{7ab}{3}$.

17. $\frac{5ab^3}{2xy}$.

Write do

21. $2ab^2$.

25. $-5ab^3$.

29. $\frac{1}{3y^2}$.

Write do

33. $(3a^2b^3)^4$.

37. $\left(\frac{3x^4}{2y^3}\right)^5$.

112. By

These results

Rule 1.
sum of their

Rule 2.
to the sum of

Examp.

$$(-2x^3)^5 = -32x^{15}.$$

$$(-3ab^3)^6 = 729a^6b^{18}.$$

$$\left(\frac{2ab^3}{3x^2y}\right)^4 = \frac{16a^4b^{12}}{81x^8y^4}.$$

It will be seen that in the last case the numerator and the denominator are operated upon separately.

EXAMPLES XV. a.

Write down the square of each of the following expressions :

1. $3ab^3.$

2. $a^3c.$

3. $7ab^2.$

4. $11b^2c^3.$

5. $4a^4b^5x^2.$

6. $5x^2y^5.$

7. $-2abc^2.$

8. $-3cx^3.$

9. $4xyz^3.$

10. $-\frac{2}{3}a^3b^3.$

11. $\frac{2x^2}{3y^3}.$

12. $-\frac{4}{3x^2y}.$

13. $-\frac{7ab}{3}.$

14. $\frac{3a^2b^3}{4c^5x^4}.$

15. $-\frac{1}{2xy}.$

16. $-2xy^2.$

17. $\frac{5ab^3}{2xy}.$

18. $13c^5x^3.$

19. $-\frac{1}{4a^4}.$

20. $-\frac{3a^5}{5x^3}.$

Write down the cube of each of the following expressions :

21. $2ab^3.$

22. $3x^3.$

23. $4x^4.$

24. $-3a^3b.$

25. $-5ab^3.$

26. $-b^3c^2x.$

27. $-6a^6.$

28. $-2a^7c^2.$

29. $\frac{1}{3y^3}.$

30. $-\frac{3x^5}{5a^3}.$

31. $7x^3y^4.$

32. $-\frac{2}{3}a^5.$

Write down the value of each of the following expressions :

33. $(3a^2b^3)^4.$

34. $(-a^2x)^6.$

35. $(-2x^3y)^5.$

36. $\left(\frac{1}{2a^3}\right)^7.$

37. $\left(\frac{3x^4}{2y^3}\right)^5.$

38. $\left(\frac{2x^3}{3y}\right)^8.$

39. $\left(-\frac{x^3}{3}\right)^7.$

40. $\left(-\frac{2x^5}{3a^4}\right)^6.$

112. By multiplication we have

$$(a+b)^2 = (a+b)(a+b) \\ = a^2 + 2ab + b^2 \dots \dots \dots (1),$$

$$(a-b)^2 = (a-b)(a-b) \\ = a^2 - 2ab + b^2 \dots \dots \dots (2).$$

These results are embodied in the following rules :

Rule 1. *The square of the sum of two quantities is equal to the sum of their squares increased by twice their product.*

Rule 2. *The square of the difference of two quantities is equal to the sum of their squares diminished by twice their product.*

Example 1. $(x+2y)^2 = x^2 + 2 \cdot x \cdot 2y + (2y)^2$
 $= x^2 + 4xy + 4y^2.$

Example 2. $(2a^3 - 3b^2)^2 = (2a^3)^2 - 2 \cdot 2a^3 \cdot 3b^2 + (3b^2)^2$
 $= 4a^6 - 12a^3b^2 + 9b^4.$

113. These rules may sometimes be conveniently applied to find the squares of numerical quantities.

Example 1. The square of 1012 $= (1000 + 12)^2$
 $= (1000)^2 + 2 \cdot 1000 \cdot 12 + (12)^2$
 $= 1000000 + 24000 + 144$
 $= 1024144.$

Example 2. The square of 98 $= (100 - 2)^2$
 $= (100)^2 - 2 \cdot 100 \cdot 2 + (2)^2$
 $= 10000 - 400 + 4$
 $= 9604.$

The work is considerably shortened by the omission of the first two steps.

114. We may now extend the rules of Art 112 thus :

$$(a+b+c)^2 = \{(a+b)+c\}^2$$

$$= (a+b)^2 + 2(a+b)c + c^2 \quad [\text{Art. 112. Rule 1.}]$$

$$= a^2 + b^2 + c^2 + 2ab + 2ac + 2bc.$$

In the same way we may prove

$$(a-b+c)^2 = a^2 + b^2 + c^2 - 2ab + 2ac - 2bc$$

$$(a+b+c+d)^2 = a^2 + b^2 + c^2 + d^2 + 2ab + 2ac + 2ad + 2bc + 2bd + 2cd.$$

In each of these instances we observe that the square consists of

(1) the sum of the squares of the several terms of the given expression ;

(2) twice the sum of the products two and two of the several terms, taken with their proper signs ; that is, in each product the sign is + or - according as the quantities composing it have like or unlike signs.

Note. The square terms are always positive.

The same laws hold whatever be the number of terms in the expression to be squared.

Rule. To find the square of any multinominal : to the sum of the squares of the several terms add twice the product (with the proper sign) of each term into each of the terms that follow it.

Ex. 1. (

Ex. 2. (1

by co

Write do

1. $a + 3b.$

5. $3x - y.$

9. $pq - r.$

13. $a - b - c.$

16. $2a - 3b +$

19. $3p - 2q +$

22. $x - y + a$

25. $\frac{1}{2}a - 2b +$

115. By a

Also

By observ
results we ca

Example 1.

Example 2.

Write down

1. $x + a.$

5. $3x - 5y.$

9. $x^2 + 4y^2.$

13. $a - \frac{2b}{3}.$

$$\begin{aligned} \text{Ex. 1. } (x-2y-3z)^2 &= x^2 + 4y^2 + 9z^2 - 2 \cdot x \cdot 2y - 2 \cdot x \cdot 3z + 2 \cdot 2y \cdot 3z \\ &= x^2 + 4y^2 + 9z^2 - 4xy - 6xz + 12yz. \end{aligned}$$

$$\begin{aligned} \text{Ex. 2. } (1+2x-3x^2)^2 &= 1 + 4x^2 + 9x^4 + 2 \cdot 1 \cdot 2x - 2 \cdot 1 \cdot 3x^2 - 2 \cdot 2x \cdot 3x^2 \\ &= 1 + 4x^2 + 9x^4 + 4x - 6x^2 - 12x^3 \\ &= 1 + 4x - 2x^2 - 12x^3 + 9x^4, \end{aligned}$$

by collecting like terms and rearranging.

EXAMPLES XV. b.

Write down the square of each of the following expressions :

- | | | | |
|-------------------------------------|------------------------------------|--------------------------------------|---------------|
| 1. $a+3b$. | 2. $a-3b$. | 3. $x-5y$. | 4. $2x+3y$. |
| 5. $3x-y$. | 6. $3x+5y$. | 7. $9x-2y$. | 8. $5ab-c$. |
| 9. $pq-r$. | 10. $x-abc$. | 11. $ax+2by$. | 12. x^2-1 . |
| 13. $a-b-c$. | 14. $a+b-c$. | 15. $a+2b+c$. | |
| 16. $2a-3b+4c$. | 17. $x^2-y^2-z^2$. | 18. $xy+yz+zx$. | |
| 19. $3p-2q+4r$. | 20. x^2-x+1 . | 21. $2x^2+3x-1$. | |
| 22. $x-y+a-b$. | 23. $2x+3y+a-2b$. | 24. $m-n-p-q$. | |
| 25. $\frac{1}{2}a-2b+\frac{c}{4}$. | 26. $\frac{a}{3}-3b-\frac{3}{2}$. | 27. $\frac{2}{3}x^2-x+\frac{3}{2}$. | |

115. By actual multiplication we have

$$\begin{aligned} (a+b)^3 &= (a+b)(a+b)(a+b) \\ &= a^3 + 3a^2b + 3ab^2 + b^3. \end{aligned}$$

Also

$$(a-b)^3 = a^3 - 3a^2b + 3ab^2 - b^3.$$

By observing the law of formation of the terms in these results we can write down the cube of any binomial.

$$\begin{aligned} \text{Example 1. } (2x+y)^3 &= (2x)^3 + 3(2x)^2y + 3(2x)y^2 + y^3 \\ &= 8x^3 + 12x^2y + 6xy^2 + y^3. \end{aligned}$$

$$\begin{aligned} \text{Example 2. } (3x-2a)^3 &= (3x)^3 - 3(3x)^2(2a) + 3(3x)(2a)^2 - (2a)^3 \\ &= 27x^3 - 54x^2a + 36xa^2 - 8a^3. \end{aligned}$$

EXAMPLES XV. c.

Write down the cube of each of the following expressions :

- | | | | |
|------------------------|-----------------------|--------------------------|------------------------|
| 1. $x+a$. | 2. $x-a$. | 3. $x-2y$. | 4. $2x+y$. |
| 5. $3x-5y$. | 6. $ab+c$. | 7. $2ab-3c$. | 8. $5a-bc$. |
| 9. x^2+4y^2 . | 10. $4x^2-5y^2$. | 11. $2a^3-3b^2$. | 12. $5x^5-4y^4$. |
| 13. $a-\frac{2b}{3}$. | 14. $\frac{a}{3}+2$. | 15. $\frac{x^2}{3}-3x$. | 16. $\frac{a}{8}+2x$. |

EXAMPLES XVI. a.

Write down the square root of the following expressions :

- | | | | |
|----------------------------------|-----------------------------------|-------------------------------------|--|
| 1. $4a^2b^4$. | 2. $9x^4y^2$. | 3. $25x^4y^4$. | 4. $16a^4b^2c^4$. |
| 5. $81a^6b^8$. | 6. $100x^8$. | 7. $a^{20}b^{16}c^4$. | 8. $a^8b^2c^{12}$. |
| 9. $64x^6y^{18}$. | 10. $\frac{36}{a^{36}}$. | 11. $\frac{a^{16}b^8}{16}$. | 12. $\frac{289y^4}{25}$. |
| 13. $\frac{324x^{12}}{169y^8}$. | 14. $\frac{81a^{18}}{36b^{12}}$. | 15. $\frac{256x^2y^4}{289p^{14}}$. | 16. $\frac{400a^{40}b^{20}}{81x^{10}y^{18}}$. |

Write down the cube root of the following expressions :

- | | | | |
|--------------------------------|--------------------------------|-------------------------------------|------------------------------------|
| 17. $27a^6b^3c^3$. | 18. $-8a^{12}b^9$. | 19. $64x^6y^3z^{12}$. | 20. $-343a^{12}b^{18}$. |
| 21. $-\frac{x^{12}y^9}{125}$. | 22. $\frac{8x^9}{729y^{18}}$. | 23. $\frac{125a^3b^6}{216x^6y^9}$. | 24. $-\frac{27x^{27}}{64y^{63}}$. |

Write down the value of each of the following expressions :

- | | | |
|--|---|---|
| 25. $\sqrt[4]{(a^8x^{12})}$. | 26. $\sqrt[7]{(x^{14}y^{21})}$. | 27. $\sqrt[5]{(32x^5y^{10})}$. |
| 28. $\sqrt[6]{(729a^{18}b^6)}$. | 29. $\sqrt[8]{(256a^8x^6)}$. | 30. $\sqrt[5]{(-x^{10}y^{15})}$. |
| 31. $\sqrt[7]{\frac{128}{a^{63}b^{56}}}$. | 32. $\sqrt[10]{\frac{a^{80}x^{50}}{b^{100}}}$. | 33. $\sqrt[9]{\frac{a^{18}}{b^{27}c^{36}}}$. |

119. To find the square root of a compound expression.

Since the square of $a+b$ is $a^2+2ab+b^2$, we have to discover a process by which a and b , the terms of the root, can be found when $a^2+2ab+b^2$ is given.

The first term, a , is the square root of a^2 .

Arrange the terms according to powers of one letter a . The first term is a^2 , and its square root is a . Set this down as the first term of the required root. Subtract a^2 from the given expression and the remainder is $2ab+b^2$ or $(2a+b) \times b$.

Thus, b , the second term of the root, will be the quotient when the remainder is divided by $2a+b$.

This divisor consists of two terms :

1. The double of a , the term of the root already found.
2. b , the new term itself.

The work may be arranged as follows :

$$\begin{array}{r}
 a^2+2ab+b^2 \quad (a+b) \\
 \underline{a^2} \\
 2a+b \quad \left| \begin{array}{l} 2ab+b^2 \\ 2ab+b^2 \end{array} \right.
 \end{array}$$

Example 1. Find the square root of $9x^2 - 42xy + 49y^2$.

$$\begin{array}{r} 9x^2 - 42xy + 49y^2 \quad (3x - 7y) \\ 9x^2 \\ \hline 6x - 7y \quad \begin{array}{l} - 42xy + 49y^2 \\ - 42xy + 49y^2 \\ \hline \end{array} \end{array}$$

Explanation. The square root of $9x^2$ is $3x$, and this is the first term of the root.

By doubling this we obtain $6x$, which is the first term of the divisor. Divide $-42xy$, the first term of the remainder, by $6x$ and we get $-7y$, the new term in the root, which has to be annexed both to the root and divisor. Next multiply the complete divisor by $-7y$ and subtract the result from the first remainder. There is now no remainder and the root has been found.

The process can be extended so as to find the square root of any multinomial. The first two terms of the root will be obtained as before. When we have brought down the *second remainder*, the first part of the new divisor is obtained by doubling the terms of the root already found. We then divide the first term of the remainder by the first term of the new divisor, and set down the result as the next term in the root and in the divisor. We next multiply the complete divisor by the last term of the root and subtract the product from the last remainder. If there is now no remainder the root has been found; if there is a remainder we continue the process.

Example 2. Find the square root of

$$25x^2a^2 - 12xa^3 + 16x^4 + 4a^4 - 24x^2a.$$

Rearrange in descending powers of x .

$$\begin{array}{r} 16x^4 - 24x^3a + 25x^2a^2 - 12xa^3 + 4a^4 \quad (4x^2 - 3xa + 2a^2) \\ 16x^4 \\ \hline 8x^2 - 3xa \quad \begin{array}{l} - 24x^3a + 25x^2a^2 \\ - 24x^3a + 9x^2a^2 \\ \hline \end{array} \\ 8x^2 - 6xa + 2a^2 \quad \begin{array}{l} 16x^2a^2 - 12xa^3 + 4a^4 \\ 16x^2a^2 - 12xa^3 + 4a^4 \\ \hline \end{array} \end{array}$$

Explanation. When we have obtained two terms in the root, $4x^2 - 3xa$, we have a remainder

$$16x^2a^2 - 12xa^3 + 4a^4.$$

Double the terms of the root already found and place the result, $8x^2 - 6xa$, as the first part of the divisor. Divide $16x^2a^2$, the first term of the remainder, by $8x^2$, the first term of the divisor; we get $+2a^2$ which we annex both to the root and divisor. Now multiply the complete divisor by $2a^2$ and subtract. There is no remainder, and the root is found.

Find the

1. $x^2 + 4x$
3. $x^2 - 10x$
5. $81x^2 +$
7. $x^4 - 2x^2$
9. $a^4 - 2a^2$
11. $9x^4 - 12x^2$
13. $4a^4 + 4a^2$
15. $4x^2 + 9y^2$
16. $16x^2 + 1$
17. $x^4 - 22x^2$
18. $25x^4 - 3$
19. $4x^4 + 4x^2$
20. $6ab^2c -$
21. $-6b^2c^2 -$
22. $4x^4 + 9y^2$
23. $67x^2 + 4$
24. $1 - 4x +$
25. $6acx^5 +$

[If preferred taken after C

*120. W fractional to of the work

*121. Th expression c its reciproca

the order of

and the num

The reason

EXAMPLES XVI. b.

Find the square root of each of the following expressions :

1. $x^2 + 4x + 4y^2$.
2. $9a^2 + 12ab + 4b^2$.
3. $x^2 - 10xy + 25y^2$.
4. $4x^2 - 12xy + 9y^2$.
5. $81x^2 + 18xy + y^2$.
6. $25x^2 - 30xy + 9y^2$.
7. $x^4 - 2x^2y^2 + y^4$.
8. $1 - 2a^3 + a^6$.
9. $a^4 - 2a^3 + 3a^2 - 2a + 1$.
10. $4x^4 - 12x^3 + 20x^2 - 30x + 25$.
11. $9x^4 - 12x^3 - 2x^2 + 4x + 1$.
12. $x^4 - 4x^3 + 6x^2 - 4x + 1$.
13. $4a^4 + 4a^3 - 7a^2 - 4a + 4$.
14. $1 - 10x + 27x^2 - 10x^3 + x^4$.
15. $4x^3 + 9y^3 + 25z^3 + 12xy - 30yz - 20xz$.
16. $16x^6 + 16x^7 - 4x^8 - 4x^9 + x^{10}$.
17. $a^6 - 22x^4 + 34x^3 + 121x^2 - 374x + 289$.
18. $25x^4 - 30ax^3 + 49a^2x^2 - 24a^3x + 16a^4$.
19. $4x^4 + 4x^2y^2 - 12x^2z^2 + y^4 - 6y^2z^2 + 9z^4$.
20. $6ab^2c - 4a^2bc + a^2b^2 + 4a^2c^2 + 9b^2c^2 - 12abc^2$.
21. $-6b^2c^2 + 9c^4 + b^4 - 12c^2a^2 + 4a^4 + 4a^2b^2$.
22. $4x^4 + 9y^4 + 13x^2y^2 - 6xy^3 - 4x^2y$.
23. $67x^3 + 49 + 9x^4 - 70x - 30x^2$.
24. $1 - 4x + 10x^2 - 20x^3 + 25x^4 - 24x^5 + 16x^6$.
25. $6acx^5 + 4b^2x^4 + a^2x^{10} + 9c^3 - 12bcx^2 - 4abx^7$.

[If preferred, the remainder of this chapter may be postponed and taken after Chap. XXIV.]

*120. When the expression whose root is required contains fractional terms, we may proceed as before, the fractional part of the work being performed by the rules explained in Chap. XII.

*121. There is one important point to be observed when an expression contains powers of a certain letter and also powers of its reciprocal. Thus in the expression

$$2x + \frac{1}{x^2} + 4 + x^3 + \frac{5}{x} + 7x^2 + \frac{8}{x^3},$$

the order of *descending* powers is

$$x^3 + 7x^2 + 2x + 4 + \frac{5}{x} + \frac{1}{x^2} + \frac{8}{x^3};$$

and the numerical quantity 4 stands between x and $\frac{1}{x}$.

The reason for this arrangement will appear in Chap. XXX.

Example. Find the square root of $24 + \frac{16y^2}{x^3} - \frac{8x}{y} + \frac{x^2}{y^3} - \frac{32y}{x}$.

Arrange the expression in descending powers of y .

$$\frac{16y^2}{x^3} - \frac{32y}{x} + 24 - \frac{8x}{y} + \frac{x^2}{y^3} \left(\frac{4y}{x} - 4 + \frac{x}{y} \right)$$

$$\begin{array}{r} \frac{16y^2}{x^3} \\ \hline \frac{8y}{x} - 4 \\ \hline \frac{8y}{x} - 8 + \frac{x}{y} \\ \hline \end{array} \quad \begin{array}{r} -\frac{32y}{x} + 24 \\ -\frac{82y}{x} + 16 \\ \hline 8 - \frac{8x}{y} + \frac{x^2}{y^3} \\ 8 - \frac{8x}{y} + \frac{x^2}{y^3} \\ \hline \end{array}$$

Here the second term in the root, -4 , arises from division of $-\frac{32y}{x}$ by $\frac{8y}{x}$, and the third term, $\frac{x}{y}$, arises from division of 8 by $\frac{8y}{x}$;

$$\text{thus } 8 \div \frac{8y}{x} = 8 \times \frac{x}{8y} = \frac{x}{y}.$$

*EXAMPLES XVI. c.

Find the square root of each of the following expressions:

1. $\frac{x^2}{4} - 3x + 9.$
2. $4 - \frac{4x}{y} + \frac{x^2}{y^2}.$
3. $\frac{x^2}{25} + \frac{2xy}{5} + y^2.$
4. $\frac{x^2}{y^3} + \frac{10x}{y} + 25.$
5. $\frac{x^2}{4y^3} - \frac{2x}{y} + 4.$
6. $\frac{x^3}{y^3} - \frac{2ax}{by} + \frac{a^2}{b^3}.$
7. $\frac{64x^2}{9y^3} + \frac{32x}{3y} + 4.$
8. $\frac{9x^2}{25} - 2 + \frac{25}{9x^2}.$
9. $\frac{a^4}{64} + \frac{a^3}{8} - a + 1.$
10. $x^4 + 2x^3 - x + \frac{1}{4}.$
11. $-3a^3 + \frac{25}{9} + a^4 - 5a + \frac{67}{12}a^2.$
12. $x^4 - 2x + \frac{1}{9} + \frac{29}{3}x^2 - 6x^3.$
13. $\frac{a^4}{4} + \frac{a^3}{x} + \frac{a^2}{x^2} - ax - 2 + \frac{x^2}{a^2}.$
14. $x^4 - 2x^3 + \frac{3x^2}{2} - \frac{x}{2} + \frac{1}{16}.$
15. $\frac{x^4}{4} + 4x^2 + \frac{ax^3}{3} + \frac{a^2}{9} - 2x^3 - \frac{4ax}{3}.$

xvi.]

$$16. \frac{9a^2}{x^3} - 6\frac{a}{x} + 5.$$

$$17. 16m^4 + 32m^3 + 16m^2 + 4m + 1.$$

$$18. 4x^4 + 32x^3 + 64x^2 + 48x + 16.$$

*122. 2

Since the discover a p
be found w

The first
Arrange
the first ter
first term o
expression

Thus b,
when the re

This divis

(1) Three
found.

(2) Three
term b.

(3) The s

The work

$$3(a)^2$$

$$3 \times a$$

$$(b)^3$$

Example 1

$$3(2x)^2$$

$$3 \times 2x \times (-3)$$

$$(-3y)^2$$

$$16. \frac{9a^3}{x^4} - \frac{6a}{5x} + \frac{101}{25} - \frac{4x}{15a} + \frac{4x^2}{9a^3}.$$

$$17. 16m^4 + \frac{16}{3}m^2n + 8m^2 + \frac{4}{9}n^2 + \frac{4}{3}n + 1.$$

$$18. 4x^4 + 32x^2 + 96 + \frac{64}{x^4} + \frac{128}{x^2}.$$

*122. To find the cube root of a compound expression.

Since the cube of $a+b$ is $a^3+3a^2b+3ab^2+b^3$, we have to discover a process by which a and b , the terms of the root, can be found when $a^3+3a^2b+3ab^2+b^3$ is given.

The first term a is the cube root of a^3 .

Arrange the terms according to powers of one letter a ; then the first term is a^3 , and its cube root a . Set this down as the first term of the required root. Subtract a^3 from the given expression and the remainder is

$$3a^2b+3ab^2+b^3 \text{ or } (3a^2+3ab+b^2) \times b.$$

Thus b , the second term of the root, will be the quotient when the remainder is divided by $3a^2+3ab+b^2$.

This divisor consists of three terms :

(1) Three times the square of a , the term of the root already found.

(2) Three times the product of this first term a , and the new term b .

(3) The square of b .

The work may be arranged as follows :

$$\begin{array}{r} a^3+3a^2b+3ab^2+b^3 \ (a+b) \\ \underline{a^3} \\ 3(a)^2 = 3a^2 \\ 3 \times a \times b = +3ab \\ \underline{(b)^2 = +b^2} \\ 3a^2+3ab+b^2 \end{array}$$

Example 1. Find the cube root of $8x^3-36x^2y+54xy^2-27y^3$.

$$\begin{array}{r} 8x^3-36x^2y+54xy^2-27y^3 \ (2x-3y) \\ \underline{8x^3} \\ -36x^2y+54xy^2-27y^3 \\ \underline{-36x^2y+54xy^2-27y^3} \\ 0 \end{array}$$

Example 2.

Find the cube root of $27 + 108x + 90x^2 - 80x^3 - 60x^4 + 48x^5 - 8x^6$.

$$\begin{array}{r}
 3 \times (3)^2 = 27 \\
 3 \times 3 \times 4x = + 36x \\
 (4x)^2 = \frac{27 + 36x + 16x^2}{27 + 72x + 48x^2} \\
 3 \times (3 + 4x)^2 = 27 + 72x + 48x^2 \\
 3 \times (3 + 4x) \times (-2x^3) = -18x^2 - 24x^3 \\
 (-2x^3)^2 = \frac{27 + 72x + 30x^2 - 24x^3 + 4x^4}{27 + 108x + 90x^2 - 80x^3 - 60x^4 + 48x^5 - 8x^6}
 \end{array}$$

Explanation.

When we have obtained two terms in the root, $3 + 4x$, we have a remainder $-54x^2 - 144x^3 - 60x^4 + 48x^5 - 8x^6$.

Take 3 times the square of the root already found and place the result, $27 + 72x + 48x^2$, as the first part of the new divisor. Divide $-54x^2$, the first term of the remainder, by 27 , the first term of the divisor; this gives a new term of the root, $-2x^3$. To complete the divisor we take 3 times the product of $(3 + 4x)$ and $-2x^3$, and also the square of $-2x^3$. Now multiply the complete divisor by $-2x^3$ and subtract; there is no remainder and the root is found.

Find the

1. $a^3 + 3a^2$
3. $a^3x^3 - 3$
5. $64a^3 - 1$
6. $1 + 3x +$
7. $1 - 6x +$
8. $a^3 + 6a^2b - 8b^3$
9. $8a^3 - 36$
10. $y^6 - 3y^5$
11. $8a^6 + 12$
12. $27x^6 - 5$
13. $27x^6 - 2$
14. $24x^4y^2 +$
15. $216 + 34$

*123. W
terms occur

Example.

Arrange th

$$\begin{array}{l}
 3 \times \left(\frac{2}{x^2} \right) \\
 3 \times \frac{2}{x^2} \\
 (-3x)^2
 \end{array}$$

*EXAMPLES XVI. d.

Find the cube root of each of the following expressions :

1. $a^3 + 3a^2 + 3a + 1$.
2. $x^3 + 6x^2 + 12x + 8$.
3. $a^3x^3 - 3a^2x^2y^2 + 3axy^4 - y^6$.
4. $8m^3 - 12m^2 + 6m - 1$.
5. $64a^3 - 144a^2b + 108ab^2 - 27b^3$.
6. $1 + 3x + 6x^2 + 7x^3 + 6x^4 + 3x^5 + x^6$.
7. $1 - 6x + 21x^2 - 44x^3 + 63x^4 - 54x^5 + 27x^6$.
8. $a^3 + 6a^2b - 3a^2c + 12ab^2 - 12abc + 3ac^2 + 8b^3 - 12b^2c + 6bc^2 - c^3$.
9. $8a^6 - 36a^5 + 66a^4 - 63a^3 + 33a^2 - 9a + 1$.
10. $y^6 - 3y^5 + 6y^4 - 7y^3 + 6y^2 - 3y + 1$.
11. $8a^6 + 12x^6 - 30x^4 - 35x^3 + 45x^2 + 27x - 27$.
12. $27x^6 - 54x^5a + 117x^4a^2 - 116x^3a^3 + 117x^2a^4 - 54xa^5 + 27a^6$.
13. $27x^6 - 27x^5 - 18x^4 + 17x^3 + 6x^2 - 3x - 1$.
14. $24x^4y^2 + 96x^2y^4 - 6x^6y + x^6 - 96xy^6 + 64y^6 - 56x^3y^3$.
15. $216 + 342x^2 + 171x^4 + 27x^6 - 27x^5 - 109x^3 - 108x$.

*123. We add some examples of cube root where fractional terms occur in the given expressions.

Example. Find the cube root of $54 - 27x^3 + \frac{8}{x^6} - \frac{36}{x^3}$.

Arrange the expression in *ascending* powers of x .

$$\begin{array}{rcl}
 & & \frac{8}{x^6} - \frac{36}{x^3} + 54 - 27x^3 \left(\frac{2}{x^2} - 3x \right) \\
 & & \frac{8}{x^6} \\
 3 \times \left(\frac{2}{x^2} \right)^3 & = & \frac{12}{x^4} \\
 3 \times \frac{2}{x^2} \times (-3x) & = & -\frac{18}{x} \\
 (-3x)^2 & = & \frac{9x^2}{1} \\
 & & \frac{12}{x^4} - \frac{18}{x} + 9x^2 \\
 & & \frac{8}{x^6} - \frac{36}{x^3} + 54 - 27x^3
 \end{array}$$

***EXAMPLES XVI. e.**

Find the cube root of each of the following expressions :

1. $\frac{x^3}{8} - \frac{3x^2}{4} + \frac{3x}{2} - 1.$

2. $\frac{x^3}{27} + \frac{2x^2}{3} + 4x + 8.$

3. $8x^3 - 4x^2y^2 + \frac{2}{3}xy^4 - \frac{y^6}{27}.$

4. $\frac{27x^3}{64y^3} - \frac{27x^2}{8y^2} + \frac{9x}{y} - 8.$

5. $x^3 - 9x + \frac{27}{x} - \frac{27}{x^3}.$

6. $\frac{x^6}{y^3} - 6x^4 + 12x^2y^3 - 8y^6.$

7. $\frac{x^3}{y^3} + \frac{6x^2}{y^2} + \frac{9x}{y} - 4 - \frac{9y}{x} + \frac{6y^2}{x^2} - \frac{y^3}{x^3}.$

8. $\frac{x^3}{27} - \frac{x^2}{3} + 2x - 7 + \frac{18}{x} - \frac{27}{x^2} + \frac{27}{x^3}.$

9. $\frac{x^3}{a^3} - \frac{12x^2}{a^2} + \frac{54x}{a} - 112 + \frac{108a}{x} - \frac{48a^2}{x^2} + \frac{8a^3}{x^3}.$

10. $\frac{64a^3}{x^3} - \frac{192a^2}{x^2} + \frac{240a}{x} - 160 + \frac{60x}{a} - \frac{12x^2}{a^2} + \frac{x^3}{a^3}.$

11. $\frac{6b}{a} + \frac{6a}{b} - 7 + \frac{a^3}{b^3} - \frac{3a^2}{b^2} - \frac{3b^2}{a^2} + \frac{b^3}{a^3}.$

12. $\frac{60x^4}{y^4} - \frac{80x^3}{y^3} - \frac{90x^2}{y^2} + \frac{8x^6}{y^6} + \frac{108x}{y} - 27 + \frac{48x^5}{y^5}.$

***124.** Before leaving the subject of Evolution it may be useful to remark that the ordinary rules for extracting square and cube roots in Arithmetic are based upon the algebraical methods we have explained in the present chapter.

Example 1. Find the square root of 5329.

Since 5329 lies between 4900 and 6400, that is between $(70)^2$ and $(80)^2$, its square root consists of two figures and lies between 70 and 80. Hence, corresponding to a the first term of the root in the algebraical process of Art. 119, we here have 70.

The analogy between the algebraical and arithmetical methods will be seen by comparing the cases we give below.

$$\begin{array}{r} a^2 + 2ab + b^2 \quad (a+b) \\ a^2 \overline{) 2a+b} \\ \hline 2ab + b^2 \\ 2ab + b^2 \\ \hline \end{array}$$

$$\begin{array}{r} 5329 \quad (70+3=73) \\ 4900 \\ \hline 429 \\ 140+3=143 \quad \overline{) 429} \\ \hline 429 \end{array}$$

Example

Here 53
and $(300)^2$

2(

Example

Since 61
and $(90)^2$,
between 80

In Arith
also sever
would be
ence will h

Example 2. Find the square root of 53824.

Here 53824 lies between 40000 and 90000, that is between $(200)^2$ and $(300)^2$.

$$\begin{array}{r}
 \begin{array}{c} a \quad b \quad c \\ 53824 \end{array} \begin{array}{l} (200 + 30 + 2 = 232) \\ 40000 \\ \hline 13824 \\ 12900 \\ \hline 924 \\ 924 \\ \hline 0 \end{array} \\
 2a + b \dots\dots 400 + 30 = 430 \\
 2(a + b) + c \dots\dots 460 + 2 = 462
 \end{array}$$

Example 3. Find the cube root of 614125.

Since 614125 lies between 512000 and 729000, that is between $(80)^3$ and $(90)^3$, therefore its cube root consists of two figures and lies between 80 and 90.

$$\begin{array}{r}
 \begin{array}{c} a + b \\ 614125 \end{array} \begin{array}{l} (80 + 5 = 85) \\ 512000 \\ \hline 102125 \\ 102125 \\ \hline 0 \end{array} \\
 3a^2 = 3 \times (80)^2 = 19200 \\
 3 \times a \times b = 3 \times 80 \times 5 = 1200 \\
 b^2 = 5 \times 5 = 25 \\
 20425
 \end{array}$$

In Arithmetic the ciphers are usually omitted, and there are also several other modifications of the algebraical rules. It would be out of place to discuss these fully here: some reference will be made to them in Chap. **xxix**.

CHAPTER XVII.

RESOLUTION INTO FACTORS.

125. DEFINITION. When an algebraical expression is the product of two or more expressions each of these latter quantities is called a **factor** of it, and the determination of these quantities is called the **resolution** of the expression into its factors.

In this chapter we shall explain the principal rules by which the resolution of expressions into their component factors may be effected.

126. When each of the terms which compose an expression is divisible by a common factor, the expression may be simplified by dividing each term separately by this factor, and enclosing the quotient within brackets; the common factor being placed outside as a coefficient.

Example 1. The terms of the expression $3a^2 - 6ab$ have a common factor $3a$;

$$\therefore 3a^2 - 6ab = 3a(a - 2b).$$

Example 2. $5a^2bx^3 - 15abx^2 - 20b^2x^2 = 5bx^2(a^2x - 3a - 4b^2).$

EXAMPLES XVII. a.

Resolve into factors :

- | | | |
|------------------------------|-----------------------------------|----------------------------------|
| 1. $a^3 - ax.$ | 2. $x^3 - x^2.$ | 3. $2a - 2a^2.$ |
| 4. $a^2 - ab^2.$ | 5. $7p^2 + p.$ | 6. $8x - 2x^2.$ |
| 7. $5ax - 5a^2x^2.$ | 8. $3x^2 + x^5.$ | 9. $x^2 + xy.$ |
| 10. $x^3 - x^2y.$ | 11. $5x - 25x^2y.$ | 12. $15 + 25x^2.$ |
| 13. $16x + 64x^2y.$ | 14. $15a^3 - 225a^4.$ | 15. $54 - 81x.$ |
| 16. $10x^3 - 25x^4y.$ | 17. $3x^3 - x^2 + x.$ | 18. $6x^3 + 2x^4 + 4x^5.$ |
| 19. $x^3 - x^2y + xy^2.$ | 20. $3a^4 - 3a^3b + 6a^2b^2.$ | 21. $2x^2y^3 - 6x^2y^2 + 2xy^3.$ |
| 22. $6x^3 - 9x^2y + 12xy^2.$ | 23. $5x^5 - 10a^2x^3 - 15a^3x^2.$ | |
| 24. $7a - 7a^3 + 14a^4.$ | 25. $38a^3x^5 + 57a^4x^2.$ | |

CHAP. XVII.

127. An expression can be arranged

Example

Noticing the last two in one bracket

$$x^2 - ax + b.$$

Example 2

6a

Example 3

12a

Note. In the expression contains some different arrangements

12a

the same result factors of a product

Resolve into factors

1. $a^2 + ab + b^2$
3. $a^2c^2 + acd$
5. $2x + cx + 2$
7. $5a + ab + b^2$
9. $ax - bx - c$
11. $mx - my - z$
13. $2ax + ay + z$

F. A.

127. An expression may be resolved into factors *if the terms can be arranged in groups which have a compound factor common.*

Example 1. Resolve into factors $x^2 - ax + bx - ab$.

Noticing that the first two terms contain a common factor x , and the last two terms a common factor b , we enclose the first two terms in one bracket, and the last two in another. Thus

$$\begin{aligned} x^2 - ax + bx - ab &= (x^2 - ax) + (bx - ab) \\ &= x(x - a) + b(x - a) \\ &= (x - a) \text{ taken } x \text{ times plus } (x - a) \text{ taken } b \text{ times} \\ &= (x - a) \text{ taken } (x + b) \text{ times} \\ &= (x - a)(x + b). \end{aligned}$$

Example 2. Resolve into factors $6x^2 - 9ax + 4bx - 6ab$.

$$\begin{aligned} 6x^2 - 9ax + 4bx - 6ab &= (6x^2 - 9ax) + (4bx - 6ab) \\ &= 3x(2x - 3a) + 2b(2x - 3a) \\ &= (2x - 3a)(3x + 2b). \end{aligned}$$

Example 3. Resolve into factors $12a^2 - 4ab - 3ax^2 + bx^2$.

$$\begin{aligned} 12a^2 - 4ab - 3ax^2 + bx^2 &= (12a^2 - 4ab) - (3ax^2 - bx^2) \\ &= 4a(3a - b) - x^2(3a - b) \\ &= (3a - b)(4a - x^2). \end{aligned}$$

Note. In the first line of work it is sufficient to see that each pair contains some common factor. Thus, in the last example, by a different arrangement, we have

$$\begin{aligned} 12a^2 - 4ab - 3ax^2 + bx^2 &= (12a^2 - 3ax^2) - (4ab - bx^2) \\ &= 3a(4a - x^2) - b(4a - x^2) \\ &= (4a - x^2)(3a - b), \end{aligned}$$

the same result as before, since it is immaterial in what order the factors of a product are written.

EXAMPLES XVII. b.

Resolve into factors :

1. $a^2 + ab + ac + bc.$
2. $a^2 - ac + ab - bc.$
3. $a^2c^2 + acd + abc + bd.$
4. $a^2 + 3a + ac + 3c.$
5. $2x + cx + 2c + c^2.$
6. $x^2 - ax + 5x - 5a.$
7. $5a + ab + 5b + b^2.$
8. $ab - by - ay + y^2.$
9. $ax - bx - az + bz.$
10. $pr + qr - ps - qs.$
11. $mx - my - nx + ny.$
12. $mx - ma + nx - na.$
13. $2ax + ay + 2bx + by.$
14. $3ax - bx - 3ay + by.$

F. A.

H

Resolve into factors :

- | | |
|------------------------------------|--|
| 15. $6x^2 + 3xy - 2ax - ay.$ | 16. $mx - 2my - nx + 2ny.$ |
| 17. $ax^2 - 3bxy - axy + 3by^2.$ | 18. $x^2 + mxy - 4xy - 4my^2.$ |
| 19. $ax^2 + bx^2 + 2a + 2b.$ | 20. $x^2 - 3x - xy + 3y.$ |
| 21. $2x^4 - x^3 + 4x - 2.$ | 22. $3x^3 + 5x^2 + 3x + 5.$ |
| 23. $x^4 + x^3 + 2x + 2.$ | 24. $y^3 - y^2 + y - 1.$ |
| 25. $axy + bxy - az - bcz.$ | 26. $f^2x^2 + g^2x^2 - ag^2 - af^2.$ |
| 27. $2ax^2 + 3axy - 2bxy - 3by^2.$ | 28. $amx^2 + bmaxy - anxy - bny^2.$ |
| 29. $ax - bx + by + cy - cx - ay.$ | 30. $a^2x + abx + ac + aby + b^2y + bc.$ |

Trinomial Expressions.

128. Before proceeding to the next case of resolution into factors the student is advised to refer to Chap. v. Art. 44. Attention has there been drawn to the way in which, in forming the product of two binomials, the coefficients of the different terms combine so as to give a trinomial result. Thus, by Art. 44,

$$(x+5)(x+3) = x^2 + 8x + 15 \dots\dots\dots(1).$$

$$(x-5)(x-3) = x^2 - 8x + 15 \dots\dots\dots(2).$$

$$(x+5)(x-3) = x^2 + 2x - 15 \dots\dots\dots(3).$$

$$(x-5)(x+3) = x^2 - 2x - 15 \dots\dots\dots(4).$$

We now propose to consider the converse problem : namely, the resolution of a trinomial expression, similar to those which occur on the right-hand side of the above identities, into its component binomial factors.

By examining the above results, we notice that :

1. The first term of both the factors is x .
2. The *product* of the second terms of the two factors is equal to the *third term* of the trinomial ; e.g. in (2) above we see that 15 is the product of -5 and -3 ; while in (3) -15 is the product of $+5$ and -3 .
3. The *algebraic sum* of the second terms of the two factors is equal to the *coefficient of x* in the trinomial ; e.g. in (4) the sum of -5 and $+3$ gives -2 , the coefficient of x in the trinomial.

In applying these laws we will first consider a case where the *third term of the trinomial is positive*.

Example 1. Resolve into factors $x^2 + 11x + 24$.

The second terms of the factors must be such that their product is $+24$, and their sum $+11$. It is clear that they must be $+8$ and $+3$.

$$\therefore x^2 + 11x + 24 = (x+8)(x+3).$$

Example

The second term is $+24$, and it is easy to

Example

Example

Example

The second term is $+10a^2$, and

Note. In his results, b of the factors

Resolve in

1. $a^2 + 3a +$
4. $a^2 - 7a +$
7. $x^2 - 19x$
10. $x^2 - 21x$
13. $x^2 - 19x$
16. $x^2 + 20x$
19. $x^2 + 23x$
22. $a^2 + 30a$
25. $a^2 - 14ab$
27. $m^2 - 13m$
29. $x^2 - 23xy$
31. $x^4 + 8x^2 +$
33. $x^2y^2 - 16x$
35. $x^2y^2 + 34x$
37. $a^2 - 20ab$
39. $a^2 - 29ab$
41. $12 - 7x +$
43. $132 - 23x$
45. $130 + 31x$
- 204 - 29x

Example 2. Resolve into factors $x^2 - 10x + 24$.

The second terms of the factors must be such that their product is $+24$, and their sum -10 . Hence they must *both* be *negative*, and it is easy to see that they must be -6 and -4 .

$$\therefore x^2 - 10x + 24 = (x - 6)(x - 4).$$

Example 3. $x^3 - 18x + 81 = (x - 9)(x - 9)$
 $= (x - 9)^2$.

Example 4. $x^4 + 10x^2 + 25 = (x^2 + 5)(x^2 + 5)$
 $= (x^2 + 5)^2$.

Example 5. Resolve into factors $x^3 - 11ax + 10a^2$.

The second terms of the factors must be such that their product is $+10a^2$, and their sum $-11a$. Hence they must be $-10a$ and $-a$.

$$\therefore x^3 - 11ax + 10a^2 = (x - 10a)(x - a).$$

Note. In examples of this kind the student should always verify his results, by forming the product (*mentally*, as explained in Chap. v.) of the factors he has chosen.

EXAMPLES XVII. c.

Resolve into factors :

- | | | |
|--------------------------------|---------------------------------|-------------------------|
| 1. $a^2 + 3a + 2$. | 2. $a^2 + 2a + 1$. | 3. $a^2 + 7a + 12$. |
| 4. $a^2 - 7a + 12$. | 5. $x^2 - 11x + 30$. | 6. $x^2 - 15x + 56$. |
| 7. $x^2 - 19x + 90$. | 8. $x^2 + 13x + 42$. | 9. $x^2 - 21x + 110$. |
| 10. $x^2 - 21x + 108$. | 11. $x^2 - 21x + 80$. | 12. $x^2 + 21x + 90$. |
| 13. $x^2 - 19x + 84$. | 14. $x^2 - 19x + 78$. | 15. $x^2 - 18x + 45$. |
| 16. $x^2 + 20x + 96$. | 17. $x^2 - 26x + 165$. | 18. $x^2 - 21x + 104$. |
| 19. $x^2 + 23x + 102$. | 20. $a^2 - 24a + 95$. | 21. $a^2 - 32a + 256$. |
| 22. $a^2 + 30a + 225$. | 23. $a^2 + 54a + 729$. | 24. $a^2 - 38a + 361$. |
| 25. $a^2 - 14ab + 49b^2$. | 26. $a^2 + 5ab + 6b^2$. | |
| 27. $m^2 - 13mn + 40n^2$. | 28. $m^2 - 22mn + 105n^2$. | |
| 29. $x^2 - 23xy + 132y^2$. | 30. $x^2 - 26xy + 169y^2$. | |
| 31. $x^4 + 8x^2 + 7$. | 32. $x^4 + 9x^2y^2 + 14y^4$. | |
| 33. $x^2y^2 - 16xy + 39$. | 34. $x^2 + 49xy + 600y^2$. | |
| 35. $x^2y^2 + 34xy + 289$. | 36. $a^4b^4 + 37a^2b^2 + 300$. | |
| 37. $a^2 - 20abx + 75b^2x^2$. | 38. $x^2 + 43xy + 390y^2$. | |
| 39. $a^2 - 29ab + 54b^2$. | 40. $x^4 + 162x^2 + 6561$. | |
| 41. $12 - 7x + x^2$. | 42. $20 + 9x + x^2$. | |
| 43. $132 - 23x + x^2$. | 44. $88 + 19x + x^2$. | |
| 45. $130 + 31xy + x^2y^2$. | 46. $143 - 24xa + x^2a^2$. | |
| 47. $204 - 29x^2 + x^4$. | 48. $216 + 35x + x^2$. | |

129. Next consider a case where *the third term of the trinomial is negative.*

Example 1. Resolve into factors $x^2 + 2x - 35$.

The second terms of the factors must be such that their product is -35 , and their *algebraical sum* $+2$. Hence they must have *opposite* signs, and the greater of them must be *positive* in order to give its sign to their sum.

The required terms are therefore $+7$ and -5 .

$$\therefore x^2 + 2x - 35 = (x + 7)(x - 5).$$

Example 2. Resolve into factors $x^2 - 3x - 54$.

The second terms of the factors must be such that their product is -54 , and their *algebraical sum* -3 . Hence they must have *opposite* signs, and the greater of them must be *negative* in order to give its sign to their sum.

The required terms are therefore -9 and $+6$.

$$\therefore x^2 - 3x - 54 = (x - 9)(x + 6).$$

Remembering that in these cases the numerical quantities *must have opposite signs*, if preferred, the following method may be adopted.

Example 3. Resolve into factors $x^2y^2 + 23xy - 420$.

Find two numbers whose product is 420 , and whose *difference* is 23 . These are 35 and 12 ; hence inserting the signs so that the positive may predominate, we have

$$x^2y^2 + 23xy - 420 = (xy + 35)(xy - 12).$$

EXAMPLES XVII. d.

Resolve into factors :

- | | | |
|---------------------------|--------------------------|-----------------------------|
| 1. $x^2 - x - 2$. | 2. $x^2 + x - 2$. | 3. $x^2 - x - 6$. |
| 4. $x^2 + x - 6$. | 5. $x^2 - 2x - 3$. | 6. $x^2 + 2x - 3$. |
| 7. $x^2 + x - 56$. | 8. $x^2 + 3x - 40$. | 9. $x^2 - 4x - 12$. |
| 10. $a^2 - a - 20$. | 11. $a^2 - 4a - 21$. | 12. $a^2 + a - 20$. |
| 13. $a^2 - 4a - 117$. | 14. $x^2 + 9x - 36$. | 15. $x^2 + x - 156$. |
| 16. $x^2 + x - 110$. | 17. $x^2 - 9x - 90$. | 18. $x^2 - x - 240$. |
| 19. $a^2 - 12a - 85$. | 20. $a^2 - 11a - 152$. | 21. $x^2y^2 - 5xy - 24$. |
| 22. $x^2 + 7xy - 60y^2$. | 23. $x^2 + ax - 42a^2$. | 24. $x^2 - 32xy - 105y^2$. |
| 25. $a^2 - ay - 210y^2$. | 26. $x^2 + 18x - 115$. | 27. $x^2 - 20xy - 96y^2$. |
| 28. $x^2 + 16x - 260$. | 29. $a^2 - 11a - 26$. | 30. $a^2y^2 + 14ay - 240$. |

[XVII.]

31. $a^4 - a^3b$

34. $a^2b^2 - 3a$

36. $a^2 - 18a$

38. $x^4 - a^2x^2$

40. $x^6 + x^3 -$

43. $110 - x -$

46. $65 + 8xy$

[Ex

130. We
nomial expres
unity.

Again, ref
following res

The conver
we have yet o

Before end
will be worth
given above.

Consider th

The first ter

The third t

The middle
two products

Again, consi

The first ter

The third t

The middle
two products
the greater of

131. The b
select the prop
enable him to
will combine so
sion to be resol

31. $a^4 - a^2b^2 - 56b^4$. 32. $x^4 - 14x^2 - 51$. 33. $y^4 + 6x^2y^2 - 27x^4$.
 34. $a^3b^2 - 3abc - 10c^2$. 35. $a^2 + 12abx - 28b^2x^2$.
 36. $a^3 - 18axy - 243x^2y^2$. 37. $x^4 + 13a^2x^2 - 300a^4$.
 38. $x^4 - a^2x^2 - 132a^4$. 39. $x^4 - a^2x^2 - 462a^4$.
 40. $x^2 + x^3 - 870$. 41. $2 + x - x^2$. 42. $6 + x - x^2$.
 43. $110 - x - x^2$. 44. $380 - x - x^2$. 45. $120 - 7ax - a^2x^2$.
 46. $65 + 8xy - x^2y^2$. 47. $98 - 7x - x^2$. 48. $204 - 5x - x^2$.

[Examples for revision will be found on page 174.]

130. We proceed now to the resolution into factors of trinomial expressions when the coefficient of the highest power is not unity.

Again, referring to Chap. v. Art. 44, we may write down the following results :

$$(3x+2)(x+4)=3x^2+14x+8\ldots\ldots(1),$$

$$(3x-2)(x-4)=3x^2-14x+8\ldots\ldots(2),$$

$$(3x+2)(x-4)=3x^2-10x-8\ldots\ldots(3),$$

$$(3x-2)(x+4)=3x^2+10x-8\ldots\ldots(4).$$

The converse problem presents more difficulty than the cases we have yet considered.

Before endeavouring to give a general method of procedure, it will be worth while to examine in detail two of the identities given above.

Consider the result $3x^2 - 14x + 8 = (3x - 2)(x - 4)$.

The first term $3x^2$ is the product of $3x$ and x .

The third term $+8$ -2 and -4 .

The middle term $-14x$ is the result of adding together the two products $3x \times -4$ and $x \times -2$.

Again, consider the result $3x^2 - 10x - 8 = (3x + 2)(x - 4)$.

The first term $3x^2$ is the product of $3x$ and x .

The third term -8 $+2$ and -4 .

The middle term $-10x$ is the result of adding together the two products $3x \times -4$ and $x \times 2$; and its sign is negative because the greater of these two products is negative.

131. The beginner will frequently find that it is not easy to select the proper factors at the first trial. Practice alone will enable him to detect at a glance whether any pair he has chosen will combine so as to give the correct coefficients of the expression to be resolved.

Example. Resolve into factors $7x^2 - 19x - 6$.

Write down $(7x \ 3)(x \ 2)$ for a first trial, noticing that 3 and 2 must have opposite signs. These factors give $7x^2$ and -6 for the first and third terms. But since $7 \times 2 - 3 \times 1 = 11$, the combination fails to give the correct coefficient of the middle term.

Next try $(7x \ 2)(x \ 3)$.

Since $7 \times 3 - 2 \times 1 = 19$, these factors will be correct if we insert the signs so that the negative shall predominate.

Thus $7x^2 - 19x - 6 = (7x + 2)(x - 3)$.

[Verify by mental multiplication.]

132. In actual work it will not be necessary to put down all these steps at length. The student will soon find that the different cases may be rapidly reviewed, and the unsuitable combinations rejected at once.

It is especially important to pay attention to the two following hints:

- (1) If the third term of the trinomial is positive, then the second terms of its factors have both the same sign, and this sign is the same as that of the middle term of the trinomial.
- (2) If the third term of the trinomial is negative, then the second terms of its factors have opposite signs.

Example 1. Resolve into factors $14x^2 + 29x - 15$ (1),
 $14x^2 - 29x - 15$ (2).

In each case we may write down $(7x \ 3)(2x \ 5)$ as a first trial, noticing that 3 and 5 must have opposite signs.

And since $7 \times 5 - 3 \times 2 = 29$, we have only now to insert the proper signs in each factor.

In (1) the positive sign must predominate,
 in 2 the negative

Therefore $14x^2 + 29x - 15 = (7x - 3)(2x + 5)$.

$14x^2 - 29x - 15 = (7x + 3)(2x - 5)$.

Example 2. Resolve into factors $5x^2 + 17x + 6$ (1),
 $5x^2 - 17x + 6$ (2).

In (1) we notice that the factors which give 6 are both positive.

In (2) negative.

And therefore for (1) we may write $(5x + \quad)(x + \quad)$.

(2) $(5x - \quad)(x - \quad)$.

And, since $5 \times 3 + 1 \times 2 = 17$, we see that

$5x^2 + 17x + 6 = (5x + 2)(x + 3)$.

$5x^2 - 17x + 6 = (5x - 2)(x - 3)$.

Note. In 6 and 1; b student wou

Example

Example

Resolve i

1. $2x^2 + 3$

4. $3x^2 + 1$

7. $2x^2 + 7$

10. $5x^2 + 1$

13. $4x^2 + 1$

16. $2x^2 - x$

19. $3x^2 + 1$

22. $3x^2 + 7$

25. $3x^2 + 1$

28. $4x^2 + x$

31. $4x^2 + 2$

34. $12x^2 - 2$

37. $12x^2 - 3$

40. $24x^2 - 2$

43. $6 + 5x$

46. $7 + 10x$

49. $20 - 9x$

133. By

a result wh

The prod
 is equal to t

Conversel
 is equal to
 quantities.

Thus any
 may at once

Note. In each expression the third term 6 also admits of factors 6 and 1; but this is one of the cases referred to above which the student would reject at once as unsuitable.

$$\text{Example 3. } 9x^2 - 48xy + 64y^2 = (3x - 8y)(3x - 8y) \\ = (3x - 8y)^2.$$

$$\text{Example 4. } 6 + 7x - 5x^2 = (3 + 5x)(2 - x).$$

EXAMPLES XVII. e. ✓

Resolve into factors :

- | | | |
|-----------------------------|--------------------------|--------------------------|
| 1. $2x^2 + 3x + 1.$ | 2. $3x^2 + 5x + 2.$ | 3. $2x^2 + 5x + 2.$ |
| 4. $3x^2 + 10x + 3.$ | 5. $2x^2 + 9x + 4.$ | 6. $3x^2 + 8x + 4.$ |
| 7. $2x^2 + 7x + 6.$ | 8. $2x^2 + 11x + 5.$ | 9. $3x^2 + 11x + 6.$ |
| 10. $5x^2 + 11x + 2.$ | 11. $2x^2 + 3x - 2.$ | 12. $3x^2 + x - 2.$ |
| ✓ 13. $4x^2 + 11x - 3.$ | 14. $3x^2 + 14x - 5.$ | 15. $2x^2 + 15x - 8.$ |
| 16. $2x^2 - x - 1.$ | 17. $3x^2 + 7x - 6.$ | 18. $2x^2 + x - 28.$ |
| 19. $3x^2 + 13x - 30.$ | 20. $6x^2 + 7x - 3.$ | 21. $6x^2 - 7x - 3.$ |
| 22. $3x^2 + 7x + 4.$ | 23. $3x^2 + 23x + 14.$ | 24. $2x^2 - x - 15.$ |
| 25. $3x^2 + 19x - 14.$ | 26. $3x^2 - 19x - 14.$ | 27. $6x^2 - 31x + 35.$ |
| 28. $4x^2 + x - 14.$ | 29. $3x^2 - 13x + 14.$ | 30. $3x^2 + 41x + 26.$ |
| 31. $4x^2 + 23x + 15.$ | 32. $2x^2 - 5xy - 3y^2.$ | 33. $8x^2 - 38x + 35.$ |
| 34. $12x^2 - 23xy + 10y^2.$ | 35. $15x^2 + 224x - 15.$ | 36. $15x^2 - 77x + 10.$ |
| 37. $12x^2 - 31x - 15.$ | 38. $24x^2 + 22x - 21.$ | 39. $72x^2 - 145x + 72.$ |
| 40. $24x^2 - 29xy - 4y^2.$ | 41. $2 - 3x - 2x^2.$ | 42. $3 + 11x - 4x^2.$ |
| 43. $6 + 5x - 6x^2.$ | 44. $4 - 5x - 6x^2.$ | 45. $5 + 32x - 21x^2.$ |
| 46. $7 + 10x + 3x^2.$ | 47. $18 - 33x + 5x^2.$ | 48. $8 + 6x - 5x^2.$ |
| 49. $20 - 9x - 20x^2.$ | 50. $24 + 37x - 72x^2.$ | |

The Difference of Two Squares.

133. By multiplying $a + b$ by $a - b$ we obtain the identity

$$(a + b)(a - b) = a^2 - b^2,$$

a result which may be verbally expressed as follows :

The product of the sum and the difference of any two quantities is equal to the difference of their squares.

Conversely, the difference of the squares of any two quantities is equal to the product of the sum and the difference of the two quantities.

Thus any expression which is the difference of two squares may at once be resolved into factors.

Example. Resolve into factors $25x^2 - 16y^2$.

$$25x^2 - 16y^2 = (5x)^2 - (4y)^2.$$

Therefore the first factor is the sum of $5x$ and $4y$,
and the second factor is the difference of $5x$ and $4y$.

$$\therefore 25x^2 - 16y^2 = (5x + 4y)(5x - 4y).$$

The intermediate steps may usually be omitted.

Example. $1 - 49c^6 = (1 + 7c^3)(1 - 7c^3)$.

The difference of the squares of two numerical quantities may be found by the formula $a^2 - b^2 = (a + b)(a - b)$.

$$\begin{aligned} \text{Example. } (329)^2 - (171)^2 &= (329 + 171)(329 - 171) \\ &= 500 \times 158 \\ &= 79000. \end{aligned}$$

EXAMPLES XVII. f.

Resolve into factors :

- | | | |
|--------------------------|-----------------------------|--------------------------|
| 1. $x^2 - 4$. | 2. $a^2 - 81$. | 3. $y^2 - 100$. |
| 4. $c^2 - 144$. | 5. $9 - a^2$. | 6. $49 - c^2$. |
| 7. $121 - x^2$. | 8. $400 - a^2$. | 9. $x^2 - 9a^2$. |
| 10. $y^2 - 25x^2$. | 11. $36x^2 - 25b^2$. | 12. $9x^2 - 1$. |
| 13. $36p^2 - 49q^2$. | 14. $4k^2 - 1$. | 15. $49 - 100k^2$. |
| 16. $1 - 25x^2$. | 17. $a^2 - 4b^2$. | 18. $9x^2 - y^2$. |
| 19. $p^2q^2 - 36$. | 20. $a^2b^2 - 4c^2d^2$. | 21. $x^4 - 9$. |
| 22. $9a^4 - 121$. | 23. $25x^2 - 64$. | 24. $81a^4 - 49x^4$. |
| 25. $x^6 - 25$. | 26. $1 - 36a^6$. | 27. $9x^4 - a^2$. |
| 28. $81x^6 - 25a^2$. | 29. $x^4a^2 - 49$. | 30. $a^2 - 64x^6$. |
| 31. $a^2b^2 - 9x^6$. | 32. $x^6y^6 - 4$. | 33. $1 - a^2b^2$. |
| 34. $4 - x^2$. | 35. $9 - 4a^2$. | 36. $9a^4 - 25b^4$. |
| 37. $x^4 - 16b^2$. | 38. $x^2 - 25y^2$. | 39. $1 - 100b^2$. |
| 40. $25 - 64x^2$. | 41. $121a^2 - 81x^2$. | 42. $p^2q^2 - 64a^4$. |
| 43. $64x^2 - 25z^6$. | 44. $49x^4 - 16y^4$. | 45. $81p^2q^6 - 25b^2$. |
| 46. $16x^{16} - 9y^6$. | 47. $36x^{36} - 49a^{14}$. | 48. $1 - 100a^6b^2c^2$. |
| 49. $25x^{10} - 16a^8$. | 50. $a^2b^4c^8 - x^{16}$. | |

Find by resolving into factors the value of

- | | | |
|----------------------------|-----------------------------|----------------------------|
| 51. $(575)^2 - (425)^2$. | 52. $(121)^2 - (120)^2$. | 53. $(750)^2 - (250)^2$. |
| 54. $(339)^2 - (319)^2$. | 55. $(753)^2 - (253)^2$. | 56. $(101)^2 - (99)^2$. |
| 57. $(1723)^2 - (277)^2$. | 58. $(1639)^2 - (739)^2$. | 59. $(1811)^2 - (689)^2$. |
| 60. $(2731)^2 - (269)^2$. | 61. $(8133)^2 - (8131)^2$. | 62. $(10001)^2 - 1$. |

134. W
tity the sa

Example

The sum
and

Example

The sum
and

If the fac
to give the

Example

Resolve i

1. $(a + b)^2$
4. $(x + 2y)^2$
7. $(x + 5c)^2$
10. $a^2 - (b - c)^2$
13. $9x^2 - (2 - 3y)^2$
16. $(a + b)^2$
19. $(a + b)^2$
22. $(4a + x)^2$
24. $1 - (7a - b)^2$
26. $(a - 3x)^2$
28. $(a + b - c)^2$

Resolve i

30. $(x + y)^2$
33. $(24x + y)^2$
35. $9x^2 - (3 - 2y)^2$
37. $(3a + 1)^2$
39. $(2a + b)^2$
41. $(x + y - z)^2$

134. When one or both of the squares is a compound quantity the same method is employed.

Example 1. Resolve into factors $(a+2b)^2 - 16x^2$.

The sum of $a+2b$ and $4x$ is $a+2b+4x$,

and their difference is $a+2b-4x$.

$$\therefore (a+2b)^2 - 16x^2 = (a+2b+4x)(a+2b-4x).$$

Example 2. Resolve into factors $x^2 - (2b-3c)^2$.

The sum of x and $2b-3c$ is $x+2b-3c$,

and their difference is $x-(2b-3c)=x-2b+3c$,

$$\therefore x^2 - (2b-3c)^2 = (x+2b-3c)(x-2b+3c).$$

If the factors contain like terms they should be collected so as to give the result in its simplest form

$$\begin{aligned} \text{Example 3. } (3x+7y)^2 - (2x-3y)^2 \\ &= \{(3x+7y) + (2x-3y)\} \{(3x+7y) - (2x-3y)\} \\ &= (3x+7y+2x-3y)(3x+7y-2x+3y) \\ &= (5x+4y)(x+10y). \end{aligned}$$

EXAMPLES XVII. g.

Resolve into factors :

- | | | |
|-------------------------------|--------------------------------|---------------------------|
| 1. $(a+b)^2 - c^2$. | 2. $(a-b)^2 - c^2$. | 3. $(x+y)^2 - 4z^2$. |
| 4. $(x+2y)^2 - a^2$. | 5. $(a+3b)^2 - 16x^2$. | 6. $(x+5a)^2 - 9y^2$. |
| 7. $(x+5c)^2 - 1$. | 8. $(a-2x)^2 - b^2$. | 9. $(2x-3a)^2 - 9c^2$. |
| 10. $a^2 - (b-c)^2$. | 11. $x^2 - (y+z)^2$. | 12. $4a^2 - (y-z)^2$. |
| 13. $9x^2 - (2a-3b)^2$. | 14. $1 - (a-b)^2$. | 15. $c^2 - (5a-3b)^2$. |
| 16. $(a+b)^2 - (c+d)^2$. | 17. $(a-b)^2 - (x+y)^2$. | 18. $(7x+y)^2 - 1$. |
| 19. $(a+b)^2 - (m-n)^2$. | 20. $(a-n)^2 - (b+m)^2$. | 21. $(b-c)^2 - (a-x)^2$. |
| 22. $(4a+x)^2 - (b+y)^2$. | 23. $(a+2b)^2 - (3x+4y)^2$. | |
| 24. $1 - (7a-3b)^2$. | 25. $(a-b)^2 - (x-y)^2$. | |
| 26. $(a-3x)^2 - 16y^2$. | 27. $(2a-5x)^2 - 1$. | |
| 28. $(a+b-c)^2 - (x-y+z)^2$. | 29. $(3a+2b)^2 - (c+x-2y)^2$. | |

Resolve into factors and simplify :

- | | | |
|--------------------------------|-------------------------------|-------------------------|
| 30. $(x+y)^2 - x^2$. | 31. $x^2 - (y-x)^2$. | 32. $(x+3y)^2 - 4y^2$. |
| 33. $(24x+y)^2 - (23x-y)^2$. | 34. $(5x+2y)^2 - (3x-y)^2$. | |
| 35. $9x^2 - (3x-5y)^2$. | 36. $(7x+3)^2 - (5x-4)^2$. | |
| 37. $(3a+1)^2 - (2a-1)^2$. | 38. $16a^2 - (3a+1)^2$. | |
| 39. $(2a+b-c)^2 - (a-b+c)^2$. | 40. $(x-7y+z)^2 - (7y-z)^2$. | |
| 41. $(x+y-8)^2 - (x-8)^2$. | 42. $(2x+a-3)^2 - (3-2x)^2$. | |

135. By suitably grouping together the terms, compound expressions can often be expressed as the difference of two squares, and so be resolved into factors.

Example 1. Resolve into factors $a^2 - 2ax + x^2 - 4b^2$.

$$\begin{aligned} a^2 - 2ax + x^2 - 4b^2 &= (a^2 - 2ax + x^2) - 4b^2 \\ &= (a - x)^2 - (2b)^2 \\ &= (a - x + 2b)(a - x - 2b). \end{aligned}$$

Example 2. Resolve into factors $9a^2 - c^2 + 4cx - 4x^2$.

$$\begin{aligned} 9a^2 - c^2 + 4cx - 4x^2 &= 9a^2 - (c^2 - 4cx + 4x^2) \\ &= (3a)^2 - (c - 2x)^2 \\ &= (3a + c - 2x)(3a - c + 2x). \end{aligned}$$

Example 3. Resolve into factors $2bd - a^2 - c^2 + b^2 + d^2 + 2ac$.

Here the terms $2bd$ and $2ac$ suggest the proper preliminary arrangement of the expression. Thus

$$\begin{aligned} 2bd - a^2 - c^2 + b^2 + d^2 + 2ac &= b^2 + 2bd + d^2 - a^2 + 2ac - c^2 \\ &= b^2 + 2bd + d^2 - (a^2 - 2ac + c^2) \\ &= (b + d)^2 - (a - c)^2 \\ &= (b + d + a - c)(b + d - a + c). \end{aligned}$$

Example 4. Resolve into factors $x^4 + x^2y^2 + y^4$.

$$\begin{aligned} x^4 + x^2y^2 + y^4 &= (x^4 + 2x^2y^2 + y^4) - x^2y^2 \\ &= (x^2 + y^2)^2 - (xy)^2 \\ &= (x^2 + y^2 + xy)(x^2 + y^2 - xy) \\ &= (x^2 + xy + y^2)(x^2 - xy + y^2). \end{aligned}$$

This result is very important and will be referred to again in Chapter XXVIII.

EXAMPLES XVII. h.

Resolve into factors :

- | | |
|--|--|
| 1. $x^2 + 2xy + y^2 - a^2$. | 2. $a^2 - 2ab + b^2 - x^2$. |
| 3. $x^2 - 6ax + 9a^2 - 16b^2$. | 4. $4a^2 + 4ab + b^2 - 9c^2$. |
| 5. $x^2 + a^2 + 2ax - y^2$. | 6. $2ay + a^2 + y^2 - x^2$. |
| 7. $x^2 - a^2 - 2ab - b^2$. | 8. $y^2 - c^2 + 2cx - x^2$. |
| 9. $1 - x^2 - 2xy - y^2$. | 10. $c^2 - x^2 - y^2 + 2xy$. |
| 11. $x^2 + y^2 + 2xy - 4x^2y^2$. | 12. $a^2 - 4ab + 4b^2 - 9a^2c^2$. |
| 13. $x^2 + 2xy + y^2 - a^2 - 2ab - b^2$. | 14. $a^2 - 2ab + b^2 - c^2 - 2cd - d^2$. |
| 15. $x^2 - 4ax + 4a^2 - b^2 + 2by - y^2$. | 16. $y^2 + 2by + b^2 - a^2 - 6ax - 9x^2$. |

17. $x^2 - 2x$

19. $x^2 - a^2$

21. $4x^2 - 1$

22. $a^2 + 6b$

23. $a^4 - 25$

24. $x^4 - x^2$

25. $a^4 + a^2b^2$

28. $c^4 + 3c^2$

136. If
and if we d
We have

These re
which can

Example

Note. T

Example

We may
the factors

Examples

Resolve

1. $x^3 - y^3$

5. $8x^3 - y^3$

9. $a^3b^3 - c^3$

13. $125 + a$

17. $x^2 - 2x + 1 - a^2 - 4ab - 4b^2$. 18. $9a^2 - 6a + 1 - x^2 - 8dx - 16d^2$.
 19. $x^2 - a^2 + y^2 - b^2 - 2xy + 2ab$. 20. $a^2 + b^2 - 2ab - c^2 - d^2 - 2cd$.
 21. $4x^2 - 12ax - c^2 - k^2 - 2ck + 9a^2$.
 22. $a^2 + 6bx - 9b^2x^2 - 10ab - 1 + 25b^2$.
 23. $a^4 - 25x^2 + 8a^2x^2 - 9 + 30x^3 + 16x^4$.
 24. $x^4 - x^2 - 9 - 2a^2x^2 + a^4 + 6x$.
 25. $a^4 + a^2b^2 + b^4$. 26. $x^4 + 4x^2y^2 + 16y^4$. 27. $p^4 + 9p^2q^2 + 81q^4$.
 28. $c^4 + 3c^2d^2 + 4d^4$. 29. $x^4 + y^4 - 11x^2y^2$. 30. $4m^4 - 5m^2n^2 + n^4$.

The Sum or Difference of Two Cubes.

136. If we divide $a^3 + b^3$ by $a + b$ the quotient is $a^2 - ab + b^2$; and if we divide $a^3 - b^3$ by $a - b$ the quotient is $a^2 + ab + b^2$.

We have therefore the following identities :

$$\begin{aligned} a^3 + b^3 &= (a + b)(a^2 - ab + b^2); \\ a^3 - b^3 &= (a - b)(a^2 + ab + b^2). \end{aligned}$$

These results enable us to resolve into factors any expression which can be written as the sum or the difference of two cubes.

Example 1. $8x^3 - 27y^3 = (2x)^3 - (3y)^3$
 $= (2x - 3y)(4x^2 + 6xy + 9y^2).$

Note. The middle term $6xy$ is the product of $2x$ and $3y$.

Example 2. $64a^3 + 1 = (4a)^3 + (1)^3$
 $= (4a + 1)(16a^2 - 4a + 1).$

We may usually omit the intermediate step and write down the factors at once.

Examples. $343a^3 - 27x^3 = (7a^2 - 3x)(49a^4 + 21a^2x + 9x^2)$.
 $8x^3 + 729 = (2x^3 + 9)(4x^3 - 18x^3 + 81).$

EXAMPLES XVII. k.

Resolve into factors :

1. $x^3 - y^3$. 2. $x^3 + y^3$. 3. $x^3 - 1$. 4. $1 + a^3$
 5. $8x^3 - y^3$. 6. $x^3 + 8y^3$. 7. $27x^3 + 1$. 8. $1 - 8y^3$.
 9. $a^3b^3 - c^3$. 10. $8a^3 + 27y^3$. 11. $1 - 343x^3$. 12. $64 + y^3$.
 13. $125 + a^3$. 14. $216 - a^3$. 15. $a^3b^3 + 512$. 16. $1000y^3 - 1$.

Resolve into factors :

- | | | |
|--------------------------|--------------------------|-------------------------|
| 17. $x^3 + 64y^3$. | 18. $27 - 1000x^3$. | 19. $a^3b^3 + 216c^3$. |
| 20. $343 - 8x^3$. | 21. $a^3 + 27b^3$. | 22. $27x^3 - 64y^3$. |
| 23. $125x^3 - 1$. | 24. $216p^3 - 343$. | 25. $x^3y^3 + z^3$. |
| 26. $a^3b^3c^3 - 1$. | 27. $343x^3 + 1000y^3$. | 28. $729a^3 - 64b^3$. |
| 29. $8a^3b^3 + 125x^3$. | 30. $x^3y^3 - 216z^3$. | 31. $x^6 - 27y^3$. |
| 32. $64x^6 + 125y^3$. | 33. $8x^3 - z^6$. | 34. $216x^6 - b^3$. |
| 35. $a^3 + 343b^3$. | 36. $a^6 + 729b^3$. | 37. $8x^3 - 729y^6$. |
| 38. $p^3q^3 - 27x^3$. | 39. $x^3 - 64y^6$. | 40. $x^3y^3 - 512$. |

137. Before concluding this chapter we shall draw attention to a few miscellaneous cases of resolution into factors.

Example 1. Resolve into factors $16a^4 - 81b^4$.

$$\begin{aligned} 16a^4 - 81b^4 &= (4a^2 + 9b^2)(4a^2 - 9b^2) \\ &= (4a^2 + 9b^2)(4a^2 - 9b^2). \end{aligned}$$

Example 2. Resolve into factors $x^6 - y^6$.

$$\begin{aligned} x^6 - y^6 &= (x^3 + y^3)(x^3 - y^3) \\ &= (x + y)(x^2 - xy + y^2)(x - y)(x^2 + xy + y^2). \end{aligned}$$

Note. When an expression can be arranged either as the difference of two squares, or as the difference of two cubes, each of the methods explained in Arts. 133, 136 will be applicable. It will, however, be found simplest to first use the rule for resolving into factors the difference of two squares.

In all cases where an expression to be resolved contains a simple factor common to each of its terms, this should be first taken outside a bracket as explained in Art. 126.

Example 3. Resolve into factors $28x^4y + 64x^2y - 60x^2y$.

$$\begin{aligned} 28x^4y + 64x^2y - 60x^2y &= 4x^2y(7x^2 + 16x - 15) \\ &= 4x^2y(7x - 5)(x + 3). \end{aligned}$$

Example 4. Resolve into factors $x^3p^2 - 8y^3p^2 - 4x^3q^2 + 32y^3q^2$.

$$\begin{aligned} \text{The expression} &= p^2(x^3 - 8y^3) - 4q^2(x^3 - 8y^3) \\ &= (x^3 - 8y^3)(p^2 - 4q^2) \\ &= (x - 2y)(x^2 + 2xy + 4y^2)(p + 2q)(p - 2q). \end{aligned}$$

Example 5. Resolve into factors $4x^2 - 25y^2 + 2x + 5y$.

$$\begin{aligned} 4x^2 - 25y^2 + 2x + 5y &= (2x + 5y)(2x - 5y) + 2x + 5y \\ &= (2x + 5y)(2x - 5y + 1). \end{aligned}$$

Resolve

1. $a^2 - y^2$
4. $729y^6 -$
6. $2mn +$
8. $a^4 + b^4$
10. $(a + b +$
12. $x^2 - 10x$
14. $x^2 - a^2$
16. $21x^2 + 8$
18. $c^5d^3 - c^4$
20. $x^2 - 6x$
22. $acx^2 - b$
24. $x^4 + 4x^2$
27. $500x^2y$
30. $b^2 + c^2 -$
32. $14a^2x^3 -$
34. $1 - (m^2 -$
37. $8x^2y + 5$
39. $729a^7b -$
41. $a^{12} - b^{12}$
43. $(a + b)^4$
45. $(c + d)^3 -$
47. $250(a -$
49. $8(x + y)$
51. $a^2 - b^2 +$
53. $a^3 + b^3 +$
55. $4(x - y)$

[Miscellaneous
furnish further

EXAMPLES XVII. 1.

Resolve into two or more factors :

1. $a^2 - y^2 - 2yz - z^2$.
2. $x^6 - y^6z^6$.
3. $6x^2 - x - 77$.
4. $729y^6 - 64x^6$.
5. $x^6 - 4096$.
6. $2mn + 2xy + m^2 + n^2 - x^2 - y^2$.
7. $33x^2 - 16x - 65$.
8. $a^4 + b^4 - c^4 - d^4 + 2a^2b^2 - 2c^2d^2$.
9. $m^3x + m^2y - n^3x - n^3y$.
10. $(a + b + c)^2 - (a - b - c)^2$.
11. $4 + 4x + 2ay + x^2 - a^2 - y^2$.
12. $x^2 - 10x - 119$.
13. $a^2 - b^2 - c^2 + d^2 - 2(ad - bc)$.
14. $x^2 - a^2 + y^2 - 2xy$.
15. $a^2 + x^2 - (y^2 + z^2) - 2(yz - ax)$.
16. $21x^2 + 82x - 39$.
17. $1 - a^2x^2 - b^2y^2 + 2abxy$.
18. $c^5d^3 - c^3 - a^2c^3d^3 + a^2$.
19. $a^2x^6 - a^2y^6 - b^2x^6 + b^2y^6$.
20. $x^2 - 6x - 247$.
21. $a^3x^3 - c^3x^3 - a^3y^2 + c^3y^2$.
22. $acx^2 - bcx + adx - bd$.
23. $a^2x - b^2x + a^2y - b^2y$.
24. $x^4 + 4x^2y^2z^2 + 4y^4z^4$.
25. $a^3b^3 + 512$.
26. $2x^2 + 17x + 35$.
27. $500x^2y - 20y^3$.
28. $a^5 - 8a^2b^3$.
29. $a^2x^3 - 16x^2y^2$.
30. $b^2 + c^2 - a^2 - 2bc$.
31. $5x^4 - 15x^3 - 90x^2$.
32. $14a^2x^3 - 35a^3x^2 + 14a^4x$.
33. $x^3 - 1$.
34. $1 - (m^2 + n^2) + 2mn$.
35. $75x^4 - 48a^4$.
36. $5a^4b^4 - 5ab$.
37. $8x^2y + 52xy + 60y$.
38. $3x^2y^2 + 26axy + 35a^2$.
39. $729a^7b - ab^7$.
40. $a^2x^6 - 64a^2y^6$.
41. $a^{12} - b^{12}$.
42. $24x^2y^2 - 30xy^3 - 36y^4$.
43. $(a + b)^4 - 1$.
44. $a^4 - (b + c)^4$.
45. $(c + d)^3 - 1$.
46. $1 - (x - y)^3$.
47. $250(a - b)^3 + 2$.
48. $(c + d)^3 + (c - d)^3$.
49. $8(x + y)^3 - (2x - y)^3$.
50. $x^3 - 4y^2 + x - 2y$.
51. $a^2 - b^2 + a - b$.
52. $(a + b)^2 + a + b$.
53. $a^3 + b^3 + a + b$.
54. $a^2 - 9b^2 + a + 3b$.
55. $4(x - y)^3 - (x - y)$.
56. $x^4y - x^2y^3 - x^2y^2 + xy^4$.

[Miscellaneous Examples IV., p. 174, and Chapter XXVIII. will furnish further practice in Resolution into Factors.]

MISCELLANEOUS EXAMPLES III.

1. Subtract $3x^3 - 7x + 1$ from $2x^2 - 5x - 3$, then subtract the difference from zero, and add this last result to $2x^2 - 2x^3 - 4$.

2. Simplify

$$2\{3a - (4b - 5c)\} + 4\{4a - (5b - 2c)\} + 4\{5a - 3(b - c)\}.$$

3. Find the product of

$$a^3 - 2a^2c + 2ac^2 - c^3$$

and

$$a^3 + 2a^2c + 2ac^2 + c^3.$$

4. Solve the equations:

$$(1) \frac{x}{2} + \frac{x}{3} = \frac{x}{4} + 7;$$

$$(2) \begin{aligned} 9x + 5y &= 75, \\ 7x - 4y &= 11. \end{aligned}$$

5. Find the square root of $8x^4 + 16x^2 + 1 - 8x - 2x^3 + x^6$.

6. Find a number whose third, fourth, sixth, and eighth parts together make up 63.

7. If $a=4$, $b=3$, $c=2$, find the value of $\frac{a^3 - b^3}{b+c} + \frac{b^3 - c^3}{c+a} + \frac{c^3 - a^3}{a+b}$.

8. Divide $x^4 + \frac{9}{4}x^3 + \frac{21}{8}x^2 + \frac{33}{16}x + \frac{5}{16}$ by $x^2 + \frac{3x}{2} + \frac{1}{4}$

9. Add $5x^2 - 6x$ to the excess of 1 over $3x^2 - 5x + 1$.

10. Find the factors of (1) $a^2x^2 - 2ax - 15$; (2) $4m^4 - 81p^2q^2$.

11. Solve the equations:

$$(1) \begin{aligned} 13x + 11y &= 18, \\ 11x + 13y &= 30. \end{aligned}$$

$$(2) \begin{aligned} 57x + 52y &= 181, \\ 76x - 39y &= 458. \end{aligned}$$

12. A train which travels a miles in b hours is p times as fast as a coach. If the coach takes m hours to cover the distance between two places, how many miles are they apart?

13. Find the continued product of $3x^2 - 2x + 3$, $4x + 5$, $7x - 2$.

14. Solve the equations:

$$(1) \frac{5x}{7} - \frac{4}{5}\left(x - \frac{3}{4}\right) - \frac{2}{21}\left(x + \frac{7}{2}\right) + 1 = 0;$$

$$(2) 2\left(\frac{5x}{3} - 1\right) + \frac{11}{5}\left(1 + \frac{14x}{33}\right) = \frac{2x+7}{5} - 7.$$

15. Write down the square of $x^3 + 7x - 11$.

16. Res

(1

17. Fin

18. A h
sum he ther

19. Sim

20. She

21. Exp

(1

(2

22. Solv

and shew th

23. Divi

24. Wh
when 60 ap
cost as muc

25. Exp
in simple f
product of

26. If x

$x -$

27. Divi
by

28. Solv

4

29. Res

30. In h
whole of w

If $m=4$,
answer?

16. Resolve into factors :

(1) $x^3 + 2ax - bx - 2ab$; (2) $x^4 + 10x^2y - 56y^2$.

17. Find the H.C.F. and L.C.M. of $49bc^3$, $21a^2b^2$, $56ca^3$, $63abc^2$.

18. A has £50, and B has £6; after B has won from A a certain sum he then has five-ninths of what A has : how much did B win ?

19. Simplify $\frac{15a^2p^3}{56mk^2} \times \frac{49ak^5}{40p^2m^3} \div \frac{7a^3k^3}{64m^4}$.

20. Shew that $a(a-1)(a-2)(a-3) = (a^2 - 3a + 1)^2 - 1$.

21. Express by means of symbols :

(1) The excess of m over n is greater than a by c ;

(2) Three times the square of ab together with the cube of c is equal to p times the sum of m and n .

22. Solve $\frac{x}{4}\left(3 - \frac{8}{x}\right) - \frac{7}{8}\left(7 - \frac{3x}{4}\right) = 15\left(\frac{1}{3} - \frac{x}{64}\right)$,

and shew that $x=2$ does not satisfy the equation.

23. Divide the product of $3x^2 - 2xy - y^2$ and $2x - y$ by $x - y$.

24. What is the price of apples per dozen, and of eggs per score, when 60 apples and 100 eggs together cost 8s. 4d., and 72 apples cost as much as 30 eggs ?

25. Express the product $(2x^2 - 13x + 15)(x^2 - 4x - 5)(2x^2 - x - 3)$ in simple factors, and thence write down its square root as the product of three binomial factors.

26. If $x=6$, $y=7$, $z=8$, find the value of

$$x - (y - z) - 2[x + z - 3\{ - 2(y - 1) \}] + 4\left[\frac{x}{2} - \left(3 - \frac{9}{2}y\right)\right].$$

27. Divide $6x^5 + 57x^4y + 128x^3y^2 - 60x^2y^3 - 130xy^4 + 63y^5$ by $3x^3 + 15x^2y + 7xy^2 - 9y^3$.

28. Solve the equations :

$$4x + 2y + z = 14, \quad 3x - y + 2z = 3, \quad x + 7y - z = 23.$$

29. Resolve into two or more factors :

(1) $x^2y - 4xy^3$; (2) $2m^4 + m^2n^2 - 3n^4$.

30. In how many days will a men do $\frac{1}{m}$ th of a piece of work, the whole of which can be done by b men in c days ?

If $m=4$, $a=24$, $b=14$, $c=18$, what is the numerical value of the answer ?

CHAPTER XVIII.

HIGHEST COMMON FACTOR.

138. DEFINITION. The **highest common factor** of two or more algebraical expressions is the *expression of highest dimensions* which divides each of them without remainder.

Note. The term *greatest common measure* is sometimes used instead of *highest common factor*; but, strictly speaking, the term *greatest common measure* ought to be confined to arithmetical quantities; for the highest common factor is not necessarily the greatest common measure in all cases, as will appear later. [Art. 145.]

In Chap. XI. we have explained how to write down by inspection the highest common factor of two or more *simple* expressions. An analogous method will enable us readily to find the highest common factor of *compound* expressions which are given as the product of factors, or which can be easily resolved into factors.

Example 1. Find the highest common factor of $4cx^3$ and $2cx^3 + 4c^2x^2$.

It will be easy to pick out the common factors if the expressions are arranged as follows:

$$\begin{aligned} 4cx^3 &= 4cx^3, \\ 2cx^3 + 4c^2x^2 &= 2cx^2(x + 2c); \end{aligned}$$

therefore the H.C.F. is $2cx^2$.

Example 2. Find the highest common factor of $3a^2 + 9ab$, $a^3 - 9ab^2$, $a^3 + 6a^2b + 9ab^2$.

Resolving each expression into its factors, we have

$$\begin{aligned} 3a^2 + 9ab &= 3a(a + 3b), \\ a^3 - 9ab^2 &= a(a + 3b)(a - 3b), \\ a^3 + 6a^2b + 9ab^2 &= a(a + 3b)(a + 3b); \end{aligned}$$

therefore the H.C.F. is $a(a + 3b)$.

139. W
different po
be careful t
the highest
all the giv

Example

x

Example

Resolving

Therefore
factor is $a(x$

Find the

1. $a^2 + ab$,
3. $2x^2 - 2x$
5. $x^3 + x^2y$
7. $a^3 - a^2x$
9. $a^2bx + a$
11. $a^2 - x^2$,
13. $20x - 4$,
15. $x^2 + x$,
17. $x^2 - 2xy$
19. $x^3 + 8y^3$,
21. $x^2 + 3x -$
23. $x^2 - 18x$
25. $12x^2 + x$
27. $c^2x^2 - d^2$
28. $x^5 - xy^2$,
29. $a^3x - a^2b$
30. $2x^2 + 9x$
31. $3x^4 + 8x^2$

E.A.

139. When there are two or more expressions containing different powers of the same *compound* factor, the student should be careful to notice that the highest common factor must contain the highest power of the compound factor which is common to all the given expressions.

Example 1. The highest common factor of

$$x(a-x)^2, a(a-x)^3, \text{ and } 2ax(a-x)^5 \text{ is } (a-x)^2.$$

Example 2. Find the highest common factor of

$$ax^2+2a^2x+a^3, 2ax^2-4a^2x-6a^3, 3(ax+a^2)^2.$$

Resolving the expressions into factors, we have

$$ax^2+2a^2x+a^3=a(x^2+2ax+a^2) \\ =a(x+a)^2 \dots\dots\dots(1),$$

$$2ax^2-4a^2x-6a^3=2a(x^2-2ax-3a^2) \\ =2a(x+a)(x-3a) \dots\dots\dots(2),$$

$$3(ax+a^2)^2=3a^2(x+a)^2 \dots\dots\dots(3).$$

Therefore from (1), (2), (3), by inspection, the highest common factor is $a(x+a)$.

EXAMPLES XVIII. a.

Find the highest common factor of

- | | |
|--|--------------------------------|
| 1. $a^2+ab, a^2-b^2.$ | 2. $(x+y)^2, x^2-y^2.$ |
| 3. $2x^2-2xy, x^3-x^2y.$ | 4. $6x^2-9xy, 4x^2-9y^2.$ |
| 5. $x^3+x^2y, x^3+y^3.$ | 6. $a^3b-ab^3, a^5b^3-a^2b^5.$ |
| 7. $a^3-a^2x, a^3-ax^2, a^4-ax^3.$ | 8. $a^2-4x^2, a^3+2ax.$ |
| 9. $a^2bx+ab^2x, a^2b-b^3.$ | 10. $2x^2y-6xy^2, x^2-9y^2.$ |
| 11. $a^2-x^2, a^2-ax, a^2x-ax^2.$ | 12. $4x^2+2xy, 12x^2y-3y^3.$ |
| 13. $20x-4, 50x^2-2.$ | 14. $6bx+4by, 9cx+6cy.$ |
| 15. $x^2+x, (x+1)^2, x^3+1.$ | 16. $xy-y, x^4y-xy.$ |
| 17. $x^2-2xy+y^2, (x-y)^2.$ | 18. $x^3+a^2x, x^4-a^4.$ |
| 19. $x^3+8y^3, x^3+xy-2y^2.$ | 20. $x^4-27a^3x, (x-3a)^2.$ |
| 21. $x^2+3x+2, x^2-4.$ | 22. $x^3-x-20, x^2-9x+20.$ |
| 23. $x^2-18x+45, x^2-9.$ | 24. $2x^2-7x+3, 3x^2-7x-6.$ |
| 25. $12x^2+x-1, 15x^2+8x+1.$ | 26. $2x^2-x-1, 3x^2-x-2.$ |
| 27. $c^2x^2-d^2, acx^2-bcx+adx-bd.$ | |
| 28. $x^5-xy^2, x^3+x^2y+xy+y^3.$ | |
| 29. $a^3x-a^2bx-6ab^2x, a^2bx^2-4ab^3x^2+3b^5x^2.$ | |
| 30. $2x^2+9x+4, 2x^2+11x+5, 2x^2-3x-2.$ | |
| 31. $3x^4+8x^3+4x^2, 3x^5+11x^4+6x^3, 3x^4-16x^3-12x^2.$ | |

140. The highest common factor should always be determined by inspection when possible, but it will sometimes happen that expressions cannot be readily resolved into factors. To find the highest common factor in such cases, we adopt a method analogous to that used in Arithmetic, for finding the greatest common measure of two or more numbers.

141. We shall now work out examples illustrative of the algebraical process of finding the highest common factor, postponing for the present the complete proof of the rules we use. But we may conveniently *enunciate* two principles, which the student should bear in mind in reading the examples which follow.

I. If an expression contain a certain factor, any multiple of the expression is divisible by that factor.

II. If two expressions have a common factor, it will divide their sum and their difference; and also the sum and the difference of any multiples of them.

Example. Find the highest common factor of

$$\begin{array}{r|l}
 4x^3 - 3x^2 - 24x - 9 & 8x^3 - 2x^2 - 53x - 39 \\
 4x^3 - 5x^2 - 21x & 8x^3 - 6x^2 - 48x - 18 \\
 \hline
 2x & \begin{array}{r} 2x^2 - 3x - 9 \\ 2x^2 - 0x \\ \hline 3x - 9 \end{array} \\
 3 & \begin{array}{r} 3x - 9 \\ 3x - 9 \\ \hline 0 \end{array}
 \end{array}
 \quad
 \begin{array}{r|l}
 8x^3 - 2x^2 - 53x - 39 & 2 \\
 8x^3 - 6x^2 - 48x - 18 & \\
 \hline
 4x^2 - 5x - 21 & 2 \\
 4x^2 - 6x - 18 & \\
 \hline
 x - 3 &
 \end{array}$$

Therefore the H.C.F. is $x - 3$.

Explanation. First arrange the given expressions according to descending or ascending powers of x . The expressions so arranged having their first terms of the same order, we take for divisor that whose highest power has the smaller coefficient. Arrange the work in parallel columns as above. When the first remainder $4x^2 - 5x - 21$ is made the divisor we put the quotient x to the *left* of the dividend. Again, when the second remainder $2x^2 - 3x - 9$ is in turn made the divisor, the quotient 2 is placed to the *right*; and so on. As in Arithmetic, the last divisor $x - 3$ is the highest common factor required.

142. This method is only useful to determine the *compound* factor of the highest common factor. Simple factors of the given expressions must be first removed from them, and the highest common factor of these, if any, must be observed and multiplied into the *compound* factor given by the rule.

Example.

24

We have

and

Also $2x$ and
factors $2x$ and
as in Art. 14

$2x$

2

Therefore

143. So far
applicable to
But in many
method will
understood
the remainder
common factor

Example 1.

Here on r
not contained
But noticing
 $9(3x^2 - 10x + 8)$
the course of
H.C.F. we a
two original
H.C.F. can h
go on with di

Example. Find the highest common factor of

$$24x^4 - 2x^3 - 60x^2 - 32x \text{ and } 18x^4 - 6x^3 - 39x^2 - 18x.$$

$$\text{We have } 24x^4 - 2x^3 - 60x^2 - 32x = 2x(12x^3 - x^2 - 30x - 16),$$

$$\text{and } 18x^4 - 6x^3 - 39x^2 - 18x = 3x(6x^3 - 2x^2 - 13x - 6).$$

Also $2x$ and $3x$ have the common factor x . Removing the simple factors $2x$ and $3x$, and *reserving* their common factor x , we continue as in Art. 141.

$2x$	$\begin{array}{r} 6x^3 - 2x^2 - 13x - 6 \\ 6x^3 - 8x^2 - 8x \\ \hline 6x^2 - 5x - 6 \\ 6x^2 - 8x - 8 \\ \hline 3x + 2 \end{array}$	$\begin{array}{r} 12x^3 - x^2 - 30x - 16 \\ 12x^3 - 4x^2 - 26x - 12 \\ \hline 3x^2 - 4x - 4 \\ 3x^2 + 2x \\ \hline - 6x - 4 \\ - 6x - 4 \\ \hline 0 \end{array}$	$\left \begin{array}{l} 2 \\ x \\ -2 \end{array} \right.$
------	--	--	--

Therefore the H.C.F. is $x(3x+2)$.

143. So far the process of Arithmetic has been found exactly applicable to the algebraical expressions we have considered. But in many cases certain modifications of the arithmetical method will be found necessary. These will be more clearly understood if it is remembered that, at every stage of the work, the remainder must contain as a factor of itself the highest common factor we are seeking. [See Art. 141, I. & II.]

Example 1. Find the highest common factor of

$$3x^3 - 13x^2 + 23x - 21 \text{ and } 6x^3 + x^2 - 44x + 21.$$

$$\begin{array}{r|l} 3x^3 - 13x^2 + 23x - 21 & 6x^3 + x^2 - 44x + 21 \\ \hline & 6x^3 - 26x^2 + 46x - 42 \\ \hline & 27x^2 - 90x + 63 \end{array} \quad \left| \begin{array}{l} 2 \\ 27x^2 - 90x + 63 \end{array} \right.$$

Here on making $27x^2 - 90x + 63$ a divisor, we find that it is not contained in $3x^3 - 13x^2 + 23x - 21$ with an *integral* quotient. But noticing that $27x^2 - 90x + 63$ may be written in the form $9(3x^2 - 10x + 7)$, and also bearing in mind that every remainder in the course of the work contains the H.C.F., we conclude that the H.C.F. we are seeking is contained in $9(3x^2 - 10x + 7)$. But the two original expressions have no *simple* factors, therefore their H.C.F. can have none. We may therefore *reject* the factor 9 and go on with divisor $3x^2 - 10x + 7$.

Resuming the work, we have

$$\begin{array}{r|l}
 x & \begin{array}{r} 3x^3 - 13x^2 + 23x - 21 \\ 3x^3 - 10x^2 + 7x \\ \hline - 3x^2 + 16x - 21 \\ - 3x^2 + 10x - 7 \\ \hline 2) 6x - 14 \\ 3x - 7 \end{array} & \begin{array}{r} 3x^2 - 10x + 7 \\ 3x^2 - 7x \\ \hline - 3x + 7 \\ - 3x + 7 \\ \hline 0 \end{array} x \\
 -1 & & -1
 \end{array}$$

Therefore the H.C.F. is $3x - 7$.

The factor 2 has been removed on the same grounds as the factor 9 above.

Example 2. Find the highest common factor of

$$2x^3 + x^2 - x - 2 \dots\dots\dots(1),$$

and

$$3x^3 - 2x^2 + x - 2 \dots\dots\dots(2),$$

As the expressions stand we cannot begin to divide one by the other without using a fractional quotient. The difficulty may be obviated by *introducing* a suitable factor, just as in the last case we found it useful to remove a factor when we could no longer proceed with the division in the ordinary way. The given expressions have no common *simple* factor, hence their H.C.F. cannot be affected if we multiply either of them by any simple factor.

Multiply (2) by 2, and use (1) as a divisor :

$$\begin{array}{r|l}
 \begin{array}{r} 2x^3 + x^2 - x - 2 \\ 7 \\ \hline -2x \quad 14x^3 + 7x^2 - 7x - 14 \\ \hline 14x^3 - 10x^2 - 4x \\ \hline 17x \quad 17x^3 - 3x - 14 \\ \hline 17x^3 - 17x \\ \hline 14 \quad 14x - 14 \\ \hline 14x - 14 \end{array} & \begin{array}{r} 6x^3 - 4x^2 + 2x - 4 \quad 3 \\ 6x^3 + 3x^2 - 3x - 6 \\ \hline - 7x^2 + 5x + 2 \\ 17 \\ \hline -119x^2 + 85x + 34 \\ -119x^2 + 21x + 98 \\ \hline 64) 64x - 64 \\ x - 1 \end{array}
 \end{array}$$

Therefore the H.C.F. is $x - 1$.

After the first division the factor 7 is introduced because the first remainder $-7x^2 + 5x + 2$ will not divide $2x^3 + x^2 - x - 2$. At the next stage the factor 17 is introduced for a similar reason, and finally the factor 64 is removed as explained in Example 1.

Note. Here the highest common factor might have been more easily obtained by arranging the expressions in *ascending* powers of x . In this case it will be found that there is no need to introduce a numerical factor in the course of the work. Detached coefficients, as explained in Art. 45, may also be used with advantage here, and will often effect a considerable saving of labour.

144. F
multiply
remainder
which doe

145. Le
written in

Then thei
 $2x^2 + 3x + 2$
if, howeve

and
and the gr
5 is the nu
factor. Th
common fa
do not in t

The rea
expressions
116 respec
whereas th

It will t
two expres
when the
this reason
when appli

Find the

1. $x^3 + 2x^2$
2. $x^3 - 5x^2$
3. $x^3 + 2x^2$
4. $x^3 + 4x^2$
5. $x^3 - x^2$
6. $x^3 + 3x^2$
7. $x^3 - 5x^2$
8. $x^4 - 2x^3$
9. $2x^3 - 5x^2$

144. From the last two examples it appears that we may multiply or divide either of the given expressions, or any of the remainders which occur in the course of the work, by any factor which does not divide both of the given expressions.

145. Let the two expressions in Example 2, Art. 143, be written in the form

$$2x^3 + x^2 - x - 2 = (x-1)(2x^2 + 3x + 2),$$

$$3x^3 - 2x^2 + x - 2 = (x-1)(3x^2 + x + 2).$$

Then their highest common factor is $x-1$, and therefore $2x^2 + 3x + 2$ and $3x^2 + x + 2$ have no algebraical common divisor. If, however, we put $x=6$, then

$$2x^3 + x^2 - x - 2 = 460,$$

and

$$3x^3 - 2x^2 + x - 2 = 580;$$

and the greatest common measure of 460 and 580 is 20; whereas 5 is the numerical value of $x-1$, the algebraical highest common factor. Thus the numerical values of the algebraical highest common factor and of the arithmetical greatest common measure do not in this case agree.

The reason may be explained as follows: when $x=6$, the expressions $2x^2 + 3x + 2$ and $3x^2 + x + 2$ become equal to 92 and 116 respectively, and have a common arithmetical factor 4; whereas the expressions have no algebraical common factor.

It will thus often happen that the highest common factor of two expressions, and their numerical greatest common measure, when the letters have particular values, are not the same; for this reason the term *greatest common measure* is inappropriate when applied to algebraical quantities.

EXAMPLES XVIII. b.

Find the highest common factor of the following expressions:

1. $x^3 + 2x^2 - 13x + 10$, $x^3 + x^2 - 10x + 8$.
2. $x^3 - 5x^2 - 99x + 40$, $x^3 - 6x^2 - 86x + 35$.
3. $x^3 + 2x^2 - 8x - 16$, $x^3 + 3x^2 - 8x - 24$.
4. $x^3 + 4x^2 - 5x - 20$, $x^3 + 6x^2 - 5x - 30$.
5. $x^3 - x^2 - 5x - 3$, $x^3 - 4x^2 - 11x - 6$.
6. $x^3 + 3x^2 - 8x - 24$, $x^3 + 3x^2 - 3x - 9$.
7. $a^3 - 5a^2x + 7ax^2 - 3x^3$, $a^3 - 3ax^2 + 2x^3$.
8. $x^4 - 2x^3 - 4x - 7$, $x^4 + x^3 - 3x^2 - x + 2$.
9. $2x^3 - 5x^2 + 11x + 7$, $4x^3 - 11x^2 + 25x + 7$.

Find the highest common factor of the following expressions:

10. $2x^3 + 4x^2 - 7x - 14$, $6x^3 - 10x^2 - 21x + 35$.
11. $3x^4 - 3x^3 - 2x^2 - x - 1$, $9x^4 - 3x^3 - x - 1$.
12. $2x^4 - 2x^3 + x^2 + 3x - 6$, $4x^4 - 2x^3 + 3x - 9$.
13. $3x^3 - 3ax^2 + 2a^2x - 2a^3$, $3x^3 + 12ax^2 + 2a^2x + 8a^3$.
14. $2x^3 - 9ax^2 + 9a^2x - 7a^3$, $4x^3 - 20ax^2 + 20a^2x - 16a^3$.
15. $10x^3 + 25ax^2 - 5a^3$, $4x^3 + 9ax^2 - 2a^2x - a^3$.
16. $6a^3 + 13a^2x - 9ax^2 - 10x^3$, $9a^3 + 12a^2x - 11ax^2 - 10x^3$.
17. $24x^4y + 72x^3y^2 - 6a^2y^3 - 90xy^4$, $6x^4y^2 + 13x^3y^3 - 4x^2y^4 - 15xy^5$.
18. $4x^5a^2 + 10x^4a^3 - 60x^3a^4 + 54x^2a^5$, $24x^5a^3 + 30x^3a^5 - 126x^2a^6$.
19. $4x^5 + 14x^4 + 20x^3 + 70x^2$, $8x^7 + 28x^6 - 8x^5 - 12x^4 + 56x^3$.
20. $72x^3 - 12ax^2 + 72a^2x - 420a^3$, $18x^3 + 42ax^2 - 282a^2x + 270a^3$.
21. $9x^4 + 2x^2y^2 + y^4$, $3x^4 - 8x^2y + 5x^2y^2 - 2xy^3$.
22. $x^5 - x^3 - x + 1$, $x^7 + x^6 + x^4 - 1$.
23. $1 + x + x^3 - x^5$, $1 - x^4 - x^6 + x^7$.
24. $6 - 8a - 32a^2 - 18a^3$, $20 - 35a - 95a^2 - 40a^3$.
25. $9x^2 - 15x^3 - 45x^4 - 12x^5$, $42x - 49x^2 - 203x^3 - 84x^4$.
26. $3x^5 - 5x^3 + 2$, $2x^5 - 5x^3 + 3$.
27. $4x^5 - 6x^3 - 28x$, $6x^4 + 10x^3 - 17x^2 - 35x - 14$.

*146. The statements of Art. 141 may be proved as follows.

I. If F divides A it will also divide mA .

For suppose $A = aF$, then $mA = maF$.

Thus F is a factor of mA .

II. If F divides A and B , then it will divide $mA \pm nB$.

For suppose $A = aF$, $B = bF$,

$$\begin{aligned} \text{then} \quad mA \pm nB &= maF \pm nbF \\ &= F(ma \pm nb). \end{aligned}$$

Thus F divides $mA \pm nB$.

*147. We may now enunciate and prove the rule for finding the highest common factor of any two compound algebraical expressions.

We suppose that any simple factors are first removed. [See Example, Art. 142.]

Let A and B be the two expressions after the simple factors have been removed. Let them be arranged in descending or ascending powers of some common letter; also let the highest power of that letter in B be not less than the highest power in A .

Divide
Suppose C
so obtain
make A d
factor n .
Finally, d
there is no

The wo

First, to
By exam
 D , therefo
 A , since n

Again,
divides A
divides B

Secondly,

If not, l

Then X
fore D (sin

Thus X
 X is of hi

Therefor

*148. A, B, C m

First de
next find C
be the req

For F c
and G is t
is the high

Divide B by A ; let p be the quotient, and C the remainder. Suppose C to have a *simple* factor m . Remove this factor, and so obtain a new divisor D . Further, suppose that in order to make A divisible by D it is necessary to multiply A by a *simple* factor n . Let q be the next quotient and E the remainder. Finally, divide D by E ; let r be the quotient, and suppose that there is no remainder. Then E will be the H.C.F. required.

The work will stand thus :

$$\begin{array}{r}
 A)B(p \\
 \quad pA \\
 \hline
 \quad m)C \\
 \quad \quad D)nA(q \\
 \quad \quad \quad qD \\
 \quad \quad \quad \hline
 \quad \quad \quad E)D(r \\
 \quad \quad \quad \quad rE \\
 \quad \quad \quad \quad \hline
 \quad \quad \quad \quad \quad 0
 \end{array}$$

First, to shew that E is a common factor of A and B .

By examining the steps of the work, it is clear that E divides D , therefore also qD ; therefore $qD + E$, therefore nA ; therefore A , since n is a *simple* factor.

Again, E divides D , therefore mD , that is, C . And since E divides A and C , it also divides $pA + C$, that is, B . Hence E divides both A and B .

Secondly, to show that E is the *highest* common factor.

If not, let there be a factor X of higher dimensions than E .

Then X divides A and B , therefore $B - pA$ that is, C ; therefore D (since m is a *simple* factor); therefore $nA - qD$, that is, E .

Thus X divides E ; which is impossible since by hypothesis, X is of higher dimensions than E .

Therefore E is the *highest* common factor.

*148. The highest common factor of three expressions A , B , C may be obtained as follows.

First determine F the highest common factor of A and B ; next find G the highest common factor of F and C ; then G will be the required highest common factor of A , B , C .

For F contains *every* factor which is common to A and B , and G is the highest common factor of F and C . Therefore G is the highest common factor of A , B , C .

CHAPTER XIX.

FRACTIONS.

[On first reading the subject, the student may omit the general proofs of the rules given in this chapter.]

149. In Chapter XII. we discussed the simpler kinds of fractions, using the ordinary arithmetical rules. We here propose to give proofs of those rules, and shew that they are applicable to algebraical fractions.

DEFINITION. If a quantity x be divided into b equal parts, and a of these parts be taken, the result is called *the fraction* $\frac{a}{b}$ of x . If x be the unit, the fraction $\frac{a}{b}$ of x is called simply "the fraction $\frac{a}{b}$ "; so that *the fraction* $\frac{a}{b}$ *represents a equal parts, b of which make up the unit.*

Note. This definition requires that a and b should be positive whole numbers. In Art. 155 we shall adopt a definition which will enable us to remove this restriction.

150. To prove that $\frac{a}{b} = \frac{ma}{mb}$, where a, b, m are positive integers.

By $\frac{a}{b}$ we mean a equal parts, b of which make up the unit ... (1);

by $\frac{ma}{mb}$ ma mb (2).

But

b parts in (1) = mb parts in (2);

$\therefore$ 1 part = m

$\therefore$ a parts = ma

that is,

$$\frac{a}{b} = \frac{ma}{mb}.$$

Conversely,

$$\frac{ma}{mb} = \frac{a}{b}.$$

CHAP. XIX.]

Hence, *the*
divide the n

151. An
equivalent
by any co
common fac
its lowest

Example 1

Example 2

Note. Th
until he has
convenient f

Reduce to

$$1. \frac{3a^2 - 6}{2a^2b - 4}$$

$$4. \frac{15a^2b}{100(a^2 - 4)}$$

$$7. \frac{x(2a^2 - 5a)}{a(4a^2x - 4)}$$

$$10. \frac{x^2 - 5x}{x^2 - 4x - 4}$$

$$13. \frac{x^2y + 2x^2}{x^3}$$

$$15. \frac{x^4 - 14x^2}{x^4 - 2x^2}$$

$$18. \frac{a^2x^2 - 9a}{ax^2 + 9a}$$

Hence, the value of a fraction is not altered if we multiply or divide the numerator and denominator by the same quantity.

Reduction to Lowest Terms.

151. An algebraical fraction may be changed into an equivalent fraction by dividing numerator and denominator by any common factor; if this factor be the highest common factor the resulting fraction is said to be **reduced to its lowest terms**.

Example 1. Reduce to lowest terms $\frac{24a^3c^2x^2}{18a^2x^2 - 12a^2x^3}$.

$$\text{The expression} = \frac{24a^3c^2x^2}{6a^2x^2(3a - 2x)} = \frac{4ac^2}{3a - 2x}.$$

Example 2. Reduce to lowest terms $\frac{6x^2 - 8xy}{9xy - 12y^2}$.

$$\text{The expression} = \frac{2x(3x - 4y)}{3y(3x - 4y)} = \frac{2x}{3y}.$$

Note. The beginner should be careful not to begin cancelling until he has expressed both numerator and denominator in the most convenient form, by resolution into factors where necessary.

EXAMPLES XIX. a.

Reduce to lowest terms :

1. $\frac{3a^2 - 6ab}{2a^2b - 4ab^2}$
2. $\frac{abx + bx^2}{acx + cx^2}$
3. $\frac{ax}{a^2x^2 - ax}$
4. $\frac{15a^2b^2c}{100(a^3 - a^2b)}$
5. $\frac{4x^2 - 9y^2}{4x^2 + 6xy}$
6. $\frac{20(x^3 - y^3)}{5x^2 + 5xy + 5y^2}$
7. $\frac{x(2a^2 - 3ax)}{a(4a^2x - 9x^2)}$
8. $\frac{x^3 - 2xy^2}{x^4 - 4x^2y^2 + 4y^4}$
9. $\frac{(xy - 3y^2)^2}{x^3y^2 - 27y^5}$
10. $\frac{x^2 - 5x}{x^2 - 4x - 5}$
11. $\frac{3x^2 + 6x}{x^2 + 4x + 4}$
12. $\frac{5a^2b + 10a^2b^2}{3a^2b^2 + 6ab^3}$
13. $\frac{x^2y + 2x^2y + 4xy}{x^3 - 8}$
14. $\frac{3a^4 + 9a^3b + 6a^2b^2}{a^4 + a^3b - 2a^2b^2}$
15. $\frac{x^4 - 14x^2 - 51}{x^4 - 2x^2 - 15}$
16. $\frac{x^2 + xy - 2y^2}{x^2 - y^2}$
17. $\frac{2x^2 + 17x + 21}{3x^2 + 26x + 35}$
18. $\frac{a^2x^2 - 16a^2}{ax^2 + 9ax + 20a}$
19. $\frac{3x^2 + 23x + 14}{3x^2 + 41x + 26}$
20. $\frac{27a + a^4}{18a - 6a^2 + 2a^3}$

152. When the factors of the numerator and denominator cannot be determined by inspection, the fraction may be reduced to its lowest terms by dividing both numerator and denominator by the highest common factor, which may be found by the rules given in Chap. xviii.

Example. Reduce to lowest terms $\frac{3x^3 - 13x^2 + 23x - 21}{15x^3 - 38x^2 - 2x + 21}$.

First Method. The H.C.F. of numerator and denominator is $3x - 7$.

Dividing numerator and denominator by $3x - 7$, we obtain as respective quotients $x^2 - 2x + 3$ and $5x^2 - x - 3$.

$$\text{Thus } \frac{3x^3 - 13x^2 + 23x - 21}{15x^3 - 38x^2 - 2x + 21} = \frac{(3x - 7)(x^2 - 2x + 3)}{(3x - 7)(5x^2 - x - 3)} = \frac{x^2 - 2x + 3}{5x^2 - x - 3}.$$

This is the simplest solution for the beginner; but in this and similar cases we may often effect the reduction without actually going through the process of finding the highest common factor.

Second Method. By Art. 141, the H.C.F. of numerator and denominator must be a factor of their sum $18x^3 - 51x^2 + 21x$, that is, of $3x(3x - 7)(2x - 1)$. If there be a common divisor it must clearly be $3x - 7$; hence arranging numerator and denominator so as to shew $3x - 7$ as a factor,

$$\begin{aligned} \text{the fraction} &= \frac{x^2(3x - 7) - 2x(3x - 7) + 3(3x - 7)}{5x^2(3x - 7) - x(3x - 7) - 3(3x - 7)} \\ &= \frac{(3x - 7)(x^2 - 2x + 3)}{(3x - 7)(5x^2 - x - 3)} \\ &= \frac{x^2 - 2x + 3}{5x^2 - x - 3}. \end{aligned}$$

153. If either numerator or denominator can readily be resolved into factors we may use the following method.

Example. Reduce to lowest terms $\frac{x^3 + 3x^2 - 4x}{7x^3 - 18x^2 + 6x + 5}$.

The numerator $= x(x^2 + 3x - 4) = x(x + 4)(x - 1)$.

Of these factors the only one which can be a common divisor is $x - 1$. Hence, arranging the denominator,

$$\begin{aligned} \text{the fraction} &= \frac{x(x + 4)(x - 1)}{7x^2(x - 1) - 11x(x - 1) - 5(x - 1)} \\ &= \frac{x(x + 4)(x - 1)}{(x - 1)(7x^2 - 11x - 5)} = \frac{x(x + 4)}{7x^2 - 11x - 5}. \end{aligned}$$

Reduce

$$1. \frac{a^3 - a}{a^3 + 3a}$$

$$3. \frac{a^3 + 2a}{a^3 + a}$$

$$5. \frac{4a^3 + 1}{6a^3 + 1}$$

$$7. \frac{x^2 - 2}{3x^2 + 7}$$

$$9. \frac{4x^3 + 1}{x^4 + a^2}$$

$$11. \frac{16x^4 - 1}{4x^2 + 1}$$

$$13. \frac{5x^3 + 1}{7x^3 - 4}$$

$$15. \frac{3x^3 - 2}{2x^3 + 1}$$

154. R
multiply th
divisible by

The rule

$$(1) \frac{a}{b} \text{ re}$$

$$\frac{ac}{b} \text{ r}$$

and the nu
the numbe

that is

$$(2)$$

EXAMPLES XIX. b.

Reduce to lowest terms :

1. $\frac{a^3 - a^2b - ab^2 - 2b^3}{a^3 + 3a^2b + 3ab^2 + 2b^3}$
2. $\frac{x^3 - 5x^2 + 7x - 3}{x^3 - 3x + 2}$
3. $\frac{a^3 + 2a^2 - 13a + 10}{a^3 + a^2 - 10a + 8}$
4. $\frac{2x^3 + 5x^2y - 30xy^2 + 27y^3}{4x^3 + 5xy^2 - 21y^3}$
5. $\frac{4a^3 + 12a^2b - ab^2 - 15b^3}{6a^3 + 13a^2b - 4ab^2 - 15b^3}$
6. $\frac{1 + 2x^2 + x^3 + 2x^4}{1 + 3x^2 + 2x^3 + 3x^4}$
7. $\frac{x^2 - 2x + 1}{3x^3 + 7x - 10}$
8. $\frac{3a^3 - 3a^2b + ab^2 - b^3}{4a^2 - 5ab + b^2}$
9. $\frac{4x^3 + 3ax^2 + a^3}{x^4 + ax^3 + a^2x + a^4}$
10. $\frac{4x^3 - 10x^2 + 4x + 2}{3x^4 - 2x^3 - 3x + 2}$
11. $\frac{16x^4 - 72x^2a^2 + 81a^4}{4x^2 + 12ax + 9a^2}$
12. $\frac{6x^3 + x^2 - 5x - 2}{6x^3 + 5x^2 - 3x - 2}$
13. $\frac{5x^3 + 2x^2 - 15x - 6}{7x^3 - 4x^2 - 21x + 12}$
14. $\frac{4x^4 + 11x^2 + 25}{4x^4 - 9x^2 + 30x - 25}$
15. $\frac{3x^3 - 27ax^2 + 78a^2x - 72a^3}{2x^3 + 10ax^2 - 4a^2x - 48a^3}$
16. $\frac{ax^3 - 5a^2x^2 - 99a^3x + 40a^4}{x^4 - 6ax^3 - 86a^2x^2 + 35a^3x}$

Multiplication and Division of Fractions.

154. **Rule I.** To multiply a fraction by an integer: multiply the numerator by that integer; or, if the denominator be divisible by the integer, divide the denominator by it.

The rule may be proved as follows :

(1) $\frac{a}{b}$ represents a equal parts, b of which make up the unit ;

$\frac{ac}{b}$ represents ac equal parts, b of which make up the unit ;

and the number of parts taken in the second fraction is c times the number taken in the first ;

that is $\frac{a}{b} \times c = \frac{ac}{b}$.

(2) $\frac{a}{ba} \times d = \frac{ad}{ba}$ by the preceding case,
 $= \frac{a}{b}$.

[Art. 151.]

155. By the preceding article

$$\frac{a}{b} \times b = \frac{ab}{b} = a,$$

that is, the fraction $\frac{a}{b}$ is that which must be multiplied by b in order to obtain a . But, by Art. 46, the quantity which must be multiplied by b in order to obtain a is the quotient resulting from the division of a by b ; we may therefore define a fraction thus:

the fraction $\frac{a}{b}$ is the quotient of a divided by b .

156. Rule II. To divide a fraction by an integer: *divide the numerator, if it be divisible, by the integer; or if the numerator be not divisible, multiply the denominator by that integer.*

The rule may be proved as follows:

(1) $\frac{ac}{b}$ represents ac equal parts, b of which make up the unit;

$\frac{a}{b}$ represents a equal parts, b of which make up the unit.

The number of parts taken in the first fraction is c times the number taken in the second. Therefore the second fraction is the quotient of the first fraction divided by c ;

that is

$$\frac{ac}{b} \div c = \frac{a}{b}.$$

(2) But if the numerator be not divisible by c , we have

$$\frac{a}{b} = \frac{ac}{bc};$$

$$\therefore \frac{a}{b} \div c = \frac{ac}{bc} \div c$$

$$= \frac{a}{bc}, \text{ by the preceding case.}$$

157. Rule III. To multiply together two or more fractions: *multiply together all the numerators to form a new numerator, and all the denominators to form a new denominator.*

To find the value of

$$\frac{a}{b} \times \frac{c}{d}.$$

Let

$$x = \frac{a}{b} \times \frac{c}{d}.$$

xix.]

Multipl

Dividing

Similarly

and so for

158. R
the divisor,

Since di
the quotie

Multipl

Hence

which prov

Example

by cancelli
and denomi

Multiplying each side by $b \times d$, we have

$$x \times b \times d = \frac{a}{b} \times \frac{c}{d} \times b \times d$$

$$= \frac{a}{b} \times b \times \frac{c}{d} \times d$$

[Art. 29.]

$$= a \times c$$

[Art. 154.]

$$\therefore x \times bd = ac.$$

Dividing each side by bd , we have

$$x = \frac{ac}{bd};$$

$$\therefore \frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}.$$

Similarly

$$\frac{a}{b} \times \frac{c}{d} \times \frac{e}{f} = \frac{ace}{bdf};$$

and so for any number of fractions.

158. Rule IV. To divide one fraction by another: invert the divisor, and proceed as in multiplication.

Since division is the inverse of multiplication, we may define the quotient x , when $\frac{a}{b}$ is divided by $\frac{c}{d}$ to be such that

$$x \times \frac{c}{d} = \frac{a}{b}.$$

Multiplying by $\frac{d}{c}$ we have $x \times \frac{c}{d} \times \frac{d}{c} = \frac{a}{b} \times \frac{d}{c};$

$$\therefore x = \frac{ad}{bc}.$$

Hence

$$\frac{a}{b} \div \frac{c}{d} = \frac{ad}{bc} = \frac{a}{b} \times \frac{d}{c},$$

[Art. 157.]

which proves the rule.

Example 1. Simplify $\frac{2a^2+3a}{4a^3} \times \frac{4a^2-6a}{12a+18}.$

$$\frac{2a^2+3a}{4a^3} \times \frac{4a^2-6a}{12a+18} = \frac{a(2a+3)}{4a^3} \times \frac{2a(2a-3)}{6(2a+3)}$$

$$= \frac{2a-3}{12a},$$

by cancelling those factors which are common to both numerator and denominator.

Example 2. Simplify $\frac{6x^2 - ax - 2a^2}{ax - a^2} \times \frac{x - a}{9x^2 - 4a^2} \div \frac{2x + a}{3ax + 2a^2}$

$$\begin{aligned} \text{The expression} &= \frac{6x^2 - ax - 2a^2}{ax - a^2} \times \frac{x - a}{9x^2 - 4a^2} \times \frac{3ax + 2a^2}{2x + a} \\ &= \frac{(3x - 2a)(2x + a)}{a(x - a)} \times \frac{x - a}{(3x + 2a)(3x - 2a)} \times \frac{a(3x + 2a)}{2x + a} \\ &= 1, \end{aligned}$$

since all the factors cancel each other.

EXAMPLES XIX. c.

Simplify

1. $\frac{14x^2 - 7x}{12x^3 + 24x^2} \div \frac{2x - 1}{x^2 + 2x}$
2. $\frac{a^2b^2 + 3ab}{4a^2 - 1} \div \frac{ab + 3}{2a + 1}$
3. $\frac{x^2 - 4a^2}{ax + 2a^2} \times \frac{2a}{x - 2a}$
4. $\frac{a^2 - 121}{a^2 - 4} \div \frac{a + 11}{a + 2}$
5. $\frac{16x^2 - 9a^2}{x^2 - 4} \times \frac{x - 2}{4x - 3a}$
6. $\frac{25a^2 - b^2}{9a^2x^2 - 4x^2} \times \frac{x(3a + 2)}{5a + b}$
7. $\frac{x^2 + 5x + 6}{x^2 - 1} \times \frac{x^2 - 2x - 3}{x^2 - 9}$
8. $\frac{x^2 + 3x + 2}{x^2 + 9x + 20} \times \frac{x^2 + 7x + 12}{x^2 + 5x + 6}$
9. $\frac{2x^2 + 5x + 2}{x^2 - 4} \times \frac{x^2 + 4x}{2x^2 + 9x + 4}$
10. $\frac{2x^2 + 13x + 15}{4x^2 - 9} \div \frac{2x^2 + 11x + 5}{4x^2 - 1}$
11. $\frac{x^2 - 14x - 15}{x^2 - 4x - 45} \div \frac{x^2 - 12x - 45}{x^2 - 6x - 27}$
12. $\frac{2x^2 - x - 1}{2x^2 + 5x + 2} \times \frac{4x^2 + x - 14}{16x^2 - 49}$
13. $\frac{b^4 - 27b}{2b^2 + 5b} \times \frac{4b^2 - 25}{2b^2 - 11b + 15}$
14. $\frac{x^3 - 6x^2 + 36x}{x^2 - 49} \div \frac{x^4 + 216x}{x^2 - x - 42}$
15. $\frac{64p^2q^2 - z^2}{x^2 - 4} \div \frac{(x - 2)^2}{8pq + z^2} \times \frac{x^2 - 4}{(x + 2)^2}$
16. $\frac{x^3 - x - 20}{x^2 - 25} \times \frac{x^3 - x - 2}{x^2 + 2x - 8} \div \frac{x + 1}{x^2 + 5x}$
17. $\frac{x^2 - 18x + 80}{x^2 - 5x - 50} \times \frac{x^2 - 6x - 7}{x^2 - 15x + 56} \times \frac{x + 5}{x - 1}$
18. $\frac{x^2 - 8x - 9}{x^2 - 17x + 72} \times \frac{x^2 - 25}{x^2 - 1} \div \frac{x^3 + 4x - 5}{x^2 - 9x + 8}$

19. $\frac{4x^2 + x}{6xy}$

20. $\frac{x^2 + x}{x^2 - x}$

21. $\frac{4x^2 - 1}{2x^2 + 1}$

22. $\frac{x^4 - 8}{x^2 - 4x}$

23. $\frac{(a + b)^2}{a^2 + ab}$

24. $\frac{a^2 + 2a}{a^2 - b^2}$

25. $\frac{x^2 - 1}{x^2 + 24x}$

26. $\frac{(a^2 + a)}{a^2 - a}$

27. $\frac{m^3 + 4m}{3m^2 - 1}$

28. $\frac{1 + 8x^3}{(2 - x)^2}$

29. $\frac{x^2(x - 1)}{(x + 4)^2}$

30. $\frac{(p + q)^2}{(p + q)}$

31. $\frac{a^4 - 1}{a^2 - 2a}$

32. $\frac{a^3 + 8a}{(64a^3 - 1)}$

19. $\frac{4x^2+x-14}{6xy-14y} \times \frac{4x^2}{x^2-4} \times \frac{x-2}{4x-7} \div \frac{2x^2+4x}{3x^2-x-14}.$
20. $\frac{x^2+x-2}{x^2-x-20} \times \frac{x^2+5x+4}{x^2-x} \div \left(\frac{x^2+3x+2}{x^2-2x-15} \times \frac{x+3}{x^2} \right).$
21. $\frac{4x^2-16x+15}{2x^2+3x+1} \times \frac{x^2-6x-7}{2x^2-17x+21} \times \frac{4x^2-1}{4x^2-20x+25}.$
22. $\frac{x^4-8x}{x^3-4x-5} \times \frac{x^2+2x+1}{x^3-x^2-2x} \div \frac{x^2+2x+4}{x-5}.$
23. $\frac{(a+b)^2-c^2}{a^2+ab-ac} \times \frac{a}{(a+c)^2-b^2} \times \frac{(a-b)^2-c^2}{ab-b^2-bc}.$
24. $\frac{a^2+2ab+b^2-c^2}{a^2-b^2-c^2-2bc} \times \frac{a^2-2ac+c^2-b^2}{b^2-2bc+c^2-a^2}.$
25. $\frac{x^2-64}{x^2+24x+128} \times \frac{x^3+12x-64}{x^3-64} \div \frac{x^2-16x+64}{x^2+4x+16}.$
26. $\frac{(a^2+ax)^3}{a^2-x^2} \times \frac{(a-x)^3}{a^5+a^2x^3} \times \frac{a^2-ax+x^2}{a^3+2a^2x+ax^2}.$
27. $\frac{m^3+4m^2n+4mn^2}{3m^2n-5mn^2-2n^3} \times \frac{m^2-4n^2}{9m^2-3mn+n^2} \div \frac{(m+2n)^3}{27m^3+n^3}.$
28. $\frac{1+8x^3}{(2-x)^2} \times \frac{4x-x^3}{1-4x^2} \div \frac{(1-2x)^2+2x}{2-5x+2x^2}.$
29. $\frac{x^2(x-4)^2}{(x+4)^2-4x} \times \frac{64-x^3}{16-x^2} \div \frac{(x^2-4x)^3}{(x+4)^2}.$
30. $\frac{(p+q)^2-r^2}{(p+q+r)^2} \times \frac{p^2+pq+pr}{(p-r)^2-q^2} \div \frac{p^2-pq+pr}{(p-q)^2-r^2}.$
31. $\frac{a^4-x^4}{a^2-2ax+x^2} \div \left(\frac{a^2x+x^3}{a^3-x^3} \times \frac{a^4+a^2x^2+x^4}{a^2x-ax^2+x^3} \right).$
32. $\frac{a^3+8a^2b+15ab^2}{(64a^3-b^3)(a^3+b^3)} \times \frac{16a^4-17a^2b^2+b^4}{4a^2+21ab+5b^2} \div \frac{a^2+2ab-3b^2}{a^3-a^2b+ab^2}.$

CHAPTER XX.

LOWEST COMMON MULTIPLE.

159. DEFINITION. The lowest common multiple of two or more algebraical expressions is the expression of lowest dimensions, which is divisible by each of them without remainder.

In Chapter XI. we have explained how to write down by inspection the lowest common multiple of two or more *simple* expressions; the lowest common multiple of compound expressions which are given as the product of factors, or which can be easily resolved into factors, can be readily found by a similar method.

Example 1. The lowest common multiple of $6x^3(a-x)^2$, $8a^3(a-x)^3$ and $12ax(a-x)^5$ is $24a^3x^3(a-x)^5$.

For it consists of the product of

- (1) the L.C.M. of the numerical coefficients;
- (2) the lowest power of each factor which is divisible by every power of that factor occurring in the given expressions.

Example 2. Find the lowest common multiple of

$$3a^2 + 9ab, \quad 2a^3 - 18ab^2, \quad a^3 + 6a^2b + 9ab^2.$$

$$3a^2 + 9ab = 3a(a + 3b),$$

$$2a^3 - 18ab^2 = 2a(a + 3b)(a - 3b),$$

$$a^3 + 6a^2b + 9ab^2 = a(a + 3b)(a + 3b)$$

$$= a(a + 3b)^2.$$

Therefore the L.C.M. is $6a(a + 3b)^2(a - 3b)$.

EXAMPLES XX. a.

Find the lowest common multiple of

- | | | |
|-----------------------------------|-----------------------------------|--------------------------|
| 1. $x, x^2 + x.$ | 2. $x^2, x^2 - 3x.$ | 3. $3x^2, 4x^2 + 8x.$ |
| 4. $21x^2, 7x^2(x + 1).$ | 5. $x^2 - 1, x^2 + x.$ | 6. $a^2 + ab, ab + b^2.$ |
| 7. $4x^2y - y, 2x^2 + x.$ | 8. $6x^2 - 2x, 9x^2 - 3x.$ | |
| 9. $x^3 + 2x, x^2 + 3x + 2.$ | 10. $x^2 - 3x + 2, x^2 - 1.$ | |
| 11. $x^2 + 4x + 4, x^2 + 5x + 6.$ | 12. $x^2 - 5x + 4, x^2 - 6x + 8.$ | |

13. $x^2 - x -$
14. $x^2 + x -$
15. $x^2 + x -$
16. $2x^2 + 3x$
17. $3x^2 + 11$
18. $5x^2 + 11$
19. $2x^2 + 3x$
20. $3x^2 - x$
21. $12x^2 + 3$
22. $3x^4 + 26$
23. $60x^4 + 5$
24. $8x^2 - 38$
25. $12x^2 - 2$
26. $6ax^2 + 7$
27. $4ax^2y^2 +$
28. $(3x - 5x)$
29. $14a^4(a^3 +$
30. $m^4 + m^2y$
31. $(2c^2 - 3c$

160. Which cannot be divided by finding the lowest

Example.

$$2x^4 +$$

The highest

By division

$$2x^4 -$$

$$2x^4 +$$

Therefore the

*161. Which the lowest common multiple of the given expressions.

Let A and B be the common factors of the given expressions. Then the common factor of the quotients will be A/B . Therefore, since the common factor of the given expressions is A/B , the common factor of the quotients will be A/B .

E.A.

13. $x^2 - x - 6$, $x^2 + x - 2$, $x^2 - 4x + 3$.
14. $x^2 + x - 20$, $x^2 - 10x + 24$, $x^2 - x - 30$.
15. $x^2 + x - 42$, $x^2 - 11x + 30$, $x^2 + 2x - 35$.
16. $2x^2 + 3x + 1$, $2x^2 + 5x + 2$, $x^2 + 3x + 2$.
17. $3x^2 + 11x + 6$, $3x^2 + 8x + 4$, $x^2 + 5x + 6$.
18. $5x^2 + 11x + 2$, $5x^2 + 16x + 3$, $x^2 + 5x + 6$.
19. $2x^2 + 3x - 2$, $2x^2 + 15x - 8$, $x^2 + 10x + 16$.
20. $3x^2 - x - 14$, $3x^2 - 13x + 14$, $x^2 - 4$.
21. $12x^2 + 3x - 42$, $12x^2 + 30x^2 + 12x$, $32x^2 - 40x - 28$.
22. $3x^4 + 26x^3 + 35x^2$, $6x^3 + 38x - 28$, $27x^3 + 27x^2 - 30x$.
23. $60x^4 + 5x^3 - 5x^2$, $60x^2y + 32xy + 4y$, $40x^3y - 2x^2y - 2xy$.
24. $8x^2 - 38xy + 35y^2$, $4x^2 - xy - 5y^2$, $2x^2 - 5xy - 7y^2$.
25. $12x^2 - 23xy + 10y^2$, $4x^2 - 9xy + 5y^2$, $3x^2 - 5xy + 2y^2$.
26. $6ax^3 + 7a^2x^2 - 3a^3x$, $3a^2x^2 + 14a^3x - 5a^4$, $6x^2 + 39ax + 45a^2$.
27. $4ax^2y^2 + 11axy^2 - 3ay^2$, $3x^3y^3 + 7x^2y^2 - 6xy^3$, $24ax^2 - 22ax + 4a$.
28. $(3x - 5x^2)^2$, $6 - 7x - 5x^2$, $4x + 4x^2 + x^3$.
29. $14a^4(a^3 - b^3)$, $21a^2b^2(a - b)^3$, $6a^3b(a - b)(a^2 - b^2)$.
30. $m^4 + m^2n^2 + n^4$, $m^2n + n^4$, $(m^2 - mn)^3$.
31. $(2c^2 - 3cd)^2$, $(4c - 6d)^2$, $8c^3 - 27d^3$.

160. When the given expressions are such that their factors cannot be determined by inspection, they must be resolved by finding the highest common factor.

Example. Find the lowest common multiple of

$$2x^4 + x^3 - 20x^2 - 7x + 24 \text{ and } 2x^4 + 3x^3 - 13x^2 - 7x + 15.$$

The highest common factor is $x^2 + 2x - 3$.

By division, we obtain

$$2x^4 + x^3 - 20x^2 - 7x + 24 = (x^2 + 2x - 3)(2x^2 - 3x - 8).$$

$$2x^4 + 3x^3 - 13x^2 - 7x + 15 = (x^2 + 2x - 3)(2x^2 - x - 5).$$

Therefore the L.C.M. is $(x^2 + 2x - 3)(2x^2 - 3x - 8)(2x^2 - x - 5)$.

*161. We may now give the proof of the rule for finding the lowest common multiple of two compound algebraical expressions.

Let A and B be the two expressions, and F their highest common factor. Also suppose that a and b are the respective quotients when A and B are divided by F ; then $A = aF$, $B = bF$. Therefore, since a and b have no common factor, the lowest common multiple of A and B is abF , by inspection,

E.A.

K

***162.** There is an important relation between the highest common factor and the lowest common multiple of two expressions which it is desirable to notice.

Let F be the highest common factor, and X the lowest common multiple of A and B . Then, as in the preceding article,

$$A = aF, \quad B = bF,$$

and

$$X = abF.$$

Therefore the product

$$AB = aF \cdot bF$$

$$= F \cdot abF$$

$$= FX \dots\dots\dots(1).$$

Hence the product of two expressions is equal to the product of their highest common factor and lowest common multiple.

$$\text{Again, from (1) } X = \frac{AB}{F} = \frac{A}{F} \times B = \frac{B}{F} \times A;$$

hence the lowest common multiple of two expressions may be found by dividing their product by their highest common factor; or by dividing either of them by their highest common factor, and multiplying the quotient by the other.

***163.** The lowest common multiple of three expressions A, B, C may be obtained as follows.

First, find X the L.C.M. of A and B . Next find Y the L.C.M. of X and C ; then Y will be the required L.C.M. of A, B, C .

For Y is the expression of lowest dimensions which is divisible by X and C , and X is the expression of lowest dimensions divisible by A and B . Therefore Y is the expression of lowest dimensions divisible by all three.

EXAMPLES XX. b.

1. Find the highest common factor and the lowest common multiple of $x^2 - 5x + 6$, $x^2 - 4$, $x^3 - 3x - 2$.

2. Find the lowest common multiple of

$$ab(x^2 + 1) + x(a^2 + b^2) \text{ and } ab(x^2 - 1) + x(a^2 - b^2).$$

3. Find the lowest common multiple of

$$xy - bx, \quad xy - ay, \quad y^2 - 3by + 2b^2, \quad xy - 2bx - ay + 2ab,$$

$$xy - bx - ay + ab.$$

xx.]

4. Find multiple of

5. Find

6. Find

7. Find multiple of

8. Find

9. Find

Also find th

10. Find

Also shew three quanti

11. Find

12. Find multiple of

13. Find

14. Find

15. Find multiple of

16. Find multiple of

17. Find

18. Find

4. Find the highest common factor and the lowest common multiple of $x^3 + 2x^2 - 3x$, $2x^3 + 5x^2 - 3x$.

5. Find the lowest common multiple of
 $1 - x$, $(1 - x^2)^2$, $(1 + x)^3$.

6. Find the lowest common multiple of
 $x^2 - 10x + 24$, $x^2 - 8x + 12$, $x^2 - 6x + 8$.

7. Find the highest common factor and the lowest common multiple of $6x^3 + x^2 - 5x - 2$, $6x^3 + 5x^2 - 3x - 2$.

8. Find the lowest common multiple of
 $(bc^2 - abc)^2$, $b^2(ac^2 - a^3)$, $a^2c^2 + 2ac^3 + c^4$.

9. Find the lowest common multiple of
 $x^3 - y^3$, $x^3y - y^4$, $y^2(x - y)^3$, $x^2 + xy + y^2$.

Also find the highest common factor of the first three expressions.

10. Find the highest common factor of
 $6x^2 - 13x + 6$, $2x^2 + 5x - 12$, $6x^2 - x - 12$.

Also shew that the lowest common multiple is the product of the three quantities divided by the square of the highest common factor.

11. Find the lowest common multiple of
 $x^4 + ax^3 + a^2x + a^4$, $x^4 + a^2x^2 + a^4$.

12. Find the highest common factor and the lowest common multiple of $3x^3 - 7x^2y + 5xy^2 - y^3$, $x^2y + 3xy^2 - 3x^3 - y^3$,
 $3x^3 + 5x^2y + xy^2 - y^3$.

13. Find the highest common factor of
 $4x^3 - 10x^2 + 4x + 2$, $3x^4 - 2x^3 - 3x + 2$.

14. Find the lowest common multiple of
 $a^2 - b^2$, $a^3 - b^3$, $a^3 - a^2b - ab^2 - 2b^3$.

15. Find the highest common factor and the lowest common multiple of $(2x^2 - 3a^2)y + (2a^2 - 3y^2)x$, $(2a^2 + 3y^2)x + (2x^2 + 3a^2)y$.

16. Find the highest common factor and the lowest common multiple of $x^3 - 9x^2 + 26x - 24$, $x^3 - 12x^2 + 47x - 60$.

17. Find the highest common factor of
 $x^3 - 15ax^2 + 48a^2x + 64a^3$, $x^3 - 10ax + 16a^2$.

18. Find the lowest common multiple of
 $21x(xy - y^2)^2$, $35(x^4y^2 - x^2y^4)$, $15y(x^2 + xy)^2$.

CHAPTER XXI.

ADDITION AND SUBTRACTION OF FRACTIONS.

164. HAVING explained the rules for finding the lowest common multiple of any given expressions, we now proceed to shew how the addition and subtraction of fractions may be effected.

165. To prove $\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}$.

We have $\frac{a}{b} = \frac{ad}{bd}$ and $\frac{c}{d} = \frac{bc}{bd}$ [Art. 150].

Thus in each case we divide the unit into bd equal parts, and we take first ad of these parts, and then bc of them; that is, we take $ad+bc$ of the bd parts of the unit; and this is expressed by the fraction $\frac{ad+bc}{bd}$.

$$\therefore \frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{bd}.$$

Similarly, $\frac{a}{b} - \frac{c}{d} = \frac{ad-bc}{bd}$.

166. Here the fractions have been both expressed with a common denominator bd . But if b and d have a common factor, the product bd is not the lowest common denominator, and the fraction $\frac{ad+bc}{bd}$ will not be in its lowest terms. To avoid working with fractions which are not in their lowest terms, some modification of the above will be necessary. In practice it will be found advisable to take the *lowest* common denominator, which is the lowest common multiple of the denominators of the given fractions.

xxi.] AD

Rule. I.
denomina
take it for
of the first
the quotient
fractions.

Example.

The lower
We must
respectively
Hence the

167. W
subtraction

Rule II.
lowest comm
numerators,

Example 1

The lowest
Therefore

Example 2

The lowest
Thus the e

since the term

Note. To
brackets as in

Rule. I. To reduce fractions to their lowest common denominator: find the L.C.M. of the given denominators, and take it for the common denominator; divide it by the denominator of the first fraction, and multiply the numerator of this fraction by the quotient so obtained; and do the same with all the other given fractions.

Example. Express with lowest common denominator

$$\frac{5x}{2a(x-a)} \text{ and } \frac{4a}{3x(x^2-a^2)}.$$

The lowest common denominator is $6ax(x-a)(x+a)$.

We must therefore multiply the numerators by $3x(x+a)$ and $2a$ respectively.

Hence the equivalent fractions are

$$\frac{15x^2(x+a)}{6ax(x-a)(x+a)} \text{ and } \frac{8a^2}{6ax(x-a)(x+a)}.$$

167. We may now enunciate the rule for the addition or subtraction of fractions.

Rule II. To add or subtract fractions: reduce them to the lowest common denominator; find the algebraical sum of the numerators, and retain the common denominator.

Example 1. Find the value of $\frac{2x+a}{3a} + \frac{5x-4a}{9a}$.

The lowest common denominator is $9a$.

$$\begin{aligned} \text{Therefore the expression} &= \frac{3(2x+a) + 5x - 4a}{9a} \\ &= \frac{6x + 3a + 5x - 4a}{9a} = \frac{11x - a}{9a}. \end{aligned}$$

Example 2. Find the value of $\frac{x-2y}{xy} + \frac{3y-a}{ay} - \frac{3x-2a}{ax}$.

The lowest common denominator is axy .

$$\begin{aligned} \text{Thus the expression} &= \frac{a(x-2y) + x(3y-a) - y(3x-2a)}{axy} \\ &= \frac{ax - 2ay + 3xy - ax - 3xy + 2ay}{axy} \\ &= 0, \end{aligned}$$

since the terms in the numerator destroy each other.

Note. To ensure accuracy the beginner is recommended to use brackets as in the first line of work above.

EXAMPLES XXI. a.

Find the value of

1. $\frac{x-1}{2} + \frac{x+3}{5} + \frac{x+7}{10}$.
2. $\frac{2x-1}{3} + \frac{x-5}{6} + \frac{x-4}{4}$.
3. $\frac{5x-1}{8} - \frac{3x-2}{7} + \frac{x-5}{4}$.
4. $\frac{2x-3}{9} - \frac{x+2}{6} + \frac{5x+8}{12}$.
5. $\frac{x-7}{15} + \frac{x-9}{25} - \frac{x+3}{45}$.
6. $\frac{2x+5}{x} - \frac{x+3}{2x} - \frac{27}{8x^3}$.
7. $\frac{a-b}{ab} + \frac{b-c}{bc} + \frac{c-a}{ca}$.
8. $\frac{a-2b}{2a} - \frac{a-5b}{4a} + \frac{a+7b}{8a}$.
9. $\frac{b+c}{2a} + \frac{c+a}{4b} - \frac{a-b}{3c}$.
10. $\frac{a-x}{x} + \frac{a+x}{a} - \frac{a^2-x^2}{2ax}$.
11. $\frac{x+2}{17x} - \frac{x-5}{34x} + \frac{x+2}{51x}$.
12. $\frac{2a^2-b^2}{a^2} - \frac{b^2-c^2}{b^2} - \frac{c^2-a^2}{c^2}$.
13. $\frac{x-3}{5x} + \frac{x^2-9}{10x^2} - \frac{8-x^3}{15x^3}$.
14. $\frac{2}{xy} - \frac{3y^2-x^2}{xy^2} + \frac{xy+y^2}{x^2y^2}$.
15. $\frac{2x-3y}{xy} + \frac{3x-2z}{xz} + \frac{5}{x}$.
16. $\frac{a^2-bc}{bc} - \frac{ac-b^2}{ac} - \frac{ab-c^2}{ab}$.

Example 3. Simplify $\frac{2x-3a}{x-2a} - \frac{2x-a}{x-a}$.

The lowest common denominator is $(x-2a)(x-a)$.

Hence, multiplying the numerators by $x-a$ and $x-2a$ respectively, we have

$$\begin{aligned}
 \text{the expression} &= \frac{(2x-3a)(x-a) - (2x-a)(x-2a)}{(x-2a)(x-a)} \\
 &= \frac{2x^2 - 5ax + 3a^2 - (2x^2 - 5ax + 2a^2)}{(x-2a)(x-a)} \\
 &= \frac{2x^2 - 5ax + 3a^2 - 2x^2 + 5ax - 2a^2}{(x-2a)(x-a)} \\
 &= \frac{a^2}{(x-2a)(x-a)}.
 \end{aligned}$$

Note. In finding the value of such an expression as $-(2x-a)(x-2a)$,

the beginner should first express the product in brackets, and then remove the brackets, as we have done. After a little practice he will be able to take both steps together.

The work will sometimes be shortened by first reducing the fractions to their lowest terms.

Example

The

Find the

1. $\frac{1}{x+2} +$
4. $\frac{3}{x-6} -$
7. $\frac{x+3}{x+4} -$
10. $\frac{x-4}{x-2} -$
13. $\frac{1}{2x-3} -$
15. $\frac{4a^2+b}{4a^2-b} -$
17. $\frac{x^2}{x-x^3} -$
19. $\frac{xy}{25x^2} -$
21. $\frac{x^2-4c}{x^2-2a} -$
23. $\frac{1}{a-2x} -$
25. $\frac{3}{x^2-4} -$

Example 4. Simplify $\frac{x^2+5xy-4y^2}{x^2-16y^2} - \frac{2xy}{2x^2+8xy}$.

$$\begin{aligned}\text{The expression} &= \frac{x^2+5xy-4y^2}{x^2-16y^2} - \frac{y}{x+4y} \\ &= \frac{x^2+5xy-4y^2-y(x-4y)}{x^2-16y^2} \\ &= \frac{x^2+5xy-4y^2-xy+4y^2}{x^2-16y^2} \\ &= \frac{x^2+4xy}{x^2-16y^2} = \frac{x}{x-4y}.\end{aligned}$$

EXAMPLES XXI. b.

Find the value of

1. $\frac{1}{x+2} + \frac{1}{x+3}$.
2. $\frac{2}{x+3} - \frac{1}{x+4}$.
3. $\frac{1}{x-5} - \frac{1}{x-4}$.
4. $\frac{3}{x-6} - \frac{1}{x+2}$.
5. $\frac{a}{x+a} - \frac{b}{x+b}$.
6. $\frac{a}{x-a} + \frac{b}{x-b}$.
7. $\frac{x+3}{x+4} - \frac{x+1}{x+2}$.
8. $\frac{a+x}{a-x} - \frac{a-x}{a+x}$.
9. $\frac{x+2}{x-2} - \frac{x-2}{x+2}$.
10. $\frac{x-4}{x-2} - \frac{x-7}{x-5}$.
11. $\frac{a}{x-a} - \frac{a^2}{x^2-a^2}$.
12. $\frac{3}{x-3} + \frac{2x}{x^2-9}$.
13. $\frac{1}{2x-3y} - \frac{x+y}{4x^2-9y^2}$.
14. $\frac{x+a}{x-2a} - \frac{x^2+2a^2}{x^2-4a^2}$.
15. $\frac{4a^2+b^2}{4a^2-b^2} - \frac{2a-b}{2a+b}$.
16. $\frac{2x^2}{x^2-y^2} - \frac{2x^2}{x^2+xy}$.
17. $\frac{x^2}{x-x^3} - \frac{x}{1+x^2}$.
18. $\frac{1}{x(x-y)} + \frac{1}{y(x+y)}$.
19. $\frac{xy}{25x^2-y^2} + \frac{2x^2y}{10x^2y+2xy^2}$.
20. $\frac{y}{x(x^2-y^2)} + \frac{x}{y(x^2+y^2)}$.
21. $\frac{x^2-4a^2}{x^2-2ax} - \frac{x^3+2ax-8a^3}{x^2-4a^2}$.
22. $\frac{x^2+xy+y^2}{x+y} + \frac{x^2-xy+y^2}{x-y}$.
23. $\frac{1}{a-2x} - \frac{(a+2x)^2}{a^3-8x^3}$.
24. $\frac{a^3+b^3}{a^3-ab+b^2} - \frac{a^3-b^3}{a^3+ab+b^2}$.
25. $\frac{3}{x^2-4} + \frac{1}{(x-2)^2}$.
26. $\frac{1}{a(x^2-a^2)} - \frac{1}{x(x+a)^2}$.

168. Some modification of the foregoing general methods may sometimes be used with advantage. The most useful artifices are explained in the examples which follow, but no general rules can be given which will apply to all cases.

Example 1. Simplify $\frac{a+3}{a-4} - \frac{a+4}{a-3} - \frac{8}{a^2-16}$.

Taking the first two fractions together, we have

$$\begin{aligned} \text{the expression} &= \frac{a^2-9-(a^2-16)}{(a-4)(a-3)} - \frac{8}{a^2-16} \\ &= \frac{7}{(a-4)(a-3)} - \frac{8}{(a+4)(a-4)} \\ &= \frac{7(a+4)-8(a-3)}{(a+4)(a-4)(a-3)} \\ &= \frac{52-a}{(a+4)(a-4)(a-3)} \end{aligned}$$

Example 2. Simplify $\frac{1}{2x^2+x-1} + \frac{1}{3x^2+4x+1}$.

$$\begin{aligned} \text{The expression} &= \frac{1}{(2x-1)(x+1)} + \frac{1}{(3x+1)(x+1)} \\ &= \frac{3x+1+2x-1}{(2x-1)(x+1)(3x+1)} \\ &= \frac{5x}{(2x-1)(x+1)(3x+1)} \end{aligned}$$

Example 3. Simplify $\frac{1}{a-x} - \frac{1}{a+x} - \frac{2x}{a^2+x^2} - \frac{4x^3}{a^4+x^4}$.

Here it should be evident that the first two denominators give L.C.M. a^2-x^2 , which readily combines with a^2+x^2 to give L.C.M. a^4-x^4 , which again combines with a^4+x^4 to give L.C.M. a^8-x^8 . Hence it will be convenient to proceed as follows:

$$\begin{aligned} \text{The expression} &= \frac{a+x-(a-x)}{a^2-x^2} - \dots - \dots \\ &= \frac{2x}{a^2-x^2} - \frac{2x}{a^2+x^2} - \dots \\ &= \frac{4x^3}{a^4-x^4} - \frac{4x^3}{a^4+x^4} \\ &= \frac{8x^7}{a^8-x^8} \end{aligned}$$

Find the

1. $\frac{1}{x+y}$

3. $\frac{5}{1+2x}$

5. $\frac{10}{9-a^2}$

7. $\frac{1}{2(a-b)}$

9. $\frac{3}{x-2}$

11. $\frac{1}{x^2-9x}$

13. $\frac{1}{2x^2-x}$

15. $\frac{4}{4-7a}$

17. $\frac{1}{x+1}$

18. $\frac{5}{2(x+1)}$

19. $\frac{a}{4(a+b)}$

20. $\frac{2}{x^2-3x}$

21. $\frac{x}{x^2+5x}$

22. $\frac{3}{x^2-1}$

23. $\frac{5(2x)}{11(6x^2)}$

24. $\frac{x-3}{x+2}$

EXAMPLES XXI. c.

Find the value of

1. $\frac{1}{x+y} - \frac{1}{x-y} + \frac{2x}{x^2-y^2}$

2. $\frac{1}{2x+y} + \frac{1}{2x-y} - \frac{3x}{4x^2-y^2}$

3. $\frac{5}{1+2x} - \frac{3x}{1-2x} - \frac{4-13x}{1-4x^2}$

4. $\frac{2a}{2a+3b} + \frac{3b}{2a-3b} - \frac{8b^2}{4a^2-9b^2}$

5. $\frac{10}{9-a^2} - \frac{2}{3+a} - \frac{1}{3-a}$

6. $\frac{5x}{6(x^2-1)} - \frac{1}{2(x-1)} + \frac{1}{3(x+1)}$

7. $\frac{1}{2(a-b)} - \frac{1}{2(a+b)} - \frac{b}{a^2-b^2}$

8. $\frac{2a}{2a-3} - \frac{5}{6a+9} - \frac{4(3a+2)}{3(4a^2-9)}$

9. $\frac{3}{x-2} + \frac{2}{3x+6} + \frac{5x}{x^2-4}$

10. $\frac{x}{x^3+y^3} - \frac{y}{x^3-y^3} + \frac{x^2y+xy^2}{x^6-y^6}$

11. $\frac{1}{x^2-9x+20} + \frac{1}{x^2-11x+30}$

12. $\frac{1}{x^2-7x+12} - \frac{1}{x^2-5x+6}$

13. $\frac{1}{2x^2-x-1} - \frac{1}{2x^2+x-3}$

14. $\frac{1}{2x^2-x-1} - \frac{3}{6x^2-x-2}$

15. $\frac{4}{4-7a-2a^2} - \frac{3}{3-a-10a^2}$

16. $\frac{5}{5+x-18x^2} - \frac{2}{2+5x+2x^2}$

17. $\frac{1}{x+1} - \frac{1}{(x+1)(x+2)} + \frac{1}{(x+1)(x+2)(x+3)}$

18. $\frac{5x}{2(x+1)(x-3)} - \frac{15(x-1)}{16(x-3)(x-2)} - \frac{9(x+3)}{16(x+1)(x-2)}$

19. $\frac{a+3b}{4(a+b)(a+2b)} + \frac{a+2b}{(a+b)(a+3b)} - \frac{a+b}{4(a+2b)(a+3b)}$

20. $\frac{2}{x^2-3x+2} + \frac{2}{x^2-x-2} - \frac{1}{x^2-1}$

21. $\frac{x}{x^2+5x+6} + \frac{15}{x^2+9x+14} - \frac{12}{x^2+10x+21}$

22. $\frac{3}{x^2-1} + \frac{4}{2x+1} + \frac{4x+2}{2x^2+3x+1}$

23. $\frac{5(2x-3)}{11(6x^2+x-1)} + \frac{7x}{6x^2+7x-3} - \frac{12(3x+1)}{11(4x^2+8x+3)}$

24. $\frac{x-3}{x+2} - \frac{x-2}{x+3} + \frac{1}{x-1}$

25. $\frac{x-3}{x-4} - \frac{x+4}{x+3} - \frac{5}{x^2-16}$

Find the value of

$$26. \frac{1+2a}{1-2a} - \frac{1-2a}{1+2a} - \frac{8a}{(1-2a)^2}$$

$$28. \frac{1}{3-x} - \frac{1}{3+x} - \frac{2x}{9+x^2}$$

$$30. \frac{1}{4(1+x)} + \frac{1}{4(1-x)} + \frac{1}{2(1+x^2)}$$

$$32. \frac{2x}{4+x^2} + \frac{1}{2-x} - \frac{1}{2+x}$$

$$34. \frac{1}{2a-8x} - \frac{a}{3a^2+48x^2} + \frac{1}{2a+8x}$$

$$35. \frac{1}{6a^2+54} + \frac{1}{3a-9} - \frac{a}{3a^2-27}$$

$$36. \frac{1}{8-8x} - \frac{1}{8+8x} + \frac{x}{4+4x^2} - \frac{x}{2+2x^4}$$

$$37. \frac{1}{6a-18} - \frac{1}{6a+18} - \frac{1}{a^2+9} + \frac{18}{a^4+81}$$

$$38. \frac{x+1}{2x^3-4x^2} + \frac{x-1}{2x^3+4x^2} - \frac{1}{x^2-4}$$

$$39. \frac{1}{3x^2-4xy+y^2} + \frac{1}{x^2-4xy+3y^2} - \frac{3}{3x^2-10xy+3y^2}$$

$$40. \frac{1}{x-1} + \frac{2}{x+1} - \frac{3x-2}{x^2-1} - \frac{1}{(x+1)^2}$$

$$41. \frac{108-52x}{x(3-x)^2} - \frac{4}{3-x} - \frac{12}{x} + \left(\frac{1+x}{3-x}\right)^2$$

$$42. \frac{(a+b)^2}{(x-a)(x+a+b)} - \frac{a+2b+x}{2(x-a)} + \frac{(a+b)x}{x^2+bx-a^2-ab} + \frac{1}{2}$$

$$43. \frac{3(x^2+x-2)}{x^3-x-2} - \frac{3(x^2-x-2)}{x^3+x-2} - \frac{8x}{x^2-4}$$

169. We have thus far assumed both numerator and denominator to be positive integers, and have shewn in Art. 155 that a fraction itself is the quotient resulting from the division of the numerator by denominator. But in algebra division is a process not restricted to positive integers, and we shall extend this definition as follows:

The algebraic fraction $\frac{a}{b}$ is the quotient resulting from the division of a by b , where a and b may have any values whatever.

170. B
from the d
 a by b , and

Again,
by b ; and
of signs, p

Therefor

Likewise

a by $-b$;
rule of sign

Therefor

These re

1. If th
fraction be
changed.

2. If th
changed, th

The prim
reduction
which will

1. We m
denominator

2. We m
the sign of

Example 1

Example 2

Example 3

The inter

170. By the preceding article $\frac{-a}{-b}$ is the quotient resulting from the division of $-a$ by $-b$; and this is obtained by dividing a by b , and, by the rule of signs, prefixing $+$.

Again, $\frac{-a}{b}$ is the quotient resulting from the division of $-a$ by b ; and this is obtained by dividing a by b , and, by the rule of signs, prefixing $-$.

Therefore
$$\frac{-a}{b} = -\frac{a}{b} \dots\dots\dots(2).$$

Likewise $\frac{a}{-b}$ is the quotient resulting from the division of a by $-b$; and this is obtained by dividing a by b , and, by the rule of signs, prefixing $-$.

Therefore
$$\frac{a}{-b} = -\frac{a}{b} \dots\dots\dots(3).$$

These results may be enunciated as follows :

1. *If the signs of BOTH numerator and denominator of a fraction be changed, the sign of the whole fraction will be unchanged.*

2. *If the sign of EITHER numerator or denominator alone be changed, the sign of the whole fraction will be changed.*

The principles here involved are so useful in certain cases of reduction of fractions that we quote them in another form, which will sometimes be found more easy of application.

1. *We may change the sign of every term in the numerator and denominator of a fraction without altering its value.*

2. *We may change the sign of a fraction by simply changing the sign of every term in EITHER the numerator or denominator.*

Example 1.
$$\frac{b-a}{y-x} = \frac{-b+a}{-y+x} = \frac{a-b}{x-y}.$$

Example 2.
$$\frac{x-x^2}{2y} = -\frac{-x+x^2}{2y} = -\frac{x^2-x}{2y}.$$

Example 3.
$$\frac{3x}{4-x^2} = -\frac{3x}{-4+x^2} = -\frac{3x}{x^2-4}.$$

The intermediate step may usually be omitted.

Example 4. Simplify $\frac{a}{x+a} + \frac{2x}{x-a} + \frac{a(3x-a)}{a^2-x^2}$.

Here it is evident that the lowest common denominator of the first two fractions is $x^2 - a^2$, therefore it will be convenient to alter the sign of the denominator in the third fraction.

$$\begin{aligned}\text{Thus the expression} &= \frac{a}{x+a} + \frac{2x}{x-a} - \frac{a(3x-a)}{x^2-a^2} \\ &= \frac{a(x-a) + 2x(x+a) - a(3x-a)}{x^2-a^2} \\ &= \frac{ax - a^2 + 2x^2 + 2ax - 3ax + a^2}{x^2-a^2} \\ &= \frac{2x^2}{x^2-a^2}.\end{aligned}$$

Example 5. Simplify $\frac{5}{3x-3} + \frac{3x-1}{1-x^2} + \frac{1}{2x+3}$.

$$\begin{aligned}\text{The expression} &= \frac{5}{3(x-1)} - \frac{3x-1}{x^2-1} + \frac{1}{2(x+1)} \\ &= \frac{10(x+1) - 6(3x-1) + 3(x-1)}{6(x^2-1)} \\ &= \frac{10x+10-18x+6+3x-3}{6(x^2-1)} \\ &= \frac{13-5x}{6(x^2-1)}.\end{aligned}$$

EXAMPLES XXI. d.

Simplify

1. $\frac{1}{4x-4} - \frac{1}{5x+5} + \frac{1}{1-x^2}$
2. $\frac{3}{1+a} - \frac{2}{1-a} - \frac{5a}{a^2-1}$
3. $\frac{x-2a}{x+a} + \frac{2(a^2-4ax)}{a^2-x^2} - \frac{3a}{x-a}$
4. $\frac{x-a}{x+a} + \frac{a^2+3ax}{a^2-x^2} + \frac{x+a}{x-a}$
5. $\frac{1}{2x+1} + \frac{1}{2x-1} + \frac{4x}{1-4x^2}$
6. $\frac{3x}{1-x^2} - \frac{2}{x-1} - \frac{2}{x+1}$
7. $\frac{2-5x}{x+3} - \frac{3+x}{3-x} + \frac{2x(2x-11)}{x^2-9}$
8. $\frac{3-2x}{2x+3} - \frac{2x+3}{3-2x} + \frac{12}{4x^2-9}$
9. $\frac{5}{2b+2} - \frac{3}{4b-4} + \frac{11}{6-6b^2}$
10. $\frac{1}{6a+6} + \frac{1}{6-6a} - \frac{1}{3a^2-3}$

11. $\frac{y^3}{x^3-y^3}$

13. $\frac{x^2+y^2}{x^2-y^2}$

15. $\frac{1}{2a+5b}$

17. $\frac{ax^2+b}{2x-1}$

19. $\frac{a}{(a-b)}$

20. $\frac{2a}{(x-a)}$

21. $\frac{1}{(a^2-b^2)}$

22. $\frac{1}{x+a} +$

23. $\frac{3}{x+a}$

24. $\frac{1}{4a^3(a+)}$

25. $\frac{x}{x^2-y^2}$

26. $\frac{b}{a(a^2-)}$

27. $\frac{a^2-2a}{2(a^2-)}$

28. $\frac{2}{a+b}$

29. $\frac{3}{8(1-x)}$

30. $\frac{1}{x} + \frac{1}{x-}$

31. $\frac{a^2+ac}{a^2c-c^3}$

32. $\frac{4a+6b}{a+b}$

11. $\frac{y^3}{x^3-y^3} + \frac{x^2y^2}{y^6-x^6}$ 12. $\frac{x^3-y^3}{xy} - \frac{xy-y^2}{xy-x^2}$
13. $\frac{x^2+y^2}{x^2-y^2} + \frac{x}{x+y} + \frac{y}{y-x}$ 14. $\frac{x^2+2x+4}{x+2} - \frac{x^2-2x+4}{2-x}$
15. $\frac{1}{2a+5b} + \frac{3a}{25b^2-4a^2} + \frac{1}{2a-5b}$ 16. $\frac{2b-a}{x-b} - \frac{3x(a-b)}{b^2-x^2} + \frac{b-2a}{b+x}$
17. $\frac{ax^2+b}{2x-1} + \frac{2(bx+ax^2)}{1-4x^2} - \frac{ax^2-b}{2x+1}$ 18. $\frac{a+c}{(a-b)(x-a)} + \frac{b+c}{(b-a)(x-b)}$
19. $\frac{a-c}{(a-b)(x-a)} - \frac{b-c}{(b-a)(b-x)}$
20. $\frac{2a+y}{(x-a)(a-b)} + \frac{a+b+y}{(x-b)(b-a)} - \frac{x+y-a}{(x-a)(x-b)}$
21. $\frac{1}{(a^2-b^2)(x^2+b^2)} + \frac{1}{(b^2-a^2)(x^2+a^2)} - \frac{1}{(x^2+a^2)(x^2+b^2)}$
22. $\frac{1}{x+a} + \frac{4a}{x^2-a^2} + \frac{1}{a-x} - \frac{2a}{x^2+a^2}$
23. $\frac{3}{x+a} - \frac{1}{x+3a} + \frac{3}{a-x} + \frac{1}{x-3a}$
24. $\frac{1}{4a^3(a+x)} - \frac{1}{4a^3(x-a)} + \frac{1}{2a^2(a^2+x^2)} - \frac{a^4}{a^5-x^5}$
25. $\frac{x}{x^3-y^3} - \frac{y}{x^2+y^2} + \frac{x^2+y^2}{y^4-x^4} + \frac{xy}{(x+y)(x^2+y^2)}$
26. $\frac{b}{a(a^2-b^2)} + \frac{a}{b(a^2+b^2)} + \frac{a^4+b^4}{ab(b^4-a^4)} - \frac{a^6}{b^6-a^6}$
27. $\frac{a^3-2ax+x^2}{2(a^2-x^2)} - \frac{2ax(a+x)}{(a-x)(a^2+2ax+x^2)} - \frac{x^2-a^2}{2(x-a)^2}$
28. $\frac{2}{a+b} - \frac{1}{a-b} - \frac{3b}{b^3-a^3} + \frac{ab}{a^3+b^3}$
29. $\frac{3}{8(1-x)} + \frac{1}{8(1+x)} - \frac{1-x}{4(1+x^2)} - \frac{3}{4(x^2-1)}$
30. $\frac{1}{x} + \frac{1}{x-1} + \frac{1}{x+1} + \frac{x}{1-x^2} + \frac{3}{x(x^2-1)}$
31. $\frac{a^2+ac}{a^2c-c^3} - \frac{a^2-c^2}{a^2c+2ac^2+c^3} + \frac{2c}{c^2-a^2} - \frac{3}{a+c}$
32. $\frac{4a+6b}{a+b} + \frac{6a-4b}{a-b} + \frac{4a^2+6b^2}{b^2-a^2} + \frac{4b^2-6a^2}{a^2+b^2} - \frac{20b^4}{b^4-a^4}$

*171. Consider the expression

$$\frac{1}{(a-b)(a-c)} + \frac{1}{(b-c)(b-a)} + \frac{1}{(c-a)(c-b)}$$

Here in finding the L.C.M. of the denominators it must be observed that there are not *six* different compound factors to be considered; for three of them differ from the other three only in sign.

Thus

$$(a-c) = -(c-a),$$

$$(b-a) = -(a-b),$$

$$(c-b) = -(b-c).$$

Hence, replacing the second factor in each denominator by its equivalent, we may write the expression in the form

$$-\frac{1}{(a-b)(c-a)} - \frac{1}{(b-c)(a-b)} - \frac{1}{(c-a)(b-c)} \dots\dots\dots(1).$$

Now the L.C.M. is $(b-c)(c-a)(a-b)$;

$$\begin{aligned} \text{and the expression} &= \frac{-(b-c) - (c-a) - (a-b)}{(b-c)(c-a)(a-b)} \\ &= \frac{-b+c-c+a-a+b}{(b-c)(c-a)(a-b)} \\ &= 0. \end{aligned}$$

*172. There is a peculiarity in the arrangement of this example which it is desirable to notice. In the expression (1) the letters occur in what is known as **Cyclic Order**; that is, b follows a , a follows c , c follows b . Thus if a, b, c are arranged round the circumference of a circle, as in the annexed diagram, if we start from any letter and move round in the direction of the arrows, the other letters follow in cyclic order, namely, abc, bca, cab .

The observance of this principle is especially important in a large class of examples in which the differences of three letters are involved. Thus we are observing cyclic order when we write $b-c, c-a, a-b$; whereas we are violating cyclic order by the use of arrangements such as $b-c, a-c, a-b$, or $a-c, b-a, b-c$. It will always be found that the work is rendered shorter and easier by following cyclic order from the beginning, and adhering to it throughout the question.

In the present chapter we shall confine our attention to a few of the simpler cases, resuming the subject in Chapter XXIX.



Find the

1. $\frac{a}{(a-b)}$

2. $\frac{b}{(a-b)}$

3. $\frac{2}{(x-y)}$

4. $\frac{y}{(x-y)}$

5. $\frac{b}{(a-b)}$

6. $\frac{x^2}{(x-y)}$

7. $\frac{1}{(a-b)}$

8. $\frac{p}{(p-q)}$

9. $\frac{p+q}{(p-q)}$

10. $\frac{1}{(a^2-b^2)}$

11. $\frac{x}{(p-q)}$

12. $\frac{q}{(x-y)}$

***EXAMPLES XXI. e.**

Find the value of

$$1. \frac{a}{(a-b)(a-c)} + \frac{b}{(b-c)(b-a)} + \frac{c}{(c-a)(c-b)}.$$

$$2. \frac{b}{(a-b)(a-c)} + \frac{c}{(b-c)(b-a)} + \frac{a}{(c-a)(c-b)}.$$

$$3. \frac{z}{(x-y)(x-z)} + \frac{x}{(y-z)(y-x)} + \frac{y}{(z-x)(z-y)}.$$

$$4. \frac{y+z}{(x-y)(x-z)} + \frac{z+x}{(y-z)(y-x)} + \frac{x+y}{(z-x)(z-y)}.$$

$$5. \frac{b-c}{(a-b)(a-c)} + \frac{c-a}{(b-c)(b-a)} + \frac{a-b}{(c-a)(c-b)}.$$

$$6. \frac{x^2yz}{(x-y)(x-z)} + \frac{y^2zx}{(y-z)(y-x)} + \frac{z^2xy}{(z-x)(z-y)}.$$

$$7. \frac{1+a}{(a-b)(a-c)} + \frac{1+b}{(b-c)(b-a)} + \frac{1+c}{(c-a)(c-b)}.$$

$$8. \frac{p-a}{(p-q)(p-r)} + \frac{q-a}{(q-r)(q-p)} + \frac{r-a}{(r-p)(r-q)}.$$

$$9. \frac{p+q-r}{(p-q)(p-r)} + \frac{q+r-p}{(q-r)(q-p)} + \frac{r+p-q}{(r-p)(r-q)}.$$

$$10. \frac{a^2}{(a^2-b^2)(a^2-c^2)} + \frac{b^2}{(b^2-c^2)(b^2-a^2)} + \frac{c^2}{(c^2-a^2)(c^2-b^2)}.$$

$$11. \frac{x+y}{(p-q)(p-r)} + \frac{x+y}{(q-r)(q-p)} + \frac{x+y}{(r-p)(r-q)}.$$

$$12. \frac{q+r}{(x-y)(x-z)} + \frac{r+p}{(y-z)(y-x)} + \frac{p+q}{(z-x)(z-y)}.$$



CHAPTER XXII.

MISCELLANEOUS FRACTIONS.

173. We now propose to consider some miscellaneous questions concerning fractions of a more complicated kind than those already discussed.

In the previous chapters on Fractions, the numerator and denominator have been regarded as integers; but cases frequently occur in which the numerator or denominator of a fraction is itself fractional.

174. **Definition.** A fraction of which the numerator or denominator is itself a fraction is called a **Complex Fraction**.

Thus $\frac{\frac{a}{b}}{\frac{c}{d}}$, $\frac{\frac{a}{b}}{x}$, $\frac{\frac{a}{b}}{\frac{c}{d}}$ are Complex Fractions.

In the last of these types, the outside quantities, a and d , are sometimes referred to as the *extremes*, while the two middle quantities, b and c , are called the *means*.

175. Instead of using the horizontal line to separate numerator and denominator, it is sometimes convenient to write complex fractions in the forms

$$a/b, \quad a/x, \quad a/c$$

176. By definition (Art. 169) $\frac{\frac{a}{b}}{\frac{c}{d}}$ is the quotient resulting

from the division of $\frac{a}{b}$ by $\frac{c}{d}$; and this by Art. 158 is $\frac{ad}{bc}$;

$$\therefore \frac{\frac{a}{b}}{\frac{c}{d}} = \frac{ad}{bc}$$

177. H
of writing
Multipl
new denom

Example

by cancelli

178. T
cases, and

179. T
complex fr

Example

Simplification of Complex Fractions.

177. From the preceding article we deduce an easy method of writing down the simplified form of a complex fraction.

Multiply the extremes for a new numerator, and the means for a new denominator.

Example.
$$\frac{\frac{a+x}{b}}{\frac{a^2-x^2}{ab}} = \frac{ab(a+x)}{b(a^2-x^2)} = \frac{a}{a-x},$$

by cancelling common factors in numerator and denominator.

178. The student should especially notice the following cases, and should be able to write down the results readily.

$$\frac{1}{\frac{a}{b}} = 1 \div \frac{a}{b} = 1 \times \frac{b}{a} = \frac{b}{a},$$

$$\frac{\frac{a}{1}}{\frac{1}{b}} = a \div \frac{1}{b} = a \times b = ab.$$

$$\frac{\frac{1}{a}}{\frac{1}{b}} = \frac{1}{a} \div \frac{1}{b} = \frac{1}{a} \times \frac{b}{1} = \frac{b}{a}.$$

179. The following examples illustrate the simplification of complex fractions.

Example 1.
$$\frac{\frac{a}{b} + \frac{c}{d}}{\frac{a}{b} - \frac{c}{d}} = \left(\frac{a}{b} + \frac{c}{d} \right) \div \left(\frac{a}{b} - \frac{c}{d} \right)$$

$$= \frac{ad+bc}{bd} \div \frac{ad-bc}{bd}$$

$$= \frac{ad+bc}{bd} \times \frac{bd}{ad-bc}$$

$$= \frac{ad+bc}{ad-bc}$$

Example 2.
$$\frac{x + \frac{a^2}{x}}{x - \frac{a^2}{x}} = \left(x + \frac{a^2}{x}\right) \div \left(x - \frac{a^2}{x}\right) = \frac{x^2 + a^2}{x} \div \frac{x^2 - a^2}{x}$$

$$= \frac{x^2 + a^2}{x} \times \frac{x^2}{x^2 - a^2} = \frac{x^2}{x^2 - a^2}.$$

Example 3. Simplify
$$\frac{\frac{3}{a} + \frac{a}{3} - 2}{\frac{a}{6} + \frac{1}{2} - \frac{3}{a}}.$$

Here the reduction may be simply effected by multiplying the fractions above and below by $6a$, which is the L.C.M. of the denominators.

Thus the expression =
$$\frac{18 + 2a^2 - 12a}{a^2 + 3a - 18}$$

$$= \frac{2(a^2 - 6a + 9)}{(a+6)(a-3)} = \frac{2(a-3)}{a+6}.$$

Example 4. Simplify
$$\frac{\frac{a^2 + b^2}{a^2 - b^2} - \frac{a^2 - b^2}{a^2 + b^2}}{\frac{a+b}{a-b} - \frac{a-b}{a+b}}.$$

The numerator =
$$= \frac{(a^2 + b^2)^2 - (a^2 - b^2)^2}{(a^2 + b^2)(a^2 - b^2)}$$

$$= \frac{4a^2b^2}{(a^2 + b^2)(a^2 - b^2)}.$$

Similarly the denominator =
$$= \frac{4ab}{(a+b)(a-b)}.$$

Hence the fraction =
$$= \frac{4a^2b^2}{(a^2 + b^2)(a^2 - b^2)} \div \frac{4ab}{(a+b)(a-b)}$$

$$= \frac{4a^2b^2}{(a^2 + b^2)(a^2 - b^2)} \times \frac{(a+b)(a-b)}{4ab}$$

$$= \frac{ab}{a^2 + b^2}.$$

Note. To ensure accuracy and neatness, when the numerator and denominator are somewhat complicated, the beginner is advised to simplify each separately as in the above example.

180. Continue
simplify

Example

The exp

Find the

1.
$$\frac{\frac{m}{n} - \frac{l}{m}}{\frac{a}{m} - \frac{n}{n}}$$

5.
$$\frac{2 + \frac{3a}{4b}}{a + \frac{3}{3}}$$

9.
$$\frac{\frac{a}{b} + \frac{c}{d}}{b + \frac{c}{d}}$$

13.
$$\frac{x+5}{1 + \frac{6}{x}} +$$

180. In the case of fractions like the following, called **Continued Fractions**, we begin from the lowest fraction and simplify step by step.

Example. Simplify $\frac{9x^2 - 64}{x - 1 - \frac{1}{1 - \frac{x}{4+x}}}$.

$$\begin{aligned} \text{The expression} &= \frac{9x^2 - 64}{x - 1 - \frac{1}{\frac{4+x-x}{4+x}}} = \frac{9x^2 - 64}{x - 1 - \frac{4+x}{4}} \\ &= \frac{9x^2 - 64}{\frac{4x - 4 - (4+x)}{4}} = \frac{9x^2 - 64}{\frac{3x - 8}{4}} \\ &= \frac{4(9x^2 - 64)}{3x - 8} = 4(3x + 8). \end{aligned}$$

EXAMPLES XXII. a.

Find the value of

1. $\frac{\frac{m}{n} - \frac{l}{m}}{\frac{a}{m} - \frac{b}{n}}$

2. $\frac{\frac{1}{x} + \frac{1}{y}}{\frac{1}{x} - \frac{1}{y}}$

3. $\frac{a + \frac{b}{d}}{x - \frac{y}{d}}$

4. $\frac{1 + \frac{c}{x}}{\frac{b}{x} - 1}$

5. $\frac{2 + \frac{3a}{4b}}{a + \frac{8b}{3}}$

6. $\frac{3a + \frac{7b}{8c}}{3c + \frac{7b}{8a}}$

7. $\frac{1 - \frac{y^2}{x^2}}{1 + \frac{y^2}{x^2}}$

8. $\frac{1}{a + \frac{b}{c}}$

9. $\frac{a}{x + \frac{c}{d}}$

10. $\frac{x}{x - \frac{m}{n}}$

11. $\frac{a + \frac{c}{d}}{\frac{m}{n} + \frac{k}{p}}$

12. $\frac{x - \frac{1}{x}}{1 + \frac{1}{x}}$

13. $\frac{x + 5 + \frac{6}{x}}{1 + \frac{6}{x} + \frac{8}{x^2}}$

14. $\frac{\frac{1}{x} - \frac{2}{x^2} - \frac{3}{x^3}}{\frac{9}{x} - x}$

15. $\frac{2x^2 - x - 6}{\frac{4}{x^3} - 1}$

Find the value of

$$16. \frac{2}{1-x^2} \div \left(\frac{1}{1-x} - \frac{1}{1+x} \right). \quad 17. \left(\frac{a^3-b^3}{a-b} - \frac{a^3+b^3}{a+b} \right) \div \frac{4ab}{a^2-b^2}.$$

$$18. \left(\frac{a^3-ax+x^3}{a-x} + \frac{a^3+ax+x^3}{a+x} \right) \div \frac{x^3}{a^3-x^3}.$$

$$19. \left(y + \frac{xy}{y-x} \right) \left(y - \frac{xy}{x+y} \right) \times \frac{y^3-x^3}{y^3+x^3}.$$

$$20. \left(\frac{x}{1+x} + \frac{1-x}{x} \right) \div \left(\frac{x}{1+x} - \frac{1-x}{x} \right).$$

$$21. \frac{\frac{a+b}{a-b} - \frac{a-b}{a+b}}{1 - \frac{a^2+b^2}{(a+b)^2}}.$$

$$22. \frac{\frac{a}{x^2} + \frac{x}{a^2}}{\frac{1}{a^2} - \frac{1}{ax} + \frac{1}{x^2}}.$$

$$23. \frac{\frac{1}{3x-2} - \frac{1}{3x+2}}{\frac{1}{9 - \frac{4}{x^2}}}.$$

$$24. 1 + \frac{x}{1+x + \frac{2x^2}{1-x}}.$$

$$25. \frac{\frac{1}{a^2-1}}{a - \frac{1}{a + \frac{1}{a-1}}}.$$

$$26. \frac{1}{4x + \frac{4x}{1 + \frac{2(x+y)}{6-x}}}.$$

$$27. \frac{\frac{a}{x + \frac{m}{y + \frac{n}{z}}}}{y + \frac{n}{z}}.$$

$$28. \frac{1}{1 - \frac{1+x}{x - \frac{1}{x}}}.$$

$$29. \frac{x-2}{x-2 - \frac{x}{x - \frac{x-1}{x-2}}}.$$

$$30. \frac{1}{x - \frac{1}{x + \frac{1}{x}}} - \frac{1}{x + \frac{1}{x - \frac{1}{x}}}.$$

$$31. \frac{\frac{a-b}{1+ab} + \frac{b-c}{1+bc}}{1 - \frac{(a-b)(b-c)}{(1+ab)(1+bc)}}.$$

$$32. \frac{1 + \frac{a-b}{a+b}}{1 - \frac{a-b}{a+b}} \div \frac{1 + \frac{a^2-b^2}{a^2+b^2}}{1 - \frac{a^2-b^2}{a^2+b^2}}.$$

$$33. \frac{\quad}{a^2 - a}$$

$$35. \frac{\quad}{4x - 2}$$

181. So
as a group

Example

182. Si
by the den
valent form

Example

Example

In some

Example

By divis

Thus the

Therefore

$$33. \frac{a-x}{a^2-ax-\frac{(a-x)^2}{1-\frac{a}{x}}}.$$

$$34. \frac{2x^2+2x}{\frac{x}{x-2}-1} - \frac{3(x-1)}{\frac{x}{x-4}-1} + \frac{3}{4}.$$

$$35. \frac{2-4x}{4x-2-\frac{4x}{1+\frac{2x-1}{1+\frac{1}{4x-1}}}}.$$

$$36. \frac{x^2}{1-\frac{1}{x^2+\frac{1}{x}}} + \frac{x^2-2}{1-\frac{1}{x^2-\frac{1}{x}}}.$$

181. Sometimes it is convenient to express a single fraction as a group of fractions.

$$\begin{aligned} \text{Example. } \frac{5x^2y-10xy^2+15y^3}{10x^2y^2} &= \frac{5x^2y}{10x^2y^2} - \frac{10xy^2}{10x^2y^2} + \frac{15y^3}{10x^2y^2} \\ &= \frac{1}{2y} - \frac{1}{x} + \frac{3y}{2x^2}. \end{aligned}$$

182. Since a fraction represents the quotient of the numerator by the denominator, we may often express a fraction in an equivalent form, partly integral and partly fractional.

$$\text{Example 1. } \frac{x+7}{x+2} = \frac{(x+2)+5}{x+2} = 1 + \frac{5}{x+2}.$$

$$\text{Example 2. } \frac{3x-2}{x+5} = \frac{3(x+5)-15-2}{x+5} = \frac{3(x+5)-17}{x+5} = 3 - \frac{17}{x+5}.$$

In some cases actual division may be advisable.

$$\text{Example 3. Shew that } \frac{2x^2-7x-1}{x-3} = 2x-1 - \frac{4}{x-3}.$$

By division,

$$\begin{array}{r} x-3 \overline{) 2x^2-7x-1} \\ \underline{2x^2-6x} \\ -x-1 \\ \underline{-x+3} \\ -4 \end{array}$$

Thus the quotient is $2x-1$, and the remainder -4 .

Therefore

$$\frac{2x^2-7x-1}{x-3} = 2x-1 - \frac{4}{x-3}.$$

183. If the numerator be of lower dimensions than the denominator, we may still perform the division, and express the result in a form which is partly integral and partly fractional.

Example. Prove that $\frac{2x}{1+3x^2} = 2x - 6x^3 + 18x^5 - \frac{54x^7}{1+3x^2}$.

$$\begin{array}{r} \text{By division} \quad 1+3x^2 \overline{) 2x} \quad (2x - 6x^3 + 18x^5 \\ \underline{2x + 6x^3} \\ -6x^3 - 18x^5 \\ \underline{-6x^3 - 18x^5} \\ 18x^5 \\ \underline{18x^5 + 54x^7} \\ -54x^7 \end{array}$$

whence the result follows.

Here the division may be carried on to any number of terms in the quotient, and we can stop at any term we please by taking for our remainder the fraction whose numerator is the remainder last found, and whose denominator is the divisor.

Thus, if we carried on the quotient to four terms, we should have

$$\frac{2x}{1+3x^2} = 2x - 6x^3 + 18x^5 - 54x^7 + \frac{162x^9}{1+3x^2}.$$

The terms in the quotient may be fractional; thus if x^2 is divided by $x^3 - a^3$, the first four terms of the quotient are $\frac{1}{x} + \frac{a^3}{x^4} + \frac{a^6}{x^7} + \frac{a^9}{x^{10}}$ and the remainder is $\frac{a^{12}}{x^{10}}$.

184. Miscellaneous examples in multiplication and division occur which can be dealt with by the preceding rules for the reduction of fractions.

Example. Multiply $x + 2a - \frac{a^2}{2x+3a}$ by $2x - a - \frac{2a^2}{x+a}$.

$$\begin{aligned} \text{The product} &= \left(x + 2a - \frac{a^2}{2x+3a} \right) \times \left(2x - a - \frac{2a^2}{x+a} \right) \\ &= \frac{2x^2 + 7ax + 6a^2 - a^2}{2x+3a} \times \frac{2x^2 + ax - a^2 - 2a^2}{x+a} \\ &= \frac{2x^2 + 7ax + 5a^2}{2x+3a} \times \frac{2x^2 + ax - 3a^2}{x+a} \\ &= \frac{(2x+5a)(x+a)}{2x+3a} \times \frac{(2x+3a)(x-a)}{x+a} \\ &= (2x+5a)(x-a). \end{aligned}$$

Express
fractions

1. $\frac{3x^2y}{}$

3. $\frac{a^3 - }{}$

5. $\frac{bc + }{}$

Perform
four terms

7. $x \div ()$

10. $1 \div ()$

13. Shev

14. Shev

15. Shev

16. Shev

17. Divi

18. Mul

19. Divi

20. Divi

21. Mul

EXAMPLES XXII. b.

Express each of the following fractions as a group of simple fractions in lowest terms:

1. $\frac{3x^2y + xy^2 - y^3}{9xy}$.

2. $\frac{3a^3x - 4a^2x^2 + 6ax^3}{12ax}$.

3. $\frac{a^3 - 3a^2b + 3ab^2 + b^3}{2ab}$.

4. $\frac{a + b + c}{abc}$.

5. $\frac{bc + ca + ab}{abc}$.

6. $\frac{a^3bc - 3ab^3c + 2abc}{6abc}$.

Perform the following divisions, giving the remainder after four terms in the quotient:

7. $x \div (1 + x)$.

8. $a \div (a - b)$.

9. $(1 + x) \div (1 - x)$.

10. $1 \div (1 - x + x^2)$.

11. $x^2 \div (x + 3)$.

12. $1 \div (1 - x)^2$.

13. Shew that $\frac{a^3 - b^3}{(a - b)^2} = a + 2b + \frac{3b^2}{a - b}$.

14. Shew that $x^2 - xy + y^2 - \frac{2y^3}{x + y} = \frac{x^3 - y^3}{x + y}$.

15. Shew that $\frac{60x^3 - 17x^2 - 4x + 1}{5x^2 + 9x - 2} = 12x - 25 + \frac{49}{x + 2}$.

16. Shew that $1 + \frac{a^2 + b^2 - c^2}{2ab} = \frac{(a + b + c)(a + b - c)}{2ab}$.

17. Divide $x + \frac{16x - 27}{x^2 - 16}$ by $x - 1 + \frac{13}{x + 4}$.

18. Multiply $a^2 - 2ax + 4x^2 - \frac{16x^3}{a + 2x}$ by $3 - \frac{6x(a + 4x)}{a^2 + 2ax + 4x^2}$.

19. Divide $b^3 + 3b - 2 - \frac{12}{b - 3}$ by $3b + 6 - \frac{2b^2}{b - 3}$.

20. Divide $a^2 + 9b^2 + \frac{65b^4}{a^2 - 9b^2}$ by $a + 3b + \frac{13b^2}{a - 3b}$.

21. Multiply $4x^2 + 14x + \frac{98x - 27}{2x - 7}$ by $\frac{1}{6} - \frac{3x + 29}{12x^2 + 18x + 27}$.

185. The following exercise contains miscellaneous examples which illustrate most of the processes connected with fractions.

EXAMPLES XXII. c.

Simplify the following fractions :

1. $\frac{4a(a^2-x^2)}{3b(c^2-x^2)} \div \left[\frac{a^2-ax}{bc+bx} \times \frac{a^2+2ax+x^2}{c^2-2cx+x^2} \right]$.
2. $\frac{x(x+a)(x+2a)}{3a} - \frac{x(x+a)(2x+a)}{6a}$.
3. $\frac{1}{b} \left(\frac{1}{a-b} - \frac{1}{a+2b} \right) - \frac{2}{a^2+ab-2b^2}$.
4. $\left(\frac{x+y}{x-y} \right)^2 - \left(\frac{x-y}{x+y} \right)^2$.
5. $\frac{2}{x-1} + \frac{2}{x+1} - \frac{4x}{x^2-x+1}$.
6. $\left(\frac{x^2}{1-x^4} + \frac{2x^4}{1-x^2} \right) \div \left(\frac{x^2+1}{x} \right)^2$.
7. $\frac{1}{x} - \frac{1}{(x+1)^2} - \frac{2}{x+1} + \frac{x}{1+x+x^2}$.
8. $\frac{1+x^3}{1+2x+2x^2+x^3}$.
9. $\frac{2x^3-9x^2+27}{3x^3-81x+162}$.
10. $\frac{a}{b} - \frac{(a^2-b^2)x}{b^2} + \frac{a(a^2-b^2)x^2}{b^2(b+ax)}$.
11. $\left\{ \frac{x^4-a^4}{x^3-2ax+a^3} \div \frac{x^2+ax}{x-a} \right\} \times \frac{x^5-a^2x^3}{x^3+a^3} \div \left(\frac{x-a}{a} \right)$.
12. $\frac{a^2-x^2}{a^3+ax+x^3} \div \frac{\left(1-\frac{x}{a}\right)^3 \left(1+\frac{x}{a}\right)^3}{a^3-x^3}$.
13. $\frac{x^4-2x^2+1}{3x^5-10x^3+15x-8}$.
14. $\frac{a^3+a(1+a)y+y^2}{a^4-y^2}$.
15. $\frac{1}{a} + \frac{2}{a+1} + \frac{3}{a+2} - \frac{\frac{4}{a}}{1+\frac{1}{a}}$.
16. $\frac{x+3}{2x^2+9x+9} + \frac{1}{2} \cdot \frac{1}{2x-3} - \frac{1}{x-\frac{9}{4x}}$.

$$17. \frac{x^2}{x^3+x^2}$$

$$18. \frac{1}{(1+ax)}$$

$$19. \frac{2x^3-x}{x^3-x^2}$$

$$21. \frac{2a}{(x-2a)}$$

$$22. \frac{1}{2} \left(\frac{a^2}{a^2-x^2} \right)$$

$$23. \frac{x}{2} \left(\frac{1}{x-1} \right)$$

$$24. \frac{1}{x+y} \div$$

$$25. (3x-5)$$

$$26. \left\{ \frac{2}{x} - \frac{1}{a} \right\}$$

$$27. \frac{1}{2x-1}$$

$$29. \left\{ \frac{b+\frac{c}{1}}{1-\frac{(a)}{1}} \right\}$$

$$30. \frac{\frac{x^2+y^2}{y}}{\frac{1}{y} - \frac{1}{x}}$$

$$32. \frac{a+\frac{b-1}{1}}{1-\frac{a(b-1)}{1}}$$

examples
fractions.

$$17. \frac{2}{x^3+x^2+x+1} - \frac{2}{x^3-x^2+x-1}.$$

$$18. \frac{1-a^2}{(1+ax)^2-(a+x)^2} \div \frac{1}{2} \left(\frac{1}{1-x} + \frac{1}{1+x} \right).$$

$$19. \frac{2x^3-x^2-2x+1}{x^3-3x+2}.$$

$$20. \frac{x^2-6x+8}{4x^3-21x^2+15x+20}.$$

$$21. \frac{2a}{(x-2a)^2} - \frac{x-a}{x^3-5ax+6a^2} + \frac{2}{x-3a}.$$

$$22. \frac{1}{2} \left(\frac{a^2+x^2}{a^2-x^2} \right) - \frac{1}{2} \cdot \frac{a+x}{a-x} - \left(\frac{a}{a+x} \right)^2.$$

$$23. \frac{x}{2} \left(\frac{1}{x-y} - \frac{1}{x+y} \right) \times \frac{x^2-y^2}{x^2y+xy^2} \div \frac{1}{x+y}.$$

$$24. \frac{1}{x+y} \div \left[\frac{y}{2} \left(\frac{1}{x+y} + \frac{1}{x-y} \right) \times \frac{x^2-y^2}{x^2y+xy^2} \right].$$

$$25. \left(3x-5-\frac{2}{x} \right) \left(3x+5-\frac{2}{x} \right) \div \left(x-\frac{4}{x} \right).$$

$$26. \left\{ \frac{2}{x} - \frac{1}{a+x} + \frac{1}{a-x} \right\} \div \left(\frac{a+x}{a-x} - \frac{a-x}{a+x} \right).$$

$$27. \frac{1}{2x-1} - \frac{2x-\frac{1}{2x}}{4x^2-1}.$$

$$28. \left(b + \frac{ab}{b-a} \right) \left(b - \frac{ab}{a+b} \right) \left(\frac{b^2-a^2}{b^2+a^2} \right).$$

$$29. \left\{ \frac{b + \frac{a-b}{1+ab}}{1 - \frac{(a-b)b}{1+ab}} - \frac{a - \frac{a-b}{1-ab}}{1 - \frac{a(a-b)}{1-ab}} \right\} \div \left(\frac{a}{b} - \frac{b}{a} \right).$$

$$30. \frac{\frac{x^2+y^2}{y} - x}{\frac{1}{y} - \frac{1}{x}} \times \frac{x^2-y^2}{x^2+y^2}.$$

$$31. \frac{2 \left(x^2 - \frac{1}{4} \right)}{2x+1} + \frac{1}{2}.$$

$$32. \frac{a + \frac{b-a}{1+ab}}{1 - \frac{a(b-a)}{1+ab}} \times \frac{\frac{x+y}{1-xy} - y}{1 + \frac{y(x+y)}{1-xy}}.$$

$$33. \frac{\frac{a+b}{a-b} + \frac{a-b}{a+b}}{\frac{a-b}{a+b} - \frac{a+b}{a-b}} \times \frac{ab^2-a^2b}{a^2+b^2}.$$

Simplify the following fractions :

$$34. \frac{(1-x^2)(1-x^2)}{x(1+x)(1-x)^2} - \frac{x^3 + \frac{1}{x^3}}{x^2 + \frac{1}{x^2} - 1}.$$

$$35. \left\{ x^3 - \frac{1}{x^3} - 3\left(x - \frac{1}{x}\right) \right\} \div \left(x - \frac{1}{x}\right).$$

$$36. \frac{1 + \frac{1}{m}}{\frac{1}{m}} \times \frac{\frac{1}{m}}{m^2 + \frac{1}{m}} \div \frac{\frac{1}{m}}{m - 1 + \frac{1}{m}}.$$

$$37. \frac{\frac{x}{y} + \frac{y}{x} - 1}{\frac{x^2}{y^2} + \frac{y^2}{x^2} + 1} \times \frac{1 + \frac{y}{x}}{x - y} \div \frac{1 + \frac{y^3}{x^3}}{\frac{x^2}{y} - \frac{y^2}{x}}.$$

$$38. \frac{\left(\frac{3x+x^3}{1+3x^2}\right)^2 - 1}{\frac{3x^3-1}{x^3-3x} + 1} \div \frac{\frac{9}{x^2} - \frac{33-x^2}{2(x^2+3)}}{\frac{2(x^2+3)}{(x^3-x)^2}}.$$

$$39. \frac{1}{a^2-2} - \frac{2}{a^2-1} + \frac{2}{a^2+1} - \frac{1}{a^2+2}.$$

$$40. \frac{1}{6m-2n} + \frac{1}{3m+2n} - \frac{3}{6m+2n}.$$

$$41. \frac{3}{4(1-x)^2} + \frac{3}{3(1-x)} + \frac{1}{8(1+x)} + \frac{x-1}{4(1+x^2)}.$$

$$42. \frac{4}{9(x-2)} + \frac{5}{9(x+1)} - \frac{1}{3(x+1)^2} - \frac{1}{x+2 + \frac{1}{x}}.$$

$$43. \left(\frac{x^2}{y} + \frac{y^2}{x}\right)\left(\frac{1}{y^2-x^2}\right) - \frac{y}{x^2+xy} + \frac{x}{xy-y^2}.$$

$$44. \frac{x^2-(y-z)^2}{(x+z)^2-y^2} + \frac{y^2-(z-x)^2}{(x+y)^2-z^2} + \frac{z^2-(x-y)^2}{(y+z)^2-x^2}.$$

$$45. \frac{x^2-(y-2z)^2}{(2z+x)^2-y^2} + \frac{y^2-(2z-x)^2}{(x+y)^2-4z^2} + \frac{4z^2-(x-y)^2}{(y+2z)^2-x^2}.$$

$$46. \frac{(x-y)(y-z) + (y-z)(z-x) + (z-x)(x-y)}{x(z-x) + y(x-y) + z(y-z)}.$$

$$47. \frac{a-b}{(a-b)}$$

$$48. \frac{c}{(a-b)}$$

$$49. \frac{x^2-(2z+x)}{(3z+x)}$$

$$50. \frac{9y^2-(2x+3)}{(2x+3)}$$

$$51. \frac{\frac{1}{x} - \frac{x}{x^2}}{\frac{1}{a} - \frac{a}{a^2}}$$

$$52. \frac{(x+a)}{a^2-a}$$

$$53. \left(\frac{a}{a^2-a}\right)$$

$$54. \frac{\frac{x-1}{3} + \frac{x+2}{4}}{\frac{x+2}{4} + \frac{x-1}{3}}$$

$$55. \frac{x-1}{3} + \frac{x+2}{4}$$

$$56. (x+2)$$

$$57. (1+a)^2$$

$$58. \frac{\{ax^2 + \dots\}}{\{ax^2 + \dots\}}$$

$$47. \frac{a-b-c}{(a-b)(a-c)} + \frac{b-c-a}{(b-c)(b-a)} + \frac{c-a-b}{(c-a)(c-b)}$$

$$48. \frac{c+a}{(a-b)(a-c)} + \frac{a+b}{(b-c)(b-a)} + \frac{b+c}{(c-a)(c-b)}$$

$$49. \frac{x^2 - (2y-3z)^2}{(3z+x)^2 - 4y^2} + \frac{4y^2 - (3z-x)^2}{(x+2y)^2 - 9z^2} + \frac{9z^2 - (x-2y)^2}{(2y+3z)^2 - x^2}$$

$$50. \frac{9y^2 - (4z-2x)^2}{(2x+3y)^2 - 16z^2} + \frac{16z^2 - (2x-3y)^2}{(3y+4z)^2 - 4x^2} + \frac{4x^2 - (3y-4z)^2}{(4z+2x)^2 - 9y^2}$$

$$51. \frac{\frac{1}{x} - \frac{x+a}{x^2+a^2}}{\frac{1}{a} - \frac{a+x}{a^2+x^2}} + \frac{\frac{1}{x} - \frac{x-a}{x^2+a^2}}{\frac{1}{a} - \frac{a-x}{a^2+x^2}}$$

$$52. \frac{(x+a)(x+b) - (y+a)(y+b)}{x-y} - \frac{(x-a)(y-b) - (x-b)(y-a)}{a-b}$$

$$53. \left(\frac{a+x}{a^2-ax+x^2} - \frac{a-x}{a^2+ax+x^2} \right) \div \left(\frac{a^2+x^2}{a^3-x^3} - \frac{a^2-x^2}{a^3+x^3} \right)$$

$$54. \frac{\frac{x-1}{3} + \frac{x-1}{x-2}}{\frac{x+2}{4} + \frac{x+2}{x-3}} \div \frac{\frac{x+3}{7} - \frac{x+3}{x+4}}{\frac{x-2}{3} + \frac{x-2}{x-1}} \quad 55. \frac{x-2 + \frac{6}{x+3}}{x-4 + \frac{12}{x+3}} \times \frac{\frac{x+3}{4} - \frac{x+3}{x+1}}{\frac{x-3}{7} + \frac{x-3}{x-4}}$$

$$56. (x+2) \left\{ 1 + \frac{6(x+2)}{x^2-x-6} \right\} \left\{ 1 - \frac{5(x+1)}{x^2+3x+2} \right\}$$

$$57. (1+a)^2 \div \left\{ 1 + \frac{a}{1-a + \frac{a}{1+a+a^2}} \right\}$$

$$58. \frac{\{ax^2 + (b-c)x - f\}^2 - \{ax^2 + (b+c)x - f\}^2}{\{ax^2 + (b+e)x - f\}^2 - \{ax^2 + (b-e)x - f\}^2}$$

MISCELLANEOUS EXAMPLES IV.

[The following Examples for revision are arranged in groups under different headings; each group illustrates one or more of the principal rules and processes already discussed, and for the most part the Examples present more variety and difficulty than those of the same type which have appeared in previous exercises.]

Substitutions and Brackets.

- Find the value of $\frac{a + \sqrt{a^2 + b^2}}{a^3 - 2b(a^2 - b^2)}$ when $a = -4$, $b = -3$.
- When $a = 1$, $b = -1$, $c = 2$ evaluate the expression

$$\sqrt{3a^3(b-c) + 3b^3(c-a) + 3c^3(a-b)}.$$
- Simplify

$$a(b-c)^3 - a(b-c)(2b^2 - bc + 2c^2) + (ab+ac)(b^2 - c^2);$$
 and find its value when $a = 1$, $b = 2$, $c = 3$.
- Find the value of

$$\sqrt[3]{5(b^2 - c^2) - a^2} + \sqrt[4]{3\{a(a^2 - c^2) - 1\}}$$
 when $a = 4$, $b = 5$, $c = 3$.
- Find the value of

$$\sqrt{(x^2 + y^2 + z)(x - y - 3z)} \div \sqrt[3]{xy^2z^2}$$
 when $x = -1$, $y = -3$, $z = 1$.
- When $a = 0$, $b = 2$, $x = 1$, $y = -3$, $z = 5$, find the numerical value of
 - $(x-y)^3 - 3a(x-y)^2 + 3b(x^3 - y^3);$
 - $(x-a)^3 - b^2(x-y+z) + \sqrt{(bx^2 - axy + y^2 + z^2)}.$
- If $x = 6$, $y = 7$, $z = 8$, find the value of
 - $x\left\{\frac{x}{7} - \frac{2}{3}\left(\frac{y}{4} + 1\right)\right\} - \frac{2}{7}\left\{y(x+1) + \frac{1}{2}(x^2 - 2y) - \frac{7x}{3}\right\};$
 - $x\left\{y\left(z - \frac{1}{z}\right)\right\} - \frac{y}{z}\left\{y - 3\left(\frac{x}{3} - xz^2\right)\right\} - 2xz\left\{-y\left(1 + \frac{y}{2xz^2}\right)\right\}.$

8. Evaluate $\frac{2a-b\{c-a(b-c)\}}{a^2(b+c)+b^2(c-a)} \cdot \sqrt{\frac{a+bc}{b^2+4ac}},$

when $a=2$, $b=-1$, $c=1$.

Find the value of

$$\frac{a-[b-\{a-(b-c)\}]-\sqrt{2a^2+b^2+c^2}}{\sqrt{c^4+a^2b^2+2a^3c}},$$

when $a=-1$, $b=5$, $c=3$.

10. When $a=4$, $b=-2$, $c=\frac{3}{2}$, $d=-1$, find the value of

$$(1) \ a^2-b^2-(a-b)^2-11(3b+2c)\left(2c^2-\frac{d^2}{2}\right);$$

$$(2) \ \sqrt[3]{4c^2-a(a-2b-d)}-\sqrt[2]{b^4c+11b^3d^2}.$$

11. Find the value of

$$l-m-3\left[\left\{l-m+lm\left(\frac{1}{l}+\frac{1}{m}\right)\right\}^2-4l^2\right]$$

when $l=\frac{2}{3}$, $m=-\frac{1}{3}$.

12. When $a=1$, $b=-\frac{1}{2}$, $c=0$, evaluate

$$\frac{a-[b-c-\{2a-2b-\frac{1}{2}(3c-b)\}]}{a-\frac{b}{c-\frac{1}{a}}}.$$

13. Simplify

$$42\left\{\frac{4x-3y}{6}-\frac{3}{7}\left(x-\frac{4}{3}y\right)\right\}-56\left\{\frac{1}{7}(3x-2y)-\frac{3}{8}\left(\frac{16}{3}x-y\right)\right\};$$

and find its value when $x=\frac{1}{6}$, $y=-\frac{3}{2}$.

14. If $x=6$, $y=7$, $z=8$ find the value of

$$(1) \ \frac{z}{x}\left\{y-\left(2z-\frac{x}{z}\right)\right\}-\frac{y}{z}\left[\frac{z^2}{x}-3\left\{1+\frac{2x^3}{3xy}\right\}+\frac{x}{2}\right];$$

$$(2) \ 3y\left\{\frac{2yz}{3}-\left(\frac{y}{5}-1\right)\right\}+\frac{y^2}{5}\left[3-z\left\{10-\frac{x}{2}(y-7)\right\}\right].$$

Resolution into Factors.

(On Arts. 128-132.)

Resolve into two or more factors :

- | | | |
|-----------------------------|------------------------------|-------------------------------|
| 15. $x^3 + 21x + 108.$ | 16. $a^3 + 6a - 91.$ | 17. $x^3 - 20xy + 96y^2.$ |
| 18. $a^2b^3 - 14ab - 51.$ | 19. $c^3 + c^2 - 153c.$ | 20. $m^2n - 6mn^2 + 9n^3.$ |
| 21. $p^4 - p^2q^2 - 56q^4.$ | 22. $d^4 - 4d^2c^2 - 45c^4.$ | 23. $x^3y - x^2y^2 - 42xy^3.$ |
| 24. $m^3 + 28m + 195.$ | 25. $210 - a - a^2.$ | 26. $57 + 16pq - p^2q^2.$ |
| 27. $x^4 + 27x^2 + 176.$ | 28. $a^4 + 7a^2 - 98.$ | 29. $c^3 + 54c + 729.$ |
| 30. $72 + xy - x^2y^2.$ | 31. $a^4 + 9a^2x^2 + 14x^4.$ | 32. $p^3 - 3pq - 108q^2.$ |
| 33. $2a^5 + 2a^3 - 264.$ | 34. $x^4 - 2x^2 - 63x^2.$ | 35. $b^2c^2 + 5bc - 84.$ |
| 36. $z^3 + 34z + 289.$ | 37. $a^3 - 22ac + 57c^2.$ | 38. $y^3z + 6y^2z - 91yz.$ |
| 39. $2 + x^3 - 3x^4.$ | 40. $2a^2b^3 + ab - 15.$ | 41. $9p^3 - 24p + 16.$ |
| 42. $35 + 12mn + m^2n^2.$ | 43. $119 - 10c - c^2.$ | 44. $6x^3 - 5x^4 + x^5.$ |
| 45. $6m^2 + 7m - 3.$ | 46. $4a^2 - 8ab - 5b^2.$ | 47. $6p^2 - 13pq + 2q^2.$ |
| 48. $20x^2 - 9xz - 20z^2.$ | 49. $8x^4 + 2x^2 - 15.$ | 50. $12y^2 - 30y + 12.$ |
| 51. $12(a^2b^2 - 1) + 7ab.$ | 52. $2(a^4b^2 + 5) - 9a^2b.$ | |
| 53. $21x^2 + 2y(5x - 8y).$ | 54. $3(6m^2 - 5n^2) + 17mn.$ | |

(On Arts. 133-137.)

Resolve into two or more factors :

- | | |
|---|---|
| 55. $c^3 - a^2 - b^2 + 2ab.$ | 56. $a^3 + 2bc - b^3 - c^2.$ |
| 57. $125x^3 + 27y^3.$ | 58. $a^3b^3 + 343.$ |
| 59. $512b^3 - a^6.$ | |
| 60. $a^2 - 4(x - y)^2.$ | 61. $2mn + m^3 - 1 + n^2.$ |
| 62. $8c^4 - 2c^2(d + c)^2.$ | 63. $(a^2b^2 - 1)^2 - x^2 + 2xy - y^2.$ |
| 64. $1 - 64m^6.$ | 65. $p^3 + 1000p^2q^3.$ |
| 66. $6561 - a^4.$ | |
| 67. $x^4 - 2x^2 - y^2 - z^2 + 2yz + 1.$ | 68. $a^2 - 16(b - c)^2.$ |
| 69. $c - d - 4(c - d)^3.$ | 70. $p^3 - 16q^2 + p - 4q.$ |
| 71. $2 + 128(a + b)^3.$ | 72. $x + 3y + x^3 + 27y^3.$ |

Resolve

73. $x^3 + x$ 75. $14 - 5$ 78. $1 - (m$ 80. $9b^2 -$ 82. $3x^2 -$ 84. $a(b^2 +$

86. Expr

87. Find

Highest

88. Find t

13ab

89. Find t

90. Find t

each

91. Find t

 $a(a +$

92. Find t

(

93. Find t

of it

(Miscellaneous Factors.)

Resolve into two or more factors :

73. $a^2 + x^2y + xy^2 + y^3$. 74. $acx^2 + \dots - adx - bd$.
 75. $14 - 5a - a^2$. 76. $98x^4 - 7x^2y^2 - y^4$. 77. $51 - 14a - a^2$.
 78. $1 - (m^2 + p^2) - 2mp$. 79. $ab(x^2 + 1) - x(a^2 + b^2)$.
 80. $9b^2 - 6bc - 16 + c^2$. 81. $x^2c^3 - c^3 + x^2 - 1$.
 82. $3x^3 - 2ab - x(b - 6a)$. 83. $m^3 - n^3 - (x^2 - mn)(m - n)$.
 84. $a(b^2 + c^2 - a^2) + b(a^2 + c^2 - b^2)$. 85. $x^7 - x^3 + 8x^4 - 8$.
 86. Express in factors the square root of
 $(x^3 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)$.
 87. Find the expression whose square is
 $(2x^3 - xy - 15y^2)(4x^2 - 25y^2) - 11xy + 15y^2$.


Highest Common Factor and Lowest Common Multiple.

88. Find the lowest common multiple of
 $13ab^2(x^3 - 3a^2x + 2a^3)$, $65a^3b(x^2 + ax - 2a^2)$, $25b^3(x^2 - a^2)^2$.
 89. Find the highest common factor of
 $2(x^4 + 9) - 5x^2(x + 1)$, $2x^3(2x - 9) + 81(x - 1)$.
 90. Find the expression of lowest dimensions which is divisible by
 each of the following expressions :
 $(2x^4 + 4x^3)(x^2 + 2x - 8)$, $(2x^3 - 4x^2)(x^3 - 2x - 8)$,
 $(x^2 - 4x)(x^2 + 2x - 8)$.
 91. Find the H.C.F. and the L.C.M. of the three expressions
 $a(a + c) - b(b + c)$, $b(b + a) - c(c + a)$, $c(c + b) - a(a + b)$.
 92. Find the divisor of highest dimensions of the expressions
 $(a + b)(a - b) + c(c - 2a)$, $(a + c)(a - c) + b(b + 2a)$.
 93. Find the expression of lowest dimensions such that the L.C.M.
 of it and $2a^3 - 3ab + b^3$ is
 $2a^4 - 3a^3b - a^2b^2 + 3ab^3 - b^4$.



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

94. Shew that
- $x^2 - 4y^2$
- is the H.C.F. of the expressions

$$x^4 - 3x^2y^2 - 4y^4, \quad x^6 - 64y^6,$$

$$\text{and} \quad x^5 + 32y^5 - 8x^3y^2 + 2x^4y + 16xy^4 - 16x^2y^3.$$

95. Find the lowest common multiple of

$$(a-c)^2 - (b-c)^2, \quad a^2 + b^2 - 2ac - 2bc + 2ab, \quad a^4 - b^4.$$

96. Shew that the lowest common multiple of

$$a(a-b)^2 - ac^2, \quad a^2b - b(b-c)^2, \quad (a+c)^2c - b^2c$$

$$\text{is} \quad abc(a^4 + b^4 + c^4 - 2b^2c^2 - 2c^2a^2 - 2a^2b^2).$$

97. Prove that

$$x^4 - 15x^3 + 75x^2 - 145x + 84$$

and

$$x^4 - 17x^3 + 101x^2 - 247x + 210$$

have the same H.C.F. and L.C.M. as

$$x^4 - 13x^3 + 53x^2 - 83x + 42$$

and

$$x^4 - 19x^3 + 131x^2 - 389x + 420.$$

Simplification of Fractions.

Simplify

98.
$$\frac{1+x+x^2}{1-x^3} + \frac{x-x^2}{(1-x)^3}$$

99.
$$\frac{2x-7}{(x-3)^2} - \frac{2(x+2)}{x^2-9}$$

100.
$$\frac{1}{2x^2-\frac{1}{2}} + \frac{1}{(2x+1)^2}$$

101.
$$\frac{(x+1)^3 - (x-1)^3}{3x^3+x}$$

102.
$$\frac{1}{(1-x)^2} + \frac{2}{1-x^2} + \frac{1}{(1+x)^2}$$

103.
$$\frac{(x^3-2x)^2 - (x^2-2)^2}{(x-1)(x+1)(x^2-2)^2}$$

104.
$$\frac{1}{6x-2} - \frac{1}{2\left(x-\frac{1}{3}\right)} - \frac{1}{1-3x}$$

105.
$$\frac{x}{9} + \frac{2}{3} + \frac{4}{x-6} - \frac{2}{3} \cdot \frac{1}{1-\frac{6}{x}}$$

106.
$$\frac{x}{x^3-y^3} - \frac{1}{x-y} + \frac{1}{x+y} + \frac{1}{x} - \frac{1}{y} + \frac{x^2-xy+y^2}{xy(x-y)}$$

107.
$$\left(x-y-\frac{4y^2}{x-y}\right)\left(x+y-\frac{4x^2}{x+y}\right) \div \left\{3(x+y) - \frac{8xy}{x-y}\right\}.$$

108.
$$\frac{a^4+}{a^4+}$$

109.
$$\frac{(a+)}{(a+)}$$

110.
$$\left[\left(\frac{x}{a}\right)\right]$$

111.
$$\frac{x^4-}{(x^2+)}$$

112.
$$\frac{1+x}{1-x}$$

113.
$$\frac{1}{1}$$

114.
$$\frac{1}{x-}$$

115.
$$\frac{1}{1+}$$

116.
$$\frac{ab}{a+b}$$

117.
$$\frac{(a+b)}{(a+b)}$$

118.
$$\left\{\frac{1}{(a-)}\right\}$$

$$108. \frac{a^4 + b^4 + ab(a^2 + b^2)}{(a+b)^2} - \frac{a^4 + b^4 - ab(a^2 + b^2)}{(a-b)^2} + \frac{12a^2b^2}{(a+b)^2 - (a-b)^2}.$$

$$109. \frac{(ac + bd)^3 - (ad + bc)^3}{(a-b)(c-d)} - \frac{(ac + bd)^3 + (ad + bc)^3}{(a+b)(c+d)}.$$

$$110. \left[\left(\frac{x}{a} \right)^2 + \left(\frac{z-x}{b} \right)^2 \right] \div \left[\frac{z^2}{a^2 + b^2} + \frac{a^2 + b^2}{a^2 b^2} \left(x - \frac{za^2}{a^2 + b^2} \right)^2 \right].$$

$$111. \frac{x^4 - (x-1)^2}{(x^2+1)^2 - x^2} + \frac{x^2 - (x^2-1)^2}{x^2(x+1)^2 - 1} + \frac{x^2(x-1)^2 - 1}{x^4 - (x+1)^2}.$$

$$112. \frac{\frac{1+x}{1-x} + \frac{4x}{1+x^2} + \frac{8x}{1-x^2} - \frac{1-x}{1+x}}{\frac{1+x^2}{1-x^2} + \frac{4x^2}{1+x^4} - \frac{1-x^2}{1+x^2}}. \quad 113. \left\{ \frac{f}{g - \frac{g^2}{f}} + \frac{g}{f - \frac{f^2}{g}} \right\} \times \frac{1}{\frac{f^2}{g} - \frac{g^2}{f}}.$$

$$114. \frac{1}{x - \frac{2}{x + \frac{1}{2}}} \times \frac{1}{2 + \frac{1}{x}} \div \frac{x}{2x - \frac{x+4}{x+1}}.$$

$$115. \frac{\frac{x}{1 + \frac{x}{1-x + \frac{x}{1+x}}}}{1 + 3x + 3x^2 + 2x^3}.$$

$$116. \frac{ab}{a+b} \left(3c + \frac{b}{a} \right) - \frac{b^2}{(a+b)^3} (a^2 + b^2) - 2a \left(\frac{b}{a+b} \right)^3.$$

$$117. \frac{\frac{(a+b)^2 + (a-b)^2}{b-a} - (a+b)}{\frac{1}{b-a} - \frac{1}{a+b}} \div \frac{(a+b)^3 + (b-a)^3}{(a+b)^2 - (a-b)^2}.$$

$$118. \left\{ \frac{c-b}{(a-b)(a-c)} - \frac{c-a}{(b-c)(b-a)} + \frac{b-a}{(c-a)(c-b)} \right\} \div \frac{2(a^2 + b^2 + c^2 - bc - ca - ab)}{(a-b)(b-c)(c-a)}.$$

When
they sho

Exampl

By tra

CHAPTER XXIII.

HARDER EQUATIONS.

186. In this chapter we propose to give a miscellaneous collection of equations. Some of these will serve as a useful exercise for revision of the methods already explained in previous chapters; but we also add others presenting more difficulty, the solution of which will often be facilitated by some special artifice.

The following examples worked in full will sufficiently illustrate the most useful methods.

Example 1. Solve $\frac{6x-3}{2x+7} = \frac{3x-2}{x+5}$.

Multiplying up, we have

$$(6x-3)(x+5) = (3x-2)(2x+7),$$

$$6x^2 + 27x - 15 = 6x^2 + 17x - 14;$$

$$\therefore 10x = 1;$$

$$\therefore x = \frac{1}{10}.$$

Note. By a simple reduction many equations can be brought to the form in which the above equation is given. When this is the case, the necessary simplification is readily completed by multiplying up, or "multiplying across," as it is sometimes called.

Example 2. Solve $\frac{8x+23}{20} - \frac{5x+2}{3x+4} = \frac{2x+3}{5} - 1$.

Multiply by 20, and we have

$$8x + 23 - \frac{20(5x+2)}{3x+4} = 4x + 12 - 20.$$

By transposition, $31 = \frac{20(5x+2)}{3x+4}$.

Multiplying across, $93x + 124 = 20(5x+2),$

$$84 = 7x;$$

$$\therefore x = 12.$$

Multip

Exampl

This eq
work wou
by procee

Transpo

Simplify

$$(x-8)$$

$$\therefore \frac{x^2-1}{x-1}$$

Hence, s
equal;
that is,

When two or more fractions have the same denominator they should be taken together and simplified.

Example 3. Solve $\frac{13-2x}{x+3} + \frac{23x+8\frac{1}{3}}{4x+5} = \frac{16-\frac{1}{4}x}{x+3} + 4$.

By transposition, we have

$$\begin{aligned}\frac{23x+8\frac{1}{3}}{4x+5} - 4 &= \frac{16-\frac{1}{4}x-13+2x}{x+3} \\ \therefore \frac{7x-\frac{35}{3}}{4x+5} &= \frac{3+\frac{7x}{4}}{x+3}\end{aligned}$$

Multiplying across, we have

$$\begin{aligned}7x^2 - \frac{35x}{3} + 21x - 35 &= 12x + 7x^2 + 15 + \frac{35x}{4} \\ -\frac{137x}{12} &= 50; \\ \therefore x &= -\frac{600}{137}.\end{aligned}$$

Example 4. Solve $\frac{x-8}{x-10} + \frac{x-4}{x-6} = \frac{x-5}{x-7} + \frac{x-7}{x-9}$.

This equation might be solved by clearing of fractions, but the work would be very laborious. The solution will be much simplified by proceeding as follows:

Transposing, $\frac{x-8}{x-10} - \frac{x-5}{x-7} = \frac{x-7}{x-9} - \frac{x-4}{x-6}$

Simplifying each side *separately*, we have

$$\begin{aligned}\frac{(x-8)(x-7)-(x-5)(x-10)}{(x-10)(x-7)} &= \frac{(x-7)(x-6)-(x-4)(x-9)}{(x-9)(x-6)}; \\ \therefore \frac{x^2-15x+56-(x^2-15x+50)}{(x-10)(x-7)} &= \frac{x^2-13x+42-(x^2-13x+36)}{(x-9)(x-6)}; \\ \therefore \frac{6}{(x-10)(x-7)} &= \frac{6}{(x-9)(x-6)}.\end{aligned}$$

Hence, since the numerators are equal, the denominators must be equal;

that is,

$$\begin{aligned}(x-10)(x-7) &= (x-9)(x-6), \\ x^2-17x+70 &= x^2-15x+54; \\ \therefore 16 &= 2x; \\ \therefore x &= 8.\end{aligned}$$

The above equation may also be solved very neatly by the following artifice.

The equation may be written in the form

$$\frac{(x-10)+2}{x-10} + \frac{(x-6)+2}{x-6} = \frac{(x-7)+2}{x-7} + \frac{(x-9)+2}{x-9};$$

whence we have

$$1 + \frac{2}{x-10} + 1 + \frac{2}{x-6} = 1 + \frac{2}{x-7} + 1 + \frac{2}{x-9};$$

which gives

$$\frac{1}{x-10} + \frac{1}{x-6} = \frac{1}{x-7} + \frac{1}{x-9}.$$

Transposing,

$$\frac{1}{x-10} - \frac{1}{x-7} = \frac{1}{x-9} - \frac{1}{x-6};$$

$$\therefore \frac{3}{(x-10)(x-7)} = \frac{3}{(x-9)(x-6)},$$

and the solution may be completed as before.

Example 5. Solve $\frac{5x-64}{x-13} - \frac{2x-11}{x-6} = \frac{4x-55}{x-14} - \frac{x-6}{x-7}.$

We have $5 + \frac{1}{x-13} - \left(2 + \frac{1}{x-6}\right) = 4 + \frac{1}{x-14} - \left(1 + \frac{1}{x-7}\right);$

$$\therefore \frac{1}{x-13} - \frac{1}{x-6} = \frac{1}{x-14} - \frac{1}{x-7}.$$

The solution may now be completed as before, and we obtain $x=10.$

EXAMPLES XXIII. a.

1. $\frac{x+4}{3x-8} = \frac{x+5}{3x-7}.$

2. $\frac{3x+1}{3(x-2)} = \frac{x-2}{x-1}.$

3. $\frac{7-5x}{1+x} = \frac{11-15x}{1+3x}.$

4. $\frac{3(7+6x)}{2+9x} = \frac{35+4x}{9+2x}.$

5. $\frac{6x+13}{15} - \frac{3x+5}{5x-25} = \frac{2x}{5}.$

6. $\frac{6x+8}{2x+1} - \frac{2x+38}{x+12} = 1.$

7. $\frac{3x-1}{2x-1} - \frac{4x-2}{3x-1} = \frac{1}{6}.$

8. $\frac{x+25}{x-5} = \frac{2x+75}{2x-15}.$

9. $\frac{x}{x+2} + \frac{4}{x+6} = 1.$

10. $\frac{6x+7}{9x+6} = \frac{1}{12} + \frac{5x-5}{12x+8}.$

11. $\frac{2x}{x+1} - \frac{1}{x-2} = \frac{3x-1}{x^2-1}.$

12. $\frac{4}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

13. $\frac{2x}{x+1} - \frac{1}{x-2} = \frac{3}{x-3}.$

15. $\frac{4}{x+1} - \frac{1}{x-2} = \frac{3}{x-3}.$

16. $\frac{7}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

17. $\frac{3}{4-x} - \frac{1}{x-2} = \frac{3}{x-3}.$

18. $\frac{25}{x+1} - \frac{1}{x-2} = \frac{3}{x-3}.$

20. $\frac{x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

22. $\frac{x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

23. $\frac{x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

25. $\frac{4x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

26. $\frac{5x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

27. $\frac{2x}{3x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

29. $\frac{083}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

30. $\frac{2x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

31. $\frac{3x}{5x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

33. $\frac{3x}{x-1} - \frac{1}{x-2} = \frac{3}{x-3}.$

he follow-

$$11. \frac{2x-5}{5} + \frac{x-3}{2x-15} = \frac{4x-3}{10} - 1\frac{1}{10}.$$

$$12. \frac{4(x+3)}{9} = \frac{8x+37}{18} - \frac{7x-29}{5x-12}.$$

$$13. \frac{(2x-1)(3x+8)}{6x(x+4)} - 1 = 0.$$

$$14. \frac{2x+5}{5x+3} - \frac{2x+1}{5x+2} = 0.$$

$$15. \frac{4}{x+3} - \frac{2}{x+1} = \frac{5}{2x+6} - \frac{2\frac{1}{2}}{2x+2}.$$

$$16. \frac{7}{x-4} - \frac{60}{5x-30} = \frac{10\frac{1}{2}}{3x-12} - \frac{8}{x-6}.$$

$$17. \frac{3}{4-2x} + \frac{30}{8(1-x)} = \frac{3}{2-x} + \frac{5}{2-2x}.$$

$$18. \frac{25-\frac{x}{3}}{x+1} + \frac{16x+4\frac{1}{6}}{3x+2} = 5 + \frac{23}{x+1}.$$

$$19. \frac{30+6x}{x+1} + \frac{60+8x}{x+3} = 14 + \frac{48}{x+1}.$$

$$20. \frac{x}{x-2} - \frac{x+1}{x-1} = \frac{x-8}{x-6} - \frac{x-9}{x-7}.$$

$$21. \frac{x+5}{x+4} - \frac{x-6}{x-7} = \frac{x-4}{x-5} - \frac{x-15}{x-16}.$$

$$22. \frac{x-7}{x-9} - \frac{x-9}{x-11} = \frac{x-13}{x-15} - \frac{x-15}{x-17}.$$

$$23. \frac{x+3}{x+3} - \frac{x+6}{x+9} = \frac{x+2}{x+5} - \frac{x+5}{x+8}.$$

$$24. \frac{x+2}{x} + \frac{x-7}{x-5} - \frac{x+3}{x+1} = \frac{x-6}{x-4}.$$

$$25. \frac{4x-17}{x-4} + \frac{10x-13}{2x-3} = \frac{8x-30}{2x-7} + \frac{5x-4}{x-1}.$$

$$26. \frac{5x-8}{x-2} + \frac{6x-44}{x-7} - \frac{10x-8}{x-1} = \frac{x-8}{x-6}.$$

$$27. \frac{2x-3}{3x-4} = \frac{4x-6}{06x-07}.$$

$$28. \frac{x-2}{05} - \frac{x-4}{0625} = 56.$$

$$29. 083(x-625) = 09(x-59375).$$

$$30. (2x+1.5)(3x-2.25) = (2x-1.125)(3x+1.25).$$

$$31. \frac{3x-1}{5x-4} = \frac{5+1.2x}{2x-1}.$$

$$32. \frac{1-1.4x}{2+x} = \frac{7(x-1)}{1-5x}.$$

$$33. \frac{(3x-2)(3x-1)}{2x-1} - \frac{1}{6}(3x-2) = 4x-2.$$

we obtain

Literal Equations.

187. In the equations we have discussed hitherto the coefficients have been numerical quantities, but equations often involve *literal* coefficients. [Art. 6.] These are supposed to be known, and will appear in the solution.

Example 1. Solve $(x+a)(x+b)-c(a+c)=(x-c)(x+c)+ab$.

Multiplying out, we have

$$x^2+ax+bx+ab-ac-c^2=x^2-c^2+ab;$$

whence

$$ax+bx=ac,$$

$$(a+b)x=ac;$$

$$\therefore x=\frac{ac}{a+b}.$$

Example 2. Solve $\frac{a}{x-a}-\frac{b}{x-b}=\frac{a-b}{x-c}$.

Simplifying the left side, we have

$$\frac{a(x-b)-b(x-a)}{(x-a)(x-b)}=\frac{a-b}{x-c}$$

$$\frac{(a-b)x}{(x-a)(x-b)}=\frac{a-b}{x-c};$$

$$\therefore \frac{x}{(x-a)(x-b)}=\frac{1}{x-c}.$$

Multiplying across,

$$x^2-cx=x^2-ax-bx+ab,$$

$$ax+bx-cx=ab,$$

$$(a+b-c)x=ab;$$

$$\therefore x=\frac{ab}{a+b-c}.$$

EXAMPLES XXIII. b.

Solve the equations:

1. $ax-2b=5bx-3a.$

2. $a^2(x-a)+b^2(x-b)=abx.$

3. $x^2+a^2=(b-x)^2.$

4. $(x-a)(x+b)=(x-a+b)^2.$

5. $a(x-2)+2x=6+a.$

6. $m^2(m-x)-mnx=n^2(n+x).$

7. $(a+x)(b+x)=x(x-c).$

8. $(a-b)(x-a)=(a-c)(x-b).$

9. $\frac{2x+3a}{x+a}=\frac{2(3x+2a)}{3x+a}.$

10. $\frac{2(x-b)}{3x-c}=\frac{2x+b}{3(x-c)}.$

11. $\frac{1}{a}-$

13. $\frac{a}{x}=$

15. $\frac{x-c}{b-c}$

17. $\frac{1}{4}x($

18. $(a+$

19. $b(a$

20. $b(a$

21. x^2+

22. $(2x$

23. $\frac{x-c}{x-}$

24. $\left(\frac{x}{a}-$

25. $b(x-\frac{x^2}{x^2-}$

Examp

The no
meet wit
choose ce
correspon
in the sec
the value
often con
some con
memory.
of x ; b , b
Sometim
 a_1 , a_2 , a_3 ;

11. $\frac{1}{a} - \frac{1}{x} = \frac{1}{x} - \frac{1}{b}$. 12. $\frac{2}{3}\left(\frac{x}{a} + 1\right) = \frac{3}{4}\left(\frac{x}{a} - 1\right)$.
13. $\frac{a}{x} = c(a - b) + \frac{b}{x}$. 14. $\frac{9a}{b} - \frac{3x}{b} = \frac{4b}{a} - \frac{2x}{a}$.
15. $\frac{x - a}{b - x} = \frac{x - b}{a - x}$. 16. $\frac{x - a}{2} = \frac{(x - b)^2}{2x - a}$.
17. $\frac{1}{4}x(x - a) - \left(\frac{x + a}{2}\right)^2 = \frac{2a}{3}\left(x - \frac{a}{2}\right)$.
18. $(a + b)x^2 - a(bx + a^2) = bx(x - a) + ax(x - b)$.
19. $b(a + x) - (a + x)(b - x) = x^2 + \frac{bc^2}{a}$.
20. $b(a - x) - \frac{a}{b}(b + x)^2 + ab\left(\frac{x}{b} + 1\right)^2 = 0$.
21. $x^2 + a(2a - x) - \frac{3b^2}{4} = \left(x - \frac{b}{2}\right)^2 + a^2$.
22. $(2x - a)\left(x + \frac{2a}{3}\right) = 4x\left(\frac{a}{3} - x\right) - \frac{1}{2}(a - 4x)(2a + 3x)$.
23. $\frac{x - a + b}{x - a} + \frac{x - b}{x - 2b} = \frac{x}{x - b} + \frac{x - a}{x - a - b}$.
24. $\left(\frac{x}{a} - 3\right)\left(\frac{3x}{a} - 1\right) - \frac{1}{a^2}(x - 2a)(2x - a) = \left(\frac{x}{a} - 1\right)^2 - 1$.
25. $\frac{b(x + a)}{x^2 - b^2} + \frac{2x + 3b - a}{x + b} = \frac{2(x^2 + bx - b^2)}{x^2 - b^2}$.

Example 3. Solve $ax + by = c$ (1),
 $a'x + b'y = c'$ (2).

The notation here first used is one that the student will frequently meet with in the course of his reading. In the first equation we choose certain letters as the coefficients of x and y , and we choose *corresponding letters with accents* to denote corresponding quantities in the second equation. There is no necessary connection between the values of a and a' , and they are as different as a and b ; but it is often convenient to use the same letter thus slightly varied to mark some common meaning of such letters, and thereby assist the memory. Thus a, a' have a common property as being coefficients of x ; b, b' as being coefficients of y .

Sometimes instead of accents letters are used with a *suffix*, such as a_1, a_2, a_3 ; b_1, b_2, b_3 , etc.

To return to the equations $ax+by=c$ (1),

$$a'x+b'y=c' \text{(2).}$$

Multiply (1) by b' and (2) by b . Thus

$$ab'x+bb'y=b'c,$$

$$a'bx+bb'y=bc';$$

by subtraction,

$$(ab'-a'b)x=b'c-bc';$$

$$\therefore x=\frac{b'c-bc'}{ab'-a'b} \text{(3).}$$

As previously explained in Art. 104, we might obtain y by substituting this value of x in either of the equations (1) or (2); but y is more conveniently found by eliminating x , as follows:

Multiplying (1) by a' and (2) by a , we have

$$aa'x+a'by=a'c,$$

$$aa'x+ab'y=ac';$$

by subtraction,

$$(a'b-ab')y=a'c-ac';$$

$$\therefore y=\frac{a'c-ac'}{a'b-ab'},$$

or, changing signs in the terms of the denominator so as to have the same denominator as in (3),

$$y=\frac{ac'-a'c}{ab'-a'b}, \text{ and } x=\frac{b'c-bc'}{ab'-a'b}.$$

Example 4. Solve $\frac{x-a}{c-a}+\frac{y-b}{c-b}=1$ (1),

$$\frac{x+a}{c}+\frac{y-a}{a-b}=\frac{a}{c} \text{(2).}$$

From (1) by clearing of fractions, we have

$$x(c-b)-a(c-b)+y(c-a)-b(c-a)=(c-a)(c-b),$$

$$x(c-b)+y(c-a)=ac-ab+bc-ab+c^2-ac-bc+ab,$$

$$x(c-b)+y(c-a)=c^2-ab \text{(3).}$$

Again, from (2), we have

$$x(a-b)+a(a-b)+cy-ca=a(a-b)$$

$$x(a-b)+cy=ac \text{(4).}$$

Multiply (3) by c and (4) by $c-a$ and subtract,

$$x\{c(c-b)-(c-a)(a-b)\}=c^3-abc-ac(c-a),$$

$$x(c^2-ac+a^2-ab)=c(c^2-ab-ac+a^2);$$

$$\therefore x=c;$$

and therefore from (4)

$$y=b.$$

Solve

1. $\frac{ax+b}{bx+a}$

4. $\frac{ax+b}{bx+a}$

7. $\frac{x}{a} + \frac{x}{a}$

$\frac{x}{a} - \frac{x}{a}$

10. $\frac{qx}{qx+a}$

13. $(a-x)$

15. $\frac{x}{a}$

$\frac{x}{3a} + \frac{x}{3a}$

18. $\frac{m}{l}$

$(x+l)$

19. bx

ax

20. $(a-x)$

21. x

EXAMPLES XXIII. c.

Solve the equations :

1. $ax + by = l,$
 $bx + ay = m.$
2. $lx + my = n,$
 $px + qy = r.$
3. $ax = by,$
 $bx + ay = c.$
4. $ax + by = a^2,$
 $bx + ay = b^2.$
5. $x + ay = a',$
 $ax + a'y = 1.$
6. $px - qy = r,$
 $rx - py = q.$
7. $\frac{x}{a} + \frac{y}{b} = \frac{1}{ab},$
 $\frac{x}{a'} - \frac{y}{b'} = \frac{1}{a'b'}.$
8. $\frac{x}{a} - \frac{y}{b} = 0,$
 $bx + ay = 4ab$
9. $\frac{3x}{a} + \frac{2y}{b} = 3,$
 $\frac{9x}{a} - \frac{6y}{b} = 3.$
10. $qx - ry = p(a - y),$
 $\frac{qx}{a} + r = p\left(1 + \frac{y}{b}\right).$
11. $\frac{x}{m} + \frac{y}{m'} = 1,$
 $\frac{x}{m'} - \frac{y}{m} = 1.$
12. $px + qy = 0,$
 $lx + my = n.$
13. $(a - b)x = (a + b)y,$
 $x + y = c.$
14. $(a - b)x + (a + b)y = 2a^2 - 2b^2,$
 $(a + b)x - (a - b)y = 4ab.$
15. $\frac{x}{a} + \frac{y}{b} = 1,$
 $\frac{x}{3a} + \frac{y}{6b} = \frac{2}{3}.$
16. $\frac{x}{a} + \frac{y}{b} = 2,$
 $\frac{x}{a} = \frac{y}{b}.$
17. $\frac{x}{a} - \frac{y}{b} = 1,$
 $\frac{x}{b} + \frac{y}{a} = \frac{a}{b}.$
18. $\frac{m}{l}x + \frac{l}{m}y = \left(\frac{1}{l} + \frac{1}{m}\right)(m^2 + l^2),$
 $(x + y)(m^2 + l^2) = 2(m^3 + l^3) + ml(x + y).$
19. $bx + cy = a + b,$
 $ax\left(\frac{1}{a-b} - \frac{1}{a+b}\right) + cy\left(\frac{1}{b-a} - \frac{1}{b+a}\right) = \frac{2a}{a+b}.$
20. $(a - b)x + (a + b)y = 2(a^2 - b^2),$ $ax - by = a^2 + b^2.$
21. $x\left(a - b + \frac{ab}{a-b}\right) = y\left(a + b - \frac{ab}{a+b}\right),$ $x + y = 2a^3.$

Hence

Thus t

If the
and 5 o'clock
must also
13 divisions
gains 20

Hence

which gi

Theref

Exampl
two plac
at the ra
will A h

Suppo

A wal

hours; a
being eq

whence

Theref

Exampl
Had the
occupied
the jour

Let th
occupied
by xy mi

CHAPTER XXIV.

HARDER PROBLEMS.

188. IN previous chapters we have given collections of problems which lead to simple equations. We add here a few examples of somewhat greater difficulty.

Example 1. A grocer buys 15 lbs. of figs and 28 lbs. of currants for £1. 1s. 8d.; by selling the figs at a loss of 10 per cent., and the currants at a gain of 30 per cent., he clears 2s. 6d. on his outlay; how much per pound did he pay for each?

Let x, y denote the number of pence in the price of a pound of figs and currants respectively; then the outlay is

$$15x + 28y \text{ pence.}$$

Therefore

$$15x + 28y = 260 \dots\dots\dots(1).$$

The loss upon the figs is $\frac{1}{10} \times 15x$ pence, and the gain upon the currants is $\frac{3}{10} \times 28y$ pence; therefore the total gain is

$$\frac{42y}{5} - \frac{3x}{2} \text{ pence;}$$

$$\therefore \frac{42y}{5} - \frac{3x}{2} = 30 \dots\dots\dots(2).$$

From (1) and (2) we find that $x=8$, and $y=5$; that is the figs cost 8d. a pound, and the currants cost 5d. a pound.

Example 2. At what time between 4 and 5 o'clock will the minute-hand of a watch be 13 minutes in advance of the hour-hand?

Let x denote the required number of minutes after 4 o'clock; then, as the minute-hand travels twelve times as fast as the hour-hand, the hour-hand will move over $\frac{x}{12}$ minute divisions in x minutes.

At 4 o'clock the minute-hand is 20 divisions behind the hour-hand, and finally the minute-hand is 13 divisions in advance; therefore the minute-hand moves over $20 + 13$, or 33 divisions more than the hour-hand.

Hence

$$x = \frac{x}{12} + 33,$$

$$\frac{11}{12}x = 33;$$

$$\therefore x = 36.$$

Thus the time is 36 minutes past 4.

If the question be asked as follows: "At what times between 4 and 5 o'clock will there be 13 minutes between the two hands?" we must also take into consideration the case when the minute-hand is 13 divisions *behind* the hour-hand. In this case the minute-hand gains $20 - 13$, or 7 divisions.

Hence

$$x = \frac{x}{12} + 7,$$

which gives

$$x = 7\frac{7}{11}.$$

Therefore the times are $7\frac{7}{11}$ past 4, and 36' past 4.

Example 3. Two persons *A* and *B* start simultaneously from two places, *c* miles apart, and walk in the same direction. *A* travels at the rate of *p* miles an hour, and *B* at the rate of *q* miles; how far will *A* have walked before he overtakes *B*?

Suppose *A* has walked *x* miles, then *B* has walked *x - c* miles.

A walking at the rate of *p* miles an hour will travel *x* miles in $\frac{x}{p}$ hours; and *B* will travel *x - c* miles in $\frac{x-c}{q}$ hours: these two times being equal, we have

$$\frac{x}{p} = \frac{x-c}{q},$$

$$qx = px - pc;$$

whence

$$x = \frac{pc}{p-q}.$$

Therefore *A* has travelled $\frac{pc}{p-q}$ miles.

Example 4. A train travelled a certain distance at a uniform rate. Had the speed been 6 miles an hour more, the journey would have occupied 4 hours less; and had the speed been 6 miles an hour less, the journey would have occupied 6 hours more. Find the distance.

Let the speed of the train be *x* miles per hour, and let the time occupied be *y* hours; then the distance traversed will be represented by *xy* miles.

On the first supposition the speed per hour is $x+6$ miles, and the time taken is $y-4$ hours. In this case the distance traversed will be represented by $(x+6)(y-4)$ miles.

On the second supposition the distance traversed will be represented by $(x-6)(y+6)$ miles.

All these expressions for the distance must be equal ;

$$\therefore xy = (x+6)(y-4) = (x-6)(y+6).$$

From these equations we have

$$\begin{aligned} xy &= xy + 6y - 4x - 24, \\ \text{or} \quad 6y - 4x &= 24 \dots\dots\dots(1); \end{aligned}$$

$$\begin{aligned} \text{and} \quad xy &= xy - 6y + 6x - 36, \\ \text{or} \quad 6x - 6y &= 36 \dots\dots\dots(2). \end{aligned}$$

From (1) and (2) we obtain $x=30$, $y=24$.

Hence the distance is 720 miles.

Example 5. A person invests £3770, partly in 3 per cent. Stock at £102, and partly in Railway Stock at £84 which pays a dividend of $4\frac{1}{2}$ per cent. : if his income from these investments is £136. 5s. per annum, what sum does he invest in each ?

Let x denote the number of pounds invested in 3 per cent., y the number of pounds invested in Railway Stock ; then

$$x + y = 3770 \dots\dots\dots(1).$$

The income from 3 per cent. Stock is $\pounds \frac{3x}{102}$, or $\pounds \frac{x}{34}$; and

that from Railway Stock is $\pounds \frac{4\frac{1}{2}y}{84}$, or $\pounds \frac{3y}{56}$.

$$\text{Therefore} \quad \frac{x}{34} + \frac{3y}{56} = 136\frac{1}{4} \dots\dots\dots(2).$$

$$\text{From (2)} \quad x + \frac{51}{28}y = 4632\frac{1}{2} ;$$

therefore by subtracting (1)

$$\frac{23}{28}y = 862\frac{1}{2} ;$$

$$\text{whence} \quad y = 28 \times 37\frac{1}{2} = 1050 ;$$

$$\text{and from (1)} \quad x = 2720.$$

Therefore he invests £2720 in 3 per cent Stock, and £1050 in Railway Stock.

EXAMPLES XXIV.

1. A sum of £10 is divided among a number of persons ; if the number had been increased by one-fourth each would have received a shilling less : find the number of persons.

2. I b
one-fifth c
penny : h

3. I b
they had
less : how

4. A n
and then
afterward
originally

5. I sp
silk ; the
pence per

6. A n
digits by
number.

7. The
digits be
original n

8. A n
my preser
ago her a
were their

9. At
watch first

10. At
one minut

11. W
of 6 and 7

12. It
hand will
what is th

13. At
elevenths
the votes

14. A
ten more
each woul
what each

15. A
4s. a yard
how much

2. I bought a certain number of eggs at four a penny; I kept one-fifth of them, and sold the rest at three a penny, and gained a penny: how many did I buy?

3. I bought a certain number of articles at five for sixpence; if they had been eleven for one shilling, I should have spent sixpence less: how many did I buy?

4. A man at whist wins twice as much as he had to begin with, and then loses 16s.; he then loses four-fifths of what remained, and afterwards wins as much as he had at first: how much had he originally, if he leaves off with £4?

5. I spend £14. 5s. in buying 20 yards of calico and 30 yards of silk; the silk costs as many shillings per yard as the calico costs pence per yard: find the price of each.

6. A number of two digits exceeds five times the sum of its digits by 9, and its ten-digit exceeds its unit-digit by 1: find the number.

7. The sum of the digits of a number less than 100 is 6; if the digits be reversed the resulting number will be less by 18 than the original number: find it.

8. A man being asked his age replied, "If you take 2 years from my present age the result will be double my wife's age, and 3 years ago her age was one-third of what mine will be in 12 years." What were their ages?

9. At what time between one and two o'clock are the hands of a watch first at right angles?

10. At what time between 3 and 4 o'clock is the minute-hand one minute ahead of the hour hand?

11. When are the hands of a clock together between the hours of 6 and 7?

12. It is between 2 and 3 o'clock, and in 10 minutes the minute-hand will be as much before the hour-hand as it is now behind it: what is the time?

13. At an election the majority was 162, which was three-elevenths of the whole numbers of voters: what was the number of the votes on each side?

14. A certain number of persons paid a bill; if there had been ten more each would have paid 2s. less; if there had been 5 less each would have paid 2s. 6d. more: find the number of persons, and what each had to pay.

15. A man spends £5 in buying two kinds of silk at 4s. 6d. and 4s. a yard; by selling it all at 4s. 3d. per yard he gains 2 per cent.: how much of each did he buy?

16. Ten years ago the sum of the ages of two sons was one-third of their father's age : one is two years older than the other, and the present sum of their ages is fourteen years less than their father's age : how old are they ?

17. *A* and *B* start from the same place walking at different rates ; when *A* has walked 15 miles *B* doubles his pace, and 6 hours later passes *A* : if *A* walks at the rate of 5 miles an hour, what is *B*'s rate at first ?

18. A basket of oranges is emptied by one person taking half of them and one more, a second person taking half of the remainder and one more, and a third person taking half of the remainder and six more. How many did the basket contain at first ?

19. A person swimming in a stream which runs $1\frac{1}{2}$ miles per hour, finds that it takes him four times as long to swim a mile up the stream as it does to swim the same distance down : at what rate does he swim ?

20. At what times between 7 and 8 o'clock will the hands of a watch be at right angles to each other ? When will they be in the same straight line ?

21. The denominator of a fraction exceeds the numerator by 4 ; and if 5 is taken from each, the sum of the reciprocal of the new fraction and four times the original fraction is 5 : find the original fraction.

22. Two persons start at noon from towns 60 miles apart. One walks at the rate of four miles an hour, but stops $2\frac{1}{2}$ hours on the way ; the other walks at the rate of 3 miles an hour without stopping : when and where will they meet ?

23. *A*, *B*, and *C* travel from the same place at the rates of 4, 5, and 6 miles an hour respectively ; and *B* starts 2 hours after *A*. How long after *B* must *C* start in order that they may overtake *A* at the same instant ?

24. A dealer bought a horse, expecting to sell it again at a price that would have given him 10 per cent. profit on his purchase ; but he had to sell it for £50 less than he expected, and he then found that he had lost 15 per cent. on what it cost him : what did he pay for the horse ?

25. A man walking from a town, *A*, to another, *B*, at the rate of 4 miles an hour, starts one hour before a coach travelling 12 miles an hour, and is picked up by the coach. On arriving at *B*, he finds that his coach journey has lasted 2 hours : find the distance between *A* and *B*.

26. What is the property of a person whose income is £1140, when one-twelfth of it is invested at 2 per cent., one-half at 3 per cent., one-third at $4\frac{1}{2}$ per cent., and the remainder pays him no dividend ?

27. *A* and pays on debts was the a

28. *T* there is t as much each to fil contents n

29. *T* tains twi much win to fill a p mixed ?

30. *T* *B*, and th walks at hour : at

31. *A* Birmingham 15 miles hour, pass of each li

32. *C* proportion selling the

33. *A* another a form a mi without l

34. *A* shillings a 3 times as same mon

35. *A* of a miles If he had have ridd that

36. *A*, walk from to his star minutes r he starts takes $c + 2$

27. A person spends one-third of his income, saves one-fourth, and pays away 5 per cent. on the whole as interest at $7\frac{1}{2}$ per cent. on debts previously incurred, and then has £110 remaining: what was the amount of his debts?

28. Two vessels contain mixtures of wine and water; in one there is three times as much wine as water, in the other, five times as much water as wine. Find how much must be drawn off from each to fill a third vessel which holds seven gallons, in order that its contents may be half wine and half water.

29. There are two mixtures of wine and water, one of which contains twice as much water as wine, and the other three times as much wine as water. How much must there be taken from each to fill a pint cup, in which the water and the wine shall be equally mixed?

30. Two men set out at the same time to walk, one from A to B , and the other from B to A , a distance of a miles. The former walks at the rate of p miles, and the latter at the rate of q miles an hour: at what distance from A will they meet?

31. A train on the North Western line passes from London to Birmingham in 3 hours; a train on the Great Western line which is 15 miles longer, travelling at a speed which is less by 1 mile per hour, passes from one place to the other in $3\frac{1}{2}$ hours: find the length of each line.

32. Coffee is bought at 1s. and chicory at 3d. per lb.; in what proportion must they be mixed that 10 per cent. may be gained by selling the mixture at 11d. per lb.?

33. A man has one kind of coffee at a pence per pound, and another at b pence per pound. How much of each must he take to form a mixture of $a-b$ lbs., which he can sell at c pence a pound without loss?

34. A man spends c half-crowns in buying two kinds of silk at a shillings and b shillings a yard respectively; he could have bought 3 times as much of the first and half as much of the second for the same money: how many yards of each did he buy?

35. A man rides one-third of the distance from A to B at the rate of a miles an hour, and the remainder at the rate of $2b$ miles an hour. If he had travelled at a uniform rate of $3c$ miles an hour, he could have ridden from A to B and back again in the same time. Prove that

$$\frac{2}{c} = \frac{1}{a} + \frac{1}{b}.$$

36. A , B , C are three towns forming a triangle. A man has to walk from one to the next, ride thence to the next, and drive thence to his starting point. He can walk, ride, and drive a mile in a , b , c minutes respectively. If he starts from B he takes $a+c-b$ hours, if he starts from C he takes $b+a-c$ hours, and if he starts from A he takes $c+b-a$ hours. Find the length of the circuit.

CHAPTER XXV.

QUADRATIC EQUATIONS.

189. SUPPOSE the following problem were proposed for solution :

A dealer bought a number of horses for £280. If he had bought four less each would have cost £8 more : how many did he buy ?

We should proceed thus :

Let x = the number of horses ; then $\frac{280}{x}$ = the number of pounds each cost.

If he had bought 4 less he would have had $x - 4$ horses, and each would have cost $\frac{280}{x-4}$ pounds.

$$\therefore 8 + \frac{280}{x} = \frac{280}{x-4} ;$$

whence

$$x(x-4) + 35(x-4) = 35x ;$$

$$\therefore x^2 - 4x + 35x - 140 = 35x ;$$

$$\therefore x^2 - 4x = 140.$$

Here we have an equation which involves the *square* of the unknown quantity ; and in order to complete the solution of the problem we must discover a method of solving such equations.

190. DEFINITION. An equation which contains the square of the unknown quantity, but no higher power, is called a **quadratic equation**, or an **equation of the second degree**.

If the equation contains both the square and the first power of the unknown it is called an *adfectied* quadratic ; if it contains only the square of the unknown it is said to be a *pure* quadratic.

Thus $2x^2 - 5x = 3$ is an adfectied quadratic,
and $5x^2 = 20$ is a pure quadratic.

CHAP. XX.

191.

tion in v

Exam

Multip

and taki

Note.

side for t

192.

equation

double s

But an

necessary

and thes

$x = +6$,

the two

sign befo

193.

form of

solved in

sides, we

Taking th

taking th

$\therefore$ the s

Now th

may be w

or

Hence, by

can be s

extracting

each side

perfect sq

E.A.

191. A pure quadratic may be considered as a simple equation in which the *square* of the unknown quantity is to be found.

Example. Solve $\frac{9}{x^2-27} = \frac{25}{x^2-11}$.

Multiplying up, $9x^2 - 99 = 25x^2 - 675$;
 $\therefore 16x^2 = 576$;
 $\therefore x^2 = 36$;

and taking the square root of these equals, we have
 $x = \pm 6$.

Note. We prefix the double sign to the number on the right-hand side for the reason given in Art. 117.

192. In extracting the square root of the two sides of the equation $x^2 = 36$, it might seem that we ought to prefix the double sign to the quantities on both sides, and write $\pm x = \pm 6$. But an examination of the various cases shews this to be unnecessary. For $\pm x = \pm 6$ gives the four cases :

$$+x = +6, +x = -6, -x = +6, -x = -6,$$

and these are all included in the two already given, namely, $x = +6, x = -6$. Hence, when we extract the square root of the two sides of an equation, it is sufficient to put the double sign before the square root of *one* side.

193. The equation $x^2 = 36$ is an instance of the simplest form of quadratic equations. The equation $(x-3)^2 = 25$ may be solved in a similar way ; for taking the square root of both sides, we have two *simple* equations,

$$x-3 = \pm 5.$$

Taking the upper sign,

$$x-3 = +5, \text{ whence } x=8 ;$$

taking the lower sign,

$$x-3 = -5, \text{ whence } x = -2.$$

$\therefore$ the solution is

$$x=8, \text{ or } -2.$$

Now the given equation, $(x-3)^2 = 25$

may be written $x^2 - 6x + (3)^2 = 25,$

or

$$x^2 - 6x = 16.$$

Hence, by retracing our steps, we learn that the equation

$$x^2 - 6x = 16.$$

can be solved by first adding $(3)^2$ or 9 to each side, and then extracting the square root ; and the reason why we add 9 to each side is that this quantity added to the left side makes it a *perfect square*.

Now whatever the quantity a may be,

$$x^2 + 2ax + a^2 = (x + a)^2,$$

and

$$x^2 - 2ax + a^2 = (x - a)^2;$$

so that if a trinomial is a perfect square, and *its highest power, x^2 , has unity for its coefficient*, we must always have the term without x equal to the square of half the coefficient of x . If, therefore, the terms in x^2 and x are given, the square may be completed by adding the square of half the coefficient of x .

Note. When an expression is a perfect square, the *square terms* are always *positive*. [Art. 114, Note.] Hence, if necessary, the coefficient of x^2 must be made equal to +1 before completing the square.

Example 1. Solve $x^2 + 14x = 32$.

The square of half 14 is $(7)^2$.

$$\therefore x^2 + 14x + (7)^2 = 32 + 49;$$

that is,

$$(x + 7)^2 = 81;$$

$$\therefore x + 7 = \pm 9;$$

$$\therefore x = -7 + 9, \text{ or } -7 - 9;$$

$$\therefore x = 2, \text{ or } -16.$$

Example 2. Solve $7x = x^2 - 8$.

Transpose so as to have the terms involving x on one side, and the square term positive,

$$\text{Thus } x^2 - 7x = 8.$$

$$\text{Completing the square, } x^2 - 7x + \left(\frac{7}{2}\right)^2 = 8 + \frac{49}{4};$$

that is,

$$\left(x - \frac{7}{2}\right)^2 = \frac{81}{4};$$

$$\therefore x - \frac{7}{2} = \pm \frac{9}{2};$$

$$\therefore x = \frac{7}{2} \pm \frac{9}{2};$$

$$\therefore x = 8, \text{ or } -1.$$

Note. We do not work out $\left(\frac{7}{2}\right)^2$ on the left-hand side.

EXAMPLES XXV. a.

1. $5(x^2 + 5) = 6x^2$.

2. $3x^2 = 4(x^2 - 4)$.

3. $x^2 + 22x = 75$.

4. $x^2 + 24x = 25$.

5. $x^2 = 10x - 21$.

6. $(9 + x)(9 - x) = 17$.

7. $x^2 + 3x = 18$.

8. $x^2 + 5x = 14$.

9. $x^2 - 5x - 36 = 0$.

10. $x^2 =$

13. 68

16. $23x$

19. $x^2 -$

22. $\frac{19}{5}x$

194.
 pleted w
 reduced
 coefficient

Examp

Transp

Divide

Thus

completing

that is,

Note.

Example

Dividing

completing

that is,

10. $x^2 = x + 72$. 11. $x^2 - 341 = 20x$. 12. $9x - x^2 + 220 = 0$.
 13. $68 - x^2 = 13x$. 14. $x + 156 = x^2$. 15. $187 = x^2 + 6x$.
 16. $23x = 120 + x^2$. 17. $42 + x^2 = 13x$. 18. $22x + 23 - x^2 = 0$.
 19. $x^2 - \frac{2}{3}x = 32$. 20. $x^2 + \frac{4}{15}x = \frac{1}{5}$. 21. $x^2 - \frac{7}{6}x - \frac{1}{2} = 0$.
 22. $\frac{19}{5}x = \frac{4}{5} - x^2$. 23. $\frac{3}{5}(x+6)(x-2) = \frac{2}{3}\left(62\frac{1}{10} + \frac{18x}{5}\right)$.

194. We have shewn that the square may readily be completed when the coefficient of x^2 is unity. All cases may be reduced to this by dividing the equation throughout by the coefficient of x^2 .

Example 1. Solve $32 - 3x^2 = 10x$.

Transposing, $3x^2 + 10x = 32$.

Divide throughout by 3, so as to make the coefficient of x^2 unity.

Thus $x^2 + \frac{10}{3}x = \frac{32}{3}$;

completing the square, $x^2 + \frac{10}{3}x + \left(\frac{5}{3}\right)^2 = \frac{32}{3} + \frac{25}{9}$;

that is, $\left(x + \frac{5}{3}\right)^2 = \frac{121}{9}$;

$$\therefore x + \frac{5}{3} = \pm \frac{11}{3};$$

$$\therefore x = -\frac{5}{3} \pm \frac{11}{3} = 2, \text{ or } -5\frac{1}{3}.$$

Note. We do not add $\left(\frac{10}{6}\right)^2$ but $\left(\frac{5}{3}\right)^2$ to the left-hand side.

Example 2. Solve $5x^2 + 11x = 12$.

Dividing by 5, $x^2 + \frac{11}{5}x = \frac{12}{5}$;

completing the square, $x^2 + \frac{11}{5}x + \left(\frac{11}{10}\right)^2 = \frac{12}{5} + \frac{121}{100}$;

that is, $\left(x + \frac{11}{10}\right)^2 = \frac{361}{100}$;

$$\therefore x + \frac{11}{10} = \pm \frac{19}{10};$$

$$\therefore x = -\frac{11}{10} \pm \frac{19}{10} = \frac{4}{5}, \text{ or } -3.$$

195. We see then that the following are the steps required for solving an affected quadratic equation :

(1) If necessary, simplify the equation so that the terms in x^2 and x are on one side of the equation, and the term without x on the other.

(2) Make the coefficient of x^2 unity and positive by dividing throughout by the coefficient of x^2 .

(3) Add to each side of the equation the square of half the coefficient of x .

(4) Take the square root of each side.

(5) Solve the resulting simple equations.

196. In the examples which follow some preliminary reduction and simplification may be necessary.

Example 1. Solve $\frac{3x-2}{2x-3} = \frac{5x}{x+4} - 2$.

Simplifying, $\frac{3x-2}{2x-3} = \frac{3x-8}{x+4}$;

multiplying across, $3x^2 + 10x - 8 = 6x^2 - 25x + 24$;

that is, $-3x^2 + 35x = 32$.

Dividing by -3 , $x^2 - \frac{35}{3}x = -\frac{32}{3}$;

completing the square, $x^2 - \frac{35}{3}x + \left(\frac{35}{6}\right)^2 = \frac{1225}{36} - \frac{32}{3}$;

that is, $\left(x - \frac{35}{6}\right)^2 = \frac{841}{36}$;

$\therefore x - \frac{35}{6} = \pm \frac{29}{6}$;

$\therefore x = 10\frac{2}{3}$, or 1 .

Example 2. Solve $7(x+2a)^2 + 3a^2 = 5a(7x+23a)$.

Simplifying, $7x^2 + 28ax + 28a^2 + 3a^2 = 35ax + 115a^2$,

that is, $7x^2 - 7ax = 84a^2$.

Whence $x^2 - ax = 12a^2$;

completing the square, $x^2 - ax + \left(\frac{a}{2}\right)^2 = 12a^2 + \frac{a^2}{4}$;

that is, $\left(x - \frac{a}{2}\right)^2 = \frac{49a^2}{4}$;

$\therefore x - \frac{a}{2} = \pm \frac{7a}{2}$;

$\therefore x = 4a$, or $-3a$.

whence
similar
two equ

1. $5x^2$

4. $8x^2$

7. $15x$

10. $12x$

13. $21x$

15. $18x$

17. $15x$

19. $6x^2$

21. $12x$

23. $(x -$

25. $\frac{5x}{x -$

28. $\frac{3x}{4x}$

30. $\frac{x +}{2x}$

32. $\frac{x +}{x -$

34. $\frac{4}{x -$

36. $\frac{x -}{x -$

38. $\frac{2}{3x}$

198.

suitable

can be v

where a

therefor

197. Sometimes there is *only one solution*. Thus if

$$x^2 - 2x + 1 = 0, \text{ then } (x-1)^2 = 0,$$

whence $x=1$ is the only solution. Nevertheless, in this and similar cases we find it convenient to say that the quadratic has *two equal roots*.

EXAMPLES XXV. b.

1. $5x^2 + 14x = 55.$
2. $3x^2 + 121 = 44x.$
3. $25x = 6x^2 + 21.$
4. $8x^2 + x = 30.$
5. $3x^2 + 35 = 22x.$
6. $x + 22 - 6x^2 = 0.$
7. $15 = 17x + 4x^2.$
8. $21 + x = 2x^2.$
9. $9x^2 - 143 - 6x = 0.$
10. $12x^2 = 29x - 14.$
11. $20x^2 = 12 - x.$
12. $19x = 15 - 8x^2.$
13. $21x^2 + 22x + 5 = 0.$
14. $50x^2 - 15x = 27.$
15. $18x^2 - 27x - 26 = 0.$
16. $5x^2 = 8x + 21.$
17. $15x^2 - 2ax = a^2.$
18. $21x^2 = 2ax + 3a^2.$
19. $6x^2 = 11kx + 7k^2.$
20. $12x^2 + 23kx + 10k^2 = 0.$
21. $12x^2 - cx - 20c^2 = 0.$
22. $2(x-3) = 3(x+2)(x-3).$
23. $(x+1)(2x+3) = 4x^2 - 22.$
24. $(3x-5)(2x-5) = x^2 + 2x - 3.$
25. $\frac{5x+7}{x-1} = 3x+2.$
26. $\frac{5x-1}{x+1} = \frac{3x}{2}.$
27. $\frac{3x-8}{x-2} = \frac{5x-2}{x+5}.$
28. $\frac{3x-1}{4x+7} = 1 - \frac{6}{x+7}.$
29. $\frac{5x-7}{7x-5} = \frac{x-5}{2x-13}.$
30. $\frac{x+3}{2x-7} - \frac{2x-1}{x-3} = 0.$
31. $\frac{1}{1+x} - \frac{1}{3-x} = \frac{6}{35}.$
32. $\frac{x+4}{x-4} + \frac{x-2}{x-3} = 6\frac{1}{5}.$
33. $\frac{1}{3-x} - \frac{4}{5} = \frac{1}{9-2x}.$
34. $\frac{4}{x-1} - \frac{5}{x+2} = \frac{3}{x}.$
35. $\frac{5}{x-2} - \frac{4}{x} = \frac{3}{x+6}.$
36. $\frac{x-2}{x-3} + \frac{3x-11}{x-4} = \frac{4x+13}{x+1}.$
37. $\frac{1}{2x-5a} + \frac{5}{2x-a} = \frac{2}{a}.$
38. $\frac{2}{3x-2c} + \frac{3}{2x-3c} = \frac{7}{2c}.$
39. $\frac{a^2b}{x^2} + \left(1 + \frac{b}{x}\right)a = 2b + \frac{a^2}{x}.$

198. From the preceding examples it appears that after suitable reduction and transposition every quadratic equation can be written in the form

$$ax^2 + bx + c = 0,$$

where a, b, c may have any numerical values whatever. If therefore we can solve this quadratic we can solve any.

Transposing, $ax^2 + bx = -c$;
 dividing by a , $x^2 + \frac{b}{a}x = -\frac{c}{a}$.

Completing the square by adding to each side $\left(\frac{b}{2a}\right)^2$,

$$x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 = \frac{b^2}{4a^2} - \frac{c}{a};$$

that is, $\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2};$

extracting the square root,

$$x + \frac{b}{2a} = \frac{\pm \sqrt{(b^2 - 4ac)}}{2a};$$

$$x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a}.$$

199. Instead of going through the process of completing the square in each particular example, we may now make use of this general formula, adapting it to the case in question by substituting the values of a , b , c .

Example. Solve $5x^2 + 11x - 12 = 0$.

Here $a=5$, $b=11$, $c=-12$.

$$\therefore x = \frac{-11 \pm \sqrt{(11)^2 - 4 \cdot 5 \cdot (-12)}}{10}$$

$$= \frac{-11 \pm \sqrt{361}}{10} = \frac{-11 \pm 19}{10} = \frac{4}{5}, \text{ or } -3,$$

which agrees with the solution of Example 2, Art. 194.

200. In the result $x = \frac{-b \pm \sqrt{(b^2 - 4ac)}}{2a},$

it must be remembered that the expression $\sqrt{(b^2 - 4ac)}$ is the square root of the compound quantity $b^2 - 4ac$, taken as a whole. We cannot simplify the solution unless we know the numerical values of a , b , c . It may sometimes happen that these values do not make $b^2 - 4ac$ a perfect square. In such a case the exact numerical solution of the equation cannot be determined.

Example 1. Solve $5x^2 - 15x + 11 = 0$.

$$\begin{aligned}\text{We have } x &= \frac{15 \pm \sqrt{(-15)^2 - 4 \cdot 5 \cdot 11}}{2 \cdot 5} \\ &= \frac{15 \pm \sqrt{5}}{10}.\end{aligned}$$

Now $\sqrt{5} = 2.236$ approximately.

$$\therefore x = \frac{15 \pm 2.236}{10} = 1.7236, \text{ or } 1.2764.$$

These solutions are correct only to four places of decimals, and neither of them will be found to *exactly* satisfy the equation.

Unless the *numerical* values of the unknown quantity are required it is usual to leave the roots in the form

$$\frac{15 + \sqrt{5}}{10}, \quad \frac{15 - \sqrt{5}}{10}.$$

Example 2. Solve $x^2 - 3x + 5 = 0$.

$$\begin{aligned}\text{We have } x &= \frac{3 \pm \sqrt{(-3)^2 - 4 \cdot 1 \cdot 5}}{2} \\ &= \frac{3 \pm \sqrt{9 - 20}}{2} \\ &= \frac{3 \pm \sqrt{-11}}{2}.\end{aligned}$$

But -11 has no square root exact or approximate [Art. 117]; so that no real value of x can be found to satisfy the equation. In such a case the roots are said to be *imaginary* or *impossible*.

201. There is still one method of obtaining the solution of a quadratic which will sometimes be found shorter than either of the methods already given.

Consider the equation $x^2 + \frac{7}{3}x = 2$.

Clearing of fractions, $3x^2 + 7x - 6 = 0$ (1);
by resolving the left-hand side into factors we have

$$(3x - 2)(x + 3) = 0.$$

Now if *either* of the factors $3x - 2$, $x + 3$ be zero their product is zero. Hence the quadratic equation is satisfied by either of the suppositions

$$3x - 2 = 0, \text{ or } x + 3 = 0.$$

Thus the roots are

$$\frac{2}{3}, \quad -3.$$

It appears from this that when a quadratic equation has been simplified and brought to the form of equation (1), its solution can always be readily obtained if the expression on the left-hand side can be resolved into factors. Each of these factors equated to zero gives a simple equation, and a corresponding root of the quadratic.

Example 1. Solve $2x^2 - ax + 2bx = ab$.

Transposing, so as to have all the terms on one side of the equation, we have

$$2x^2 - ax + 2bx - ab = 0.$$

$$\begin{aligned}\text{Now} \quad 2x^2 - ax + 2bx - ab &= x(2x - a) + b(2x - a) \\ &= (2x - a)(x + b).\end{aligned}$$

$$\text{Therefore} \quad (2x - a)(x + b) = 0;$$

$$\text{whence} \quad 2x - a = 0, \text{ or } x + b = 0.$$

$$\therefore x = \frac{a}{2}, \text{ or } -b.$$

Example 2. Solve $2(x^2 - 6) = 3(x - 4)$.

We have $2x^2 - 12 = 3x - 12$;

$$\text{that is,} \quad 2x^2 = 3x \dots\dots\dots(1).$$

Transposing, $2x^2 - 3x = 0$,

$$x(2x - 3) = 0,$$

$$\therefore x = 0, \text{ or } 2x - 3 = 0.$$

Thus the roots are $0, \frac{3}{2}$.

Note. In equation (1) above we might have divided both sides by x and obtained the simple equation $2x = 3$, whence $x = \frac{3}{2}$, which is one of the solutions of the given equation. But the student must be particularly careful to notice that whenever an x is removed by division from every term of an equation it must not be neglected, since the equation is satisfied by $x = 0$, which is therefore one of the roots.

202. There are some equations which are not really quadratics, but which may be solved by the methods explained in this chapter.

Example 1. Solve $x^4 - 13x^2 + 36 = 0$.

By resolution into factors, $(x^2 - 9)(x^2 - 4) = 0$;

$$\therefore x^2 - 9 = 0, \text{ or } x^2 - 4 = 0;$$

that is,

$$x^2 = 9, \text{ or } 4,$$

and

$$x = \pm 3, \text{ or } \pm 2.$$

Example

Write

or

From

Thus

$$x = -5,$$

Solve

$$1. \quad 3x^2 =$$

$$4. \quad x^2 =$$

$$7. \quad 8x^2 =$$

$$10. \quad 3x^2 =$$

Solve

$$13. \quad 6x^2 =$$

$$15. \quad 26x =$$

$$17. \quad 4x^2 =$$

$$19. \quad 7x^2 =$$

$$21. \quad 25x^2 =$$

$$23. \quad 12x^2 =$$

$$25. \quad 35x^2 =$$

$$27. \quad x^2 =$$

$$29. \quad 3x^2 =$$

Solve

$$31. \quad 4 = 5$$

$$33. \quad x^6 +$$

$$35. \quad 16(a$$

$$37. \quad x^2(19$$

$$39. \quad x^2 -$$

Example 2. Solve $x^2 + 3x - \frac{20}{x^2 + 3x} = 8$.

Write y for $x^2 + 3x$, then we have

$$y - \frac{20}{y} = 8,$$

or

$$y^2 - 8y - 20 = 0.$$

From this quadratic

$$y = 10, \text{ or } -2;$$

$$\therefore x^2 + 3x = 10, \text{ or } -2.$$

Thus we have *two* quadratics to solve, and finally we obtain $x = -5, 2$; or $-1, -2$.

EXAMPLES XXV. c.

Solve by the aid of the formula in Art. 198 :

- | | | |
|-----------------------|---------------------------|----------------------------|
| 1. $3x^2 = 15 - 4x$. | 2. $2x^2 + 7x = 15$. | 3. $2x^2 + 7 - 9x = 0$. |
| 4. $x^2 = 3x + 5$. | 5. $5x^2 + 4 + 21x = 0$. | 6. $x^2 + 11 = 7x$. |
| 7. $8x^2 = x + 7$. | 8. $5x^2 = 17x - 10$. | 9. $35 + 9x - 2x^2 = 0$. |
| 10. $3x^2 = x + 1$. | 11. $3x^2 + 5x = 2$. | 12. $2x^2 + 5x - 33 = 0$. |

Solve by resolution into factors :

- | | |
|----------------------------------|----------------------------------|
| 13. $6x^2 = 7 + x$. | 14. $21 + 8x^2 = 26x$ |
| 15. $26x - 21 + 11x^2 = 0$. | 16. $5x^2 + 26x + 24 = 0$. |
| 17. $4x^2 = \frac{4}{15}x + 3$. | 18. $x^2 - 2 = \frac{23}{12}x$. |
| 19. $7x^2 = 28 - 96x$. | 20. $96x^2 = 4x + 15$. |
| 21. $25x^2 = 5x + 6$. | 22. $35 - 4x = 4x^2$. |
| 23. $12x^2 - 11ax = 36a^2$. | 24. $12x^2 + 36a^2 = 43ax$. |
| 25. $35b^2 = 9x^2 + 6bx$. | 26. $36x^2 - 35b^2 = 12bx$. |
| 27. $x^2 - 2ax + 4ab = 2bx$. | 28. $x^2 - 2ax + 8x = 16a$. |
| 29. $3x^2 - 2ax - bx = 0$. | 30. $ax^2 + 2x = bx$. |

Solve as explained in Art. 202 :

- | | |
|--|---|
| 31. $4 = 5x^2 - x^4$. | 32. $x^4 + 36 = 13x^2$. |
| 33. $x^3 + 7x^2 = 8$. | 34. $x^3 - 19x^2 = 216$. |
| 35. $16\left(x^2 + \frac{1}{x^2}\right) = 257$. | 36. $x^2 + \frac{a^2b^2}{x^2} = a^2 + b^2$. |
| 37. $x^3(19 + x^3) = 216$. | 38. $(x^2 + 2)^2 + 198 = 29(x^2 + 2)$. |
| 39. $x^2 - x + \frac{72}{x^2 - x} = 18$. | 40. $x(x - 2a) = \frac{8a^4}{x^2 - 2ax} + 7a^2$. |

CHAPTER XXVI.

SIMULTANEOUS QUADRATIC EQUATIONS.

203. WE shall now consider some of the most useful methods of solving simultaneous equations, one or more of which may be of a degree higher than the first ; but no fixed rules can be laid down which are applicable to all cases.

Example 1. Solve $x + y = 15$ (1),
 $xy = 36$ (2).

From (1) by squaring, $x^2 + 2xy + y^2 = 225$;
 from (2) $4xy = 144$;
 by subtraction, $x^2 - 2xy + y^2 = 81$;
 by taking the square root, $x - y = \pm 9$.

Combining this with (1) we have to consider the two cases,

$$\begin{array}{l} x + y = 15, \} \\ x - y = 9. \} \end{array} \quad \begin{array}{l} x + y = 15, \} \\ x - y = -9. \} \end{array}$$

from which we find $\begin{array}{l} x = 12, \} \\ y = 3. \} \end{array} \quad \begin{array}{l} x = 3, \} \\ y = 12. \} \end{array}$

Example 2. Solve $x - y = 12$ (1),
 $xy = 85$ (2).

From (1) $x^2 - 2xy + y^2 = 144$;
 from (2) $4xy = 340$;
 by addition, $x^2 + 2xy + y^2 = 484$;
 by taking the square root, $x + y = \pm 22$.

Combining this with (1) we have the two cases,

$$\begin{array}{l} x + y = 22, \} \\ x - y = 12. \} \end{array} \quad \begin{array}{l} x + y = -22, \} \\ x - y = 12. \} \end{array}$$

Whence $\begin{array}{l} x = 17, \} \\ y = 5. \} \end{array} \quad \begin{array}{l} x = -5, \} \\ y = -17. \} \end{array}$

CHAP. X

204.
 speciall
 other c
 As a
 metrical
 foregoi
 soon as
 either t

Exam

Multi

When

We ha

$x + y$

$x - y$

From
 and the

Exam

By sul

Equat
 Example

Solve

1. $x + y$
 xy

4. $x - y$
 xy

7. xy
 $x + y$

204. These are the simplest cases that arise, but they are specially important since the solution in a large number of other cases is dependent upon them.

As a rule our object is to solve the proposed equations *symmetrically*, by finding the values of $x+y$ and $x-y$. From the foregoing examples it will be seen that we can always do this as soon as we have obtained the product of the unknowns, and either their sum or their difference.

Example 1. Solve $x^2 + y^2 = 74$ (1),
 $xy = 35$ (2).

Multiply (2) by 2; then by addition and subtraction we have

$$x^2 + 2xy + y^2 = 144,$$

$$x^2 - 2xy + y^2 = 4;$$

Whence $x + y = \pm 12,$
 $x - y = \pm 2.$

We have now four cases to consider; namely,

$$\begin{array}{llll} x+y=12, \} & x+y=12, \} & x+y=-12, \} & x+y=-12, \} \\ x-y=2, \} & x-y=-2, \} & x-y=2, \} & x-y=-2, \} \end{array}$$

From which the values of x are 7, 5, -5, -7;
 and the corresponding values of y are 5, 7, -7, -5.

Example 2. Solve $x^2 + y^2 = 185$ (1),
 $x + y = 17$ (2).

By subtracting (1) from the square of (2) we have

$$2xy = 104;$$

$$\therefore xy = 52 \text{(3).}$$

Equations (1) and (3) can now be solved by the method of Example 1; and the solution is

$$\begin{array}{l} x=13, \text{ or } 4, \\ y=4, \text{ or } 13. \end{array}$$

EXAMPLES XXVI. a.

Solve the following equations:

1. $x+y=28,$
 $xy=187.$

2. $x+y=51,$
 $xy=518.$

3. $x+y=74,$
 $xy=1113.$

4. $x-y=5,$
 $xy=126.$

5. $x-y=8,$
 $xy=513.$

6. $xy=1075,$
 $x-y=18.$

7. $xy=923,$
 $x+y=84.$

8. $x-y=-8,$
 $xy=1353.$

9. $x-y=-22,$
 $xy=3848.$

Solve the following equations :

- | | | |
|---|--|---|
| 10. $xy = -2193,$
$x + y = -8.$ | 11. $x - y = -18,$
$xy = 1363.$ | 12. $xy = -1914,$
$x + y = -65.$ |
| 13. $x^2 + y^2 = 89,$
$xy = 40.$ | 14. $x^2 + y^2 = 170,$
$xy = 13.$ | 15. $x^2 + y^2 = 65,$
$xy = 28.$ |
| 16. $x^2 + y^2 = 178,$
$x + y = 16.$ | 17. $x + y = 15,$
$x^2 + y^2 = 125.$ | 18. $x - y = 4,$
$x^2 + y^2 = 106.$ |
| 19. $x^2 + y^2 = 180,$
$x - y = 6.$ | 20. $x^2 + y^2 = 185,$
$x - y = 3.$ | 21. $x + y = 13,$
$x^2 + y^2 = 97.$ |
| 22. $x + y = 9,$
$x^2 + xy + y^2 = 61.$ | 23. $x - y = 3,$
$x^2 - 3xy + y^2 = -19.$ | 24. $x^2 - xy + y^2 = 76,$
$x + y = 14.$ |
| 25. $\frac{1}{10}(x - y) = 1,$
$x^2 - 4xy + y^2 = 52.$ | 26. $\frac{1}{x} + \frac{1}{y} = 2,$
$x + y = 2.$ | 27. $\frac{1}{x} + \frac{1}{y} = \frac{7}{12},$
$xy = 12,$ |
| 28. $ax + by = 2,$
$abxy = 1.$ | 29. $x^2 + pxy + y^2 = p + 2,$
$qx^2 + xy + qy^2 = 2q + 1.$ | |

205. Any pair of equations of the form

$$x^2 \pm pxy + y^2 = a^2 \dots\dots\dots(1),$$

$$x \pm y = b \dots\dots\dots(2),$$

where p is any numerical quantity, can be reduced to one of the cases already considered; for by squaring (2) and combining with (1), an equation to find xy is obtained; the solution can then be completed by the aid of equation (2).

Example 1. Solve $x^3 - y^3 = 999 \dots\dots\dots(1),$

$$x - y = 3 \dots\dots\dots(2),$$

By division, $x^2 + xy + y^2 = 333 \dots\dots\dots(3);$

from (2) $x^2 - 2xy + y^2 = 9;$

by subtraction, $3xy = 324,$
 $xy = 108 \dots\dots\dots(4).$

From (2) and (4)
$$\begin{aligned} x &= 12, \text{ or } -9, \\ y &= 9, \text{ or } -12. \end{aligned}$$

Example 2. Solve $x^4 + x^2y^2 + y^4 = 2613 \dots\dots\dots(1),$

$$x^2 + xy + y^2 = 67 \dots\dots\dots(2).$$

Dividing (1) by (2) $x^2 - xy + y^2 = 39 \dots\dots\dots(3).$

From (2) and (3) by addition, $x^2 + y^2 = 53;$

by subtraction, $xy = 14;$

whence
$$\begin{aligned} x &= \pm 7, \pm 2, \\ y &= \pm 2, \pm 7. \end{aligned}$$

[Art. 204, Ex. 1.]

Exam.

From

by subtr

adding t

Comb

Solve

1. $x^3 +$
 $x +$

4. $x^3 +$
 $x -$

7. $x^4 +$
 $x^2 +$

9. $x^4 +$
 $x^2 +$

11. $\frac{1}{x^2} +$
 $\frac{1}{x} +$

14. $\frac{y}{x} +$
 $\frac{y}{x} +$

17. $4(x$

Example 3. Solve

$$\frac{1}{x} - \frac{1}{y} = \frac{1}{3} \dots\dots\dots(1),$$

$$\frac{1}{x^2} + \frac{1}{y^2} = \frac{5}{9} \dots\dots\dots(2).$$

From (1) by squaring, $\frac{1}{x^2} - \frac{2}{xy} + \frac{1}{y^2} = \frac{1}{9};$

by subtraction, $\frac{2}{xy} = \frac{4}{9};$

adding to (2), $\frac{1}{x^2} + \frac{2}{xy} + \frac{1}{y^2} = 1;$

$$\therefore \frac{1}{x} + \frac{1}{y} = \pm 1.$$

Combining with (1), $\frac{1}{x} = \frac{2}{3},$ or $-\frac{1}{3},$

$$\frac{1}{y} = \frac{1}{3},$$
 or $-\frac{2}{3};$

$$\therefore x = \frac{3}{2},$$
 or $-3,$ and $y = 3,$ or $-\frac{3}{2}.$

EXAMPLES XXVI. b.

Solve the equations :

1. $x^2 + y^3 = 407,$
 $x + y = 11.$

2. $x^3 + y^3 = 637,$
 $x + y = 13.$

3. $x + y = 23,$
 $x^3 + y^3 = 3473.$

4. $x^3 - y^3 = 218,$
 $x - y = 2.$

5. $x - y = 4,$
 $x^3 - y^3 = 988.$

6. $x^3 - y^3 = 2197,$
 $x - y = 13.$

7. $x^4 + x^2y^2 + y^4 = 2128,$
 $x^2 + xy + y^2 = 76.$

8. $x^4 + x^2y^2 + y^4 = 2923,$
 $x^2 - xy + y^2 = 37.$

9. $x^4 + x^2y^2 + y^4 = 9211,$
 $x^2 - xy + y^2 = 61.$

10. $x^4 + x^2y^2 + y^4 = 7371,$
 $x^2 - xy + y^2 = 63.$

11. $\frac{1}{x^3} + \frac{1}{y^3} = \frac{481}{576},$
 $\frac{1}{x} + \frac{1}{y} = \frac{29}{24},$

12. $\frac{1}{x^3} + \frac{1}{y^3} = \frac{61}{900},$
 $xy = 30.$

13. $\frac{x}{y} + \frac{y}{x} = 2\frac{1}{2},$
 $x + y = 6.$

14. $\frac{x}{y} + \frac{y}{x} = 2\frac{6}{11},$
 $x - y = 4.$

15. $\frac{34}{x^2 + y^2} = \frac{15}{xy},$
 $x + y = 8.$

16. $x^3 - y^3 = 56,$
 $x^2 + xy + y^2 = 28.$

17. $4(x^2 + y^2) = 17xy,$
 $x - y = 6.$

18. $x^3 + y^3 = 126,$
 $x^2 - xy + y^2 = 21.$

Solve the equations :

$$19. \frac{1}{x^3} + \frac{1}{y^3} = 1\frac{1}{125},$$

$$20. \frac{1}{x^3} - \frac{1}{y^3} = 91,$$

$$\frac{1}{x} + \frac{1}{y} = 1\frac{1}{5}.$$

$$\frac{1}{x} - \frac{1}{y} = 1.$$

206. The following method of solution may always be used when the equations are of the same degree and homogeneous.

[See Art. 24.]

Example. Solve $x^2 + xy + 2y^2 = 74$ (1),

$$2x^2 + 2xy + y^2 = 73$$
(2).

Put $y = mx$, and substitute in both equations. Thus

$$x^2(1 + m + 2m^2) = 74$$
(3).

and

$$x^2(2 + 2m + m^2) = 73$$
(4).

By division,
$$\frac{1 + m + 2m^2}{2 + 2m + m^2} = \frac{74}{73};$$

$$\therefore 73 + 73m + 146m^2 = 148 + 148m + 74m^2;$$

$$\therefore 72m^2 - 75m - 75 = 0,$$

or

$$24m^2 - 25m - 25 = 0;$$

$$\therefore (8m + 5)(3m - 5) = 0;$$

$$\therefore m = -\frac{5}{8}, \text{ or } \frac{5}{3}.$$

(i) Take $m = -\frac{5}{8}$, and substitute in either (3) or (4).

From (3)
$$x^2\left(1 - \frac{5}{8} + \frac{50}{64}\right) = 74;$$

$$\therefore x^2 = \frac{64 \times 74}{74} = 64;$$

$$\therefore x = \pm 8;$$

$$\therefore y = mx = -\frac{5}{8}x = \mp 5.$$

(ii) Take $m = \frac{5}{3}$; then from (3)

$$x^2\left(1 + \frac{5}{3} + \frac{50}{9}\right) = 74.$$

$$x^2 = \frac{74 \times 9}{74} = 9;$$

$$\therefore x = \pm 3;$$

$$\therefore y = mx = \frac{5}{3}x = \pm 5.$$

207.
the other
find the
substitu

Examp

From (

and subst

and by su

208.
general e
cases spe

Examp

From (1)
that is,
or
whence

(i) Cor

whence

and by su

(ii) Co

whence

207. When one of the equations is of the first degree and the other of a higher degree, we may from the simple equation find the value of one of the unknowns in terms of the other, and substitute in the second equation.

Example. Solve $3x - 4y = 5$ (1),
 $3x^2 - xy - 3y^2 = 21$ (2).

From (1) we have $x = \frac{5+4y}{3}$;

and substituting in (2), $\frac{3(5+4y)^2}{9} - \frac{y(5+4y)}{3} - 3y^2 = 21$;

$\therefore 75 + 120y + 48y^2 - 15y - 12y^2 - 27y^2 = 189$;

$9y^2 + 105y - 114 = 0$,

$3y^2 + 35y - 38 = 0$;

$\therefore (y-1)(3y+38) = 0$;

$\therefore y = 1$, or $-\frac{38}{3}$;

and by substituting in (1), $x = 3$, or $-\frac{137}{9}$;

208. The examples we have given will be sufficient as a general explanation of the methods to be employed; but in some cases special artifices are necessary.

Example 1. Solve $x^2 + 4xy + 3x = 40 - 6y - 4y^2$ (1),
 $2xy - x^2 = 3$ (2).

From (1) we have $x^2 + 4xy + 4y^2 + 3x + 6y = 40$;

that is, $(x+2y)^2 + 3(x+2y) - 40 = 0$,

or $(x+2y+8)(x+2y-5) = 0$;

whence $x+2y = -8$, or 5 .

(i) Combining $x+2y=5$ with (2) we obtain

$2x^2 - 5x + 3 = 0$;

whence $x = 1$, or $\frac{3}{2}$;

and by substituting in $x+2y=5$, $y = 2$, or $\frac{7}{4}$.

(ii) Combining $x+2y = -8$ with (2) we obtain

$2x^2 + 8x + 3 = 0$;

whence $x = \frac{-4 \pm \sqrt{10}}{2}$; and $y = \frac{-12 \mp \sqrt{10}}{4}$.

Example 2. Solve $x^2y^2 - 6x = 34 - 3y$(1),

$$3xy + y = 2(9 + x) \text{(2).}$$

From (1) $x^2y^2 - 6x + 3y = 34$;

from (2) $9xy - 6x + 3y = 54$;

by subtraction, $x^2y^2 - 9xy + 20 = 0$,

$$(xy - 5)(xy - 4) = 0 ;$$

$$\therefore xy = 5, \text{ or } 4.$$

(i) Substituting $xy = 5$ in (2) gives $y - 2x = 3$.

From these equations we obtain $x = 1$, or $-\frac{5}{2}$,
 $y = 5$, or -2 .

(ii) Substituting $xy = 4$ in (2) gives $y - 2x = 6$.

From these equations we obtain $x = \frac{-3 \pm \sqrt{17}}{2}$,
 and $y = 3 \pm \sqrt{17}$.

EXAMPLES XXVI. c.

Solve the equations :

- | | | |
|--|--|---|
| 1. $5x - y = 17$,
$xy = 12$. | 2. $x^2 + xy = 15$,
$y^2 + xy = 10$. | 3. $x - y = 10$,
$x^2 - 2xy - 3y^2 = 84$. |
| 4. $3x + 2y = 16$,
$xy = 10$. | 5. $3x - y = 11$,
$3x^2 - y^2 = 47$. | 6. $x - 3y = 1$,
$x^2 - 2xy + 9y^2 = 17$. |
| 7. $x + 2y = 9$,
$3y^2 - 5x^2 = 43$. | 8. $x^2 + y^2 = 5$,
$2xy - y^2 = 3$. | 9. $5x + y = 3$,
$2x^2 - 3xy - y^2 = 1$. |
| 10. $3x^2 - 5y^2 = 28$,
$3xy - 4y^2 = 8$. | 11. $3x^2 - y^2 = 23$,
$2x^2 - xy = 12$. | 12. $x^2 + xy + y^2 = 3\frac{1}{4}$,
$2x^2 - 3xy + 2y^2 = 2\frac{3}{4}$. |
| 13. $x^2 - 3xy + y^2 + 1 = 0$,
$3x^2 - xy + 3y^2 = 13$. | 14. $7xy - 8x^2 = 10$,
$8y^2 - 9xy = 18$. | 15. $x^2 - 2xy = 21$,
$xy + y^2 = 18$. |
| 16. $x^2 + 3xy = 54$,
$xy + 4y^2 = 115$. | 17. $x^2 + y^2 = 152$,
$x^2y + xy^2 = 120$. | 18. $x^3 - y^3 = 127$,
$x^2y - xy^2 = 42$. |
| 19. $x^3 - y^3 = 208$,
$xy(x - y) = 48$. | 20. $x^2y^2 + 5xy = 84$,
$x + y = 8$. | 21. $x^2 + 4y^2 + 80 = 15x + 30y$,
$xy = 6$. |
| 22. $9x^2 + y^2 - 63x - 21y + 128 = 0$,
$xy = 4$. | | |

PR

209.

to quadra

Example
rate had b
two hours

Suppose
time occu

On the

whence
or

Hence t
being inad

It will f
question le
we are disc
a suitable
present cas

Since th
write $-x$

will be sati
signs throu

and this is
A train t
5 miles an
find the rat

E.A.

.....(1),
(2).

CHAPTER XXVII.

PROBLEMS LEADING TO QUADRATIC EQUATIONS.

209. WE shall now discuss some problems which give rise to quadratic equations.

Example 1. A train travels 300 miles at a uniform rate; if the rate had been 5 miles an hour more, the journey would have taken two hours less: find the rate of the train.

Suppose the train travels at the rate of x miles per hour, then the time occupied is $\frac{300}{x}$ hours.

On the other supposition the time is $\frac{300}{x+5}$ hours;

$$\therefore \frac{300}{x+5} = \frac{300}{x} - 2 \dots\dots\dots(1);$$

whence

$$x^2 + 5x - 750 = 0,$$

or

$$(x+30)(x-25) = 0,$$

$$\therefore x = 25, \text{ or } -30.$$

Hence the train travels 25 miles per hour, the negative value being inadmissible.

It will frequently happen that the algebraical statement of the question leads to a result which does not apply to the actual problem we are discussing. But such results can sometimes be explained by a suitable modification of the conditions of the question. In the present case we may explain the negative solution as follows.

Since the values $x=25$ and -30 satisfy the equation (1), if we write $-x$ for x the resulting equation,

$$\frac{300}{-x+5} = \frac{300}{-x} - 2 \dots\dots\dots(2),$$

will be satisfied by the values $x = -25$ and 30 . Now, by changing signs throughout, equation (2) becomes $\frac{300}{x-5} = \frac{300}{x} + 2$; and this is the algebraical statement of the following question:

A train travels 300 miles at a uniform rate; if the rate had been 5 miles an hour *less*, the journey would have taken two hours *more*: find the rate of the train. The rate is 30 miles an hour.

E.A.

Example 2. A person selling a horse for £72 finds that his loss per cent. is one-eighth of the number of pounds that he paid for the horse: what was the cost price?

Suppose that the cost price of the horse is x pounds; then the loss on £100 is $\frac{x}{8}$.

Hence the loss on $\pounds x$ is $x \times \frac{x}{800}$, or $\frac{x^2}{800}$ pounds;

$\therefore$ the selling price is $x - \frac{x^2}{800}$ pounds.

Hence $x - \frac{x^2}{800} = 72$,

or $x^2 - 800x + 57600 = 0$;

that is, $(x - 80)(x - 720) = 0$;

$\therefore x = 80$, or 720 ;

and each of these values will be found to satisfy the conditions of the problem. Thus the cost is either £80, or £720.

Example 3. A cistern can be filled by two pipes in $33\frac{1}{3}$ minutes; if the larger pipe takes 15 minutes less than the smaller to fill the cistern, find in what time it will be filled by each pipe singly.

Suppose that the two pipes running singly would fill the cistern in x and $x - 15$ minutes. When running together they will fill $\left(\frac{1}{x} + \frac{1}{x-15}\right)$ of the cistern in one minute. But they fill $\frac{1}{33\frac{1}{3}}$, or $\frac{3}{100}$ of the cistern in one minute.

Hence $\frac{1}{x} + \frac{1}{x-15} = \frac{3}{100}$,

$$100(2x - 15) = 3x(x - 15),$$

$$3x^2 - 245x + 1500 = 0,$$

$$(x - 75)(3x - 20) = 0;$$

$\therefore x = 75$, or $6\frac{2}{3}$.

Thus the smaller pipe takes 75 minutes, the larger 60 minutes.

The other solution $6\frac{2}{3}$ is inadmissible.

Example 4. The small wheel of a bicycle (1885) makes 135 revolutions more than the large wheel in a distance of 260 yards; if the circumference of each were one foot more, the small wheel would make 27 revolutions more than the large wheel in a distance of 70 yards: find the circumference of each wheel.

Suppo
in circum

In a c
revolutio

Hence

or

Similar

or

From (

whence

Substit

Putting
missible;
circumfer

Exampl
rowing on
performs
hours. H
 $5\frac{2}{3}$ hours:
the stream

Suppose
hour, and

With th
current y -

Suppose the small wheel to be x feet, and the large wheel y feet in circumference.

In a distance of 260 yards the two wheels make $\frac{780}{x}$ and $\frac{780}{y}$ revolutions respectively.

$$\text{Hence} \quad \frac{780}{x} - \frac{780}{y} = 135,$$

$$\text{or} \quad \frac{1}{x} - \frac{1}{y} = \frac{9}{52} \dots\dots\dots(1).$$

Similarly from the second condition, we obtain

$$\frac{210}{x+1} - \frac{210}{y+1} = 27,$$

$$\text{or} \quad \frac{1}{x+1} - \frac{1}{y+1} = \frac{9}{70} \dots\dots\dots(2).$$

$$\text{From (1)} \quad x = \frac{52y}{52+9y};$$

$$\text{whence} \quad x+1 = \frac{61y+52}{9y+52}.$$

$$\text{Substituting in (2), } \frac{9y+52}{61y+52} - \frac{1}{y+1} = \frac{9}{70},$$

$$70 \times 9y^2 = 9(61y+52)(y+1),$$

$$9y^2 - 113y - 52 = 0,$$

$$(y-13)(9y+4) = 0;$$

$$\therefore y = 13, \text{ or } -\frac{4}{9}.$$

Putting $y=13$ we find that $x=4$. The other value of y is inadmissible; hence the small wheel is 4 feet, the large wheel 13 feet in circumference.

Example 5. On a river there are two towns 24 miles apart. By rowing one half of the distance and walking the other half, a man performs the journey down stream in 5 hours, and up stream in 7 hours. Had there been no current, each journey would have taken $5\frac{2}{3}$ hours: find the rate of his walking, and rowing, and the rate of the stream.

Suppose that the man walks x miles per hour, rows y miles per hour, and that the stream flows at the rate of z miles per hour.

With the current the man rows $y+z$ miles, and against the current $y-z$ miles per hour.

Hence we have the following equations :

$$\frac{12}{x} + \frac{12}{y+z} = 5 \dots\dots\dots (1),$$

$$\frac{12}{x} + \frac{12}{y-z} = 7 \dots\dots\dots (2),$$

$$\frac{12}{x} + \frac{12}{y} = 5\frac{2}{3} \dots\dots\dots (3).$$

From (1) and (3) by subtraction, $\frac{1}{y} - \frac{1}{y+z} = \frac{1}{18} \dots\dots\dots (4).$

Similarly, from (2) and (3) $\frac{1}{y-z} - \frac{1}{y} = \frac{1}{9} \dots\dots\dots (5).$

From (4) $18z = y(y+z) \dots\dots\dots (6);$

and from (5) $9z = y(y-z) \dots\dots\dots (7).$

From (6) and (7) by division, $2 = \frac{y+z}{y-z};$

whence $y = 3z;$

$\therefore$ from (4) $z = 1\frac{1}{2};$ and hence $y = 4\frac{1}{2}, x = 4.$

Thus the rates of walking and rowing are 4 miles and $4\frac{1}{2}$ miles per hour respectively; and the stream flows at the rate of $1\frac{1}{2}$ miles per hour.

EXAMPLES XXVII.

1. Find a number whose square diminished by 119 is equal to ten times the excess of the number over 8.

2. A man is five times as old as his son, and the sum of the squares of their ages is equal to 2106: find their ages.

3. The sum of the reciprocals of two consecutive numbers is $\frac{15}{56}$: find them.

4. Find a number which when increased by 17 is equal to 66 times the reciprocal of the number.

5. Find two numbers whose sum is 9 times their difference, and the difference of whose squares is 81.

6. The sum of a number and its square is nine times the next highest number: find it.

7. If a train travelled 5 miles an hour faster it would take one hour less to travel 210 miles: what time does it take?

8. sum is 1

9. area is 1

10. feet; and of the s

11. 22½ min less than

12. the jour at what

13. shilling the cost

14. 3 on his of a pen master g

15. lowers t

16. width ro width.

17. if each t find the

18. this is a wide. I feet, find

19. a dozen than she them at

20. The diff 4 yards:

21. the digit be multi number.

8. Find two numbers the sum of whose squares is 74, and whose sum is 12.

9. The perimeter of a rectangular field is 500 yards, and its area is 14400 square yards: find the length of the sides.

10. The perimeter of one square exceeds that of another by 100 feet; and the area of the larger square exceeds three times the area of the smaller by 325 square feet: find the length of their sides.

11. A cistern can be filled by two pipes running together in $22\frac{1}{2}$ minutes; the larger pipe would fill the cistern in 24 minutes less than the smaller one: find the time taken by each.

12. A man travels 108 miles, and finds that he could have made the journey in $4\frac{1}{2}$ hours less had he travelled 2 miles an hour faster: at what rate did he travel?

13. I buy a number of cricket balls for £5; had they cost a shilling apiece less, I should have had five more for the money: find the cost of each.

14. A boy was sent out for a shilling's worth of eggs. He broke 3 on his way home, and his master therefore had to pay at the rate of a penny more than the market price for 5. How many did the master get for a shilling?

15. What are eggs a dozen when two more in a shilling's worth lowers the price a penny per dozen?

16. A lawn 50 feet long and 34 feet broad has a path of uniform width round it; if the area of the path is 540 square feet, find its width.

17. A hall can be paved with 200 square tiles of a certain size; if each tile were one inch longer each way it would take 128 tiles: find the length of each tile.

18. In the centre of a square garden is a square lawn; outside this is a gravel walk 4 feet wide, and then a flower border 6 feet wide. If the flower border and lawn together contain 721 square feet, find the area of the lawn.

19. By lowering the price of apples and selling them one penny a dozen cheaper, an applewoman finds that she can sell 60 more than she used to do for 5s. At what price per dozen did she sell them at first?

20. Two rectangles contain the same area, 480 square yards. The difference of their lengths is 10 yards, and of their breadths 4 yards: find their sides.

21. There is a number between 10 and 100; when multiplied by the digit on the left the product is 280; if the sum of the digits be multiplied by the same digit the product is 55: required the number.

22. A farmer having sold at 75s. a head, a flock of sheep which cost him x shillings a head, finds that he has realised x per cent. profit on his outlay : find x .

23. A tradesman bought a number of yards of cloth for £5; he kept 5 yards and sold the rest at 2s. per yard more than he gave, and got £1 more than he originally spent : how many yards did he buy?

24. If a carriage wheel $14\frac{2}{3}$ ft. in circumference takes one second more to revolve, the rate of the carriage per hour will be $2\frac{2}{3}$ miles less : how fast is the carriage travelling?

25. A broker bought as many railway shares as cost him £1875; he reserved 15, and sold the remainder for £1740, gaining £4 a share on their cost price. How many shares did he buy?

26. A and B are two stations 300 miles apart. Two trains start simultaneously from A and B , each to the opposite station. The train from A reaches B nine hours, the train from B reaches A four hours after they meet : find the rate at which each train travels.

27. A train A starts to go from P to Q , two stations 240 miles apart, and travels uniformly. An hour later another train B starts from P , and after travelling for 2 hours, comes to a point that A had passed 45 minutes previously. The pace of B is now increased by 5 miles an hour, and it overtakes A just on entering Q . Find the rates at which they started.

28. A cask P is filled with 50 gallons of water, and a cask Q with 40 gallons of brandy; x gallons are drawn from each cask, mixed and replaced; and the same operation is repeated. Find x when there are $8\frac{7}{8}$ gallons of brandy in P after the second replacement.

29. Two farmers A and B have 30 cows between them; they sell at different prices, but each receives the same sum. If A had sold his at B 's price, he would have received £320; and if B had sold his at A 's price, he would have received £245. How many had each?

30. A man arrives at the railway station nearest to his house $1\frac{1}{2}$ hours before the time at which he had ordered his carriage to meet him. He sets out at once to walk at the rate of 4 miles an hour, and, meeting his carriage when it had travelled 8 miles, reaches home exactly 1 hour earlier than he had originally expected. How far is his house from the station, and at what rate was his carriage driven?

210.
resolving
chapter
greater

211.
being w
may the

Exam.

Exam.

212.
be sepa
the diff

Exam.

Exam.

sheep which
at per cent.

for £5; he
an he gave,
ards did he

one second
be $2\frac{2}{3}$ miles

him £1875;
£4 a share

trains start
ation. The
ches A four
travels.

s 240 miles
in B starts
oint that A
w increased
g Q . Find

ask Q with
ask, mixed
nd x when
acement.

n; they sell
 A had sold
had sold his
had each?

o his house
carriage to
4 miles an
d 8 miles,
y expected.
ate was his

CHAPTER XXVIII.

HARDER FACTORS.

210. In Chapter xvii. we have explained several rules for resolving algebraical expressions into factors; in the present chapter we shall continue the subject by discussing cases of greater difficulty.

211. By a slight modification some expressions admit of being written in the form of the difference of two squares, and may then be resolved into factors by the method of Art. 133.

Example 1. Resolve into factors $x^4 + x^2y^2 + y^4$.

$$\begin{aligned} x^4 + x^2y^2 + y^4 &= (x^4 + 2x^2y^2 + y^4) - x^2y^2 \\ &= (x^2 + y^2)^2 - (xy)^2 \\ &= (x^2 + y^2 + xy)(x^2 + y^2 - xy) \\ &= (x^2 + xy + y^2)(x^2 - xy + y^2). \end{aligned}$$

Example 2. Resolve into factors $x^4 - 15x^2y^2 + 9y^4$.

$$\begin{aligned} x^4 - 15x^2y^2 + 9y^4 &= (x^4 - 6x^2y^2 + 9y^4) - 9x^2y^2 \\ &= (x^2 - 3y^2)^2 - (3xy)^2 \\ &= (x^2 - 3y^2 + 3xy)(x^2 - 3y^2 - 3xy). \end{aligned}$$

212. Expressions which can be put into the form $x^3 \pm \frac{1}{y^3}$ may be separated into factors by the rules for resolving the sum or the difference of two cubes. [Art. 136.]

Example 1. $\frac{8}{a^3} - 27b^3 = \left(\frac{2}{a}\right)^3 - (3b)^3$

$$= \left(\frac{2}{a} - 3b\right) \left(\frac{4}{a^2} + \frac{6b^2}{a} + 9b^4\right).$$

Example 2. Resolve $a^2x^3 - \frac{8a^2}{y^3} - x^3 + \frac{8}{y^3}$ into four factors.

$$\begin{aligned} a^2x^3 - \frac{8a^2}{y^3} - x^3 + \frac{8}{y^3} &= x^3(a^2 - 1) - \frac{8}{y^3}(a^2 - 1) \\ &= (a^2 - 1) \left(x^3 - \frac{8}{y^3}\right) \\ &= (a + 1)(a - 1) \left(x - \frac{2}{y}\right) \left(x^2 + \frac{2x}{y} + \frac{4}{y^2}\right). \end{aligned}$$

Example 3. Resolve $a^9 - 64a^3 - a^6 + 64$ into six factors.

$$\begin{aligned}\text{The expression} &= a^3(a^6 - 64) - (a^6 - 64) \\ &= (a^6 - 64)(a^3 - 1) \\ &= (a^3 + 8)(a^3 - 8)(a^3 - 1) \\ &= (a + 2)(a^2 - 2a + 4)(a - 2)(a^2 + 2a + 4)(a - 1)(a^2 + a + 1).\end{aligned}$$

Example 4.

$$a(a-1)x^2 - (a-b-1)xy - b(b+1)y^2. \quad \{ax - (b+1)y\} \{(a-1)x + by\}.$$

Note. In examples of this kind the coefficients of x and y in the binomial factors can usually be guessed at once, and it only remains to verify the coefficient of the middle term.

213. From Example 2, Art. 53, we see that the quotient of

$$a^3 + b^3 + c^3 - 3abc \text{ by } a + b + c \text{ is } a^2 + b^2 + c^2 - bc - ca - ab.$$

$$\text{Thus } a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - bc - ca - ab) \dots (1).$$

This result is important and should be carefully remembered. We may note that the expression on the left consists of the sum of the cubes of three quantities a, b, c , diminished by 3 times the product abc . Whenever an expression admits of a similar arrangement, the above formula will enable us to resolve it into factors.

Example 1. Resolve into factors $a^3 - b^3 + c^3 + 3abc$.

$$\begin{aligned}a^3 - b^3 + c^3 + 3abc &= a^3 + (-b)^3 + c^3 - 3a(-b)c \\ &= (a - b + c)(a^2 + b^2 + c^2 + bc - ca + ab),\end{aligned}$$

$-b$ taking the place of b in formula (1).

Example 2.

$$\begin{aligned}x^3 - 8y^3 - 27 - 18xy &= x^3 + (-2y)^3 + (-3)^3 - 3x(-2y)(-3) \\ &= (x - 2y - 3)(x^2 + 4y^2 + 9 - 6y + 3x + 2xy).\end{aligned}$$

EXAMPLES XXVIII. a.

Resolve into factors :

- | | |
|------------------------------|-------------------------------|
| 1. $x^4 + 16x^2 + 256.$ | 2. $81a^4 + 9a^2b^2 + b^4.$ |
| 3. $x^4 + y^4 - 7x^2y^2.$ | 4. $m^4 + n^4 - 18m^2n^2.$ |
| 5. $x^4 - 6x^2y^2 + y^4.$ | 6. $4x^4 + 9y^4 - 93x^2y^2.$ |
| 7. $4m^4 + 9n^4 - 24m^2n^2.$ | 8. $9x^4 + 4y^4 + 11x^2y^2.$ |
| 9. $x^4 - 19x^2y^2 + 25y^4.$ | 10. $16a^4 + b^4 - 28a^2b^2.$ |

11. $\frac{27}{a^3b^3} - 1.$

12. $216a^3 - \frac{b^3}{8}.$

13. $\frac{x^3}{125} + y^3.$

14. $\frac{m^3n^3}{729} - 1.$

15. $\frac{a^3b^3}{125} + 1000.$

16. $\frac{x^3}{512} - \frac{64}{x^3}.$

Resolve into two or more factors :

17. $x^2y + 3xy^2 - 3x^3 - y^3.$

18. $4mn^2 - 20n^3 + 45nm^2 - 9m^3.$

19. $ab(x^2 + 1) + x(a^2 + b^2).$

20. $y^2z^2(x^4 - 1) + x^2(y^4 - z^4).$

21. $a^3 + (a + b)ax + bx^2.$

22. $pn(m^2 + 1) - m(p^2 + n^2).$

23. $6bx(a^2 + 1) - a(4x^2 + 9b^2).$

24. $(2a^2 + 3y^2)x + (2x^2 + 3a^2)y.$

25. $(2x^2 - 3a^2)y + (2a^2 - 3y^2)x.$

26. $a(a - 1)x^2 + (2a^2 - 1)x + a(a + 1).$

27. $3x^2 - (4a + 2b)x + a^2 + 2ab.$

28. $2a^2x^2 - 2(3b - 4c)(b - c)y^2 + abxy.$

29. $(a^2 - 3a + 2)x^2 + (2a^2 - 4a + 1)x + a(a - 1).$

30. $a(a + 1)x^2 + (a + b)xy - b(b - 1)y^2.$

31. $b^3 + c^3 - 1 + 3bc.$

32. $a^3 + 8c^3 + 1 - 6ac.$

33. $a^3 + b^3 + 8c^3 - 6abc.$

34. $a^3 - 27b^3 + c^3 + 9abc.$

35. $a^3 - b^3 - c^3 - 3abc.$

36. $8a^3 + 27b^3 + c^3 - 18abc.$

37. Resolve $x^8 + 81x^4 + 6561$ into three factors.

38. Resolve $(a^4 - 2a^2b^2 - b^4)^2 - 4a^4b^4$ into four factors.

39. Resolve $4(ab + cd)^2 - (a^2 + b^2 - c^2 - d^2)^2$ into four factors.

40. Resolve $x^8 - \frac{1}{256}$ into four factors.

41. Resolve $x^{16} - y^{16}$ into five factors.

42. Resolve $x^{18} - y^{18}$ into six factors.

Resolve into four factors :

43. $\frac{a^3}{x^2} - 8x - a^3 + 8x^3.$

44. $x^9 + x^3y^6 - 8x^6y^3 - 8y^9.$

45. $x^9 + x^6 + 64x^3 + 64.$

46. $4a - 9b + \frac{4b^3}{a^2} - \frac{9a^3}{b^2}.$

47. $\frac{xy^3}{72} - \frac{x^3y^5}{32} - \frac{1}{9x^2} + \frac{y^2}{4}.$

48. $x^6 - 25x^2 + 6\frac{1}{4} - \frac{1}{4}x^4.$

Resolve into five factors :

49. $x^7 + x^4 - 16x^3 - 16.$

50. $16x^7 - 81x^3 - 16x^4 + 81.$

214. The actual processes of multiplication and division can often be partially or wholly avoided by a skilful use of factors.

It should be observed that the formulæ which the student has seen exemplified in the preceding pages are just as useful in their converse as in their direct application. Thus the formula for resolving into factors the difference of two squares is equally useful as enabling us to write down at once the product of the sum and the difference of two quantities.

Example 1. Multiply $2a + 3b - c$ by $2a - 3b + c$.

These expressions may be arranged thus :

$$2a + (3b - c) \text{ and } 2a - (3b - c).$$

Hence the product = $\{2a + (3b - c)\} \{2a - (3b - c)\}$

$$= (2a)^2 - (3b - c)^2 \quad [\text{Art. 133.}]$$

$$= 4a^2 - (9b^2 - 6bc + c^2)$$

$$= 4a^2 - 9b^2 + 6bc - c^2.$$

Example 2. Multiply $(a^2 + a + 1)x - a - 1$ by $(a - 1)x - a^2 + a - 1$.

The product = $\{(a^2 + a + 1)x - (a + 1)\} \{(a - 1)x - (a^2 - a + 1)\}$

$$= (a^3 - 1)x^2 - \{(a^4 + a^2 + 1) + (a^3 - 1)\}x + (a^3 + 1)$$

$$= (a^3 - 1)x^2 - (a^4 + 2a^2)x + a^3 + 1$$

$$= (a^3 - 1)x^2 - a^2(a^2 + 2)x + a^3 + 1.$$

Note. The product of $a^2 + a + 1$ and $a^2 - a + 1$ is $a^4 + a^2 + 1$ and should be written down without actual multiplication.

Example 3. Multiply $(3 + x - 2x^2)^2 - (3 - x + 2x^2)^2 \dots \dots \dots (1),$
by $(3 + x + 2x^2)^2 - (3 - x - 2x^2)^2 \dots \dots \dots (2),$

The expression (1)

$$= (3 + x - 2x^2 + 3 - x + 2x^2)(3 + x - 2x^2 - 3 + x - 2x^2)$$

$$= 6(2x - 4x^2)$$

$$= 12x(1 - 2x).$$

The expression (2)

$$= (3 + x + 2x^2 + 3 - x - 2x^2)(3 + x + 2x^2 - 3 + x + 2x^2)$$

$$= 6(2x + 4x^2)$$

$$= 12x(1 + 2x).$$

Therefore the product = $12x(1 - 2x) \times 12x(1 + 2x)$

$$= 144x^2(1 - 4x^2).$$

division can
of factors.

the student
as useful in
the formula
es is equally
duct of the

Example 4. Divide the product of $2x^2+x-6$, and $6x^2-5x+1$ by $3x^2+5x-2$.

Denoting the division by means of a fraction, the required quotient

$$\begin{aligned} &= \frac{(2x^2+x-6)(6x^2-5x+1)}{3x^2+5x-2} \\ &= \frac{(2x-3)(x+2)(3x-1)(2x-1)}{(3x-1)(x+2)} \\ &= (2x-3)(2x-1). \end{aligned}$$

Example 5. Shew that $(2x+3y-z)^3+(3x+7y+z)^3$ is divisible by $5(x+2y)$.

The given expression is of the form A^3+B^3 , and therefore has a divisor of the form $A+B$.

[Art. 133.]

Therefore $(2x+3y-z)^3+(3x+7y+z)^3$
is divisible by $(2x+3y-z)+(3x+7y+z)$,
that is, by $5x+10y$,
or by $5(x+2y)$.

$-a^2+a-1$.
 $+1\}$
 $+1\}$

Example 6. Find the quotient when $a^3+8-5b(25b^2-6a)$ is divided by $a-5b+2$.

$+a^2+1$ and

$$\begin{aligned} \text{The expression} &= a^3+8-125b^3+30ab \\ &= a^3+(-5b)^3+(2)^3-3 \cdot a(-5b)(2) \\ &= (a-5b+2)(a^2+25b^2+4+10b-2a+5ab). \end{aligned}$$

[Art. 213.]

$\therefore$ the quotient is $a^2+25b^2+4+10b-2a+5ab$.

.....(1),
.....(2).

Example 7. If $x+y=a$, and $x-y=b$ shew that

x^2

$$\begin{aligned} 4(x^4-6x^2y^2+y^4) &= 6a^2b^2-a^4-b^4. \\ x^4-6x^2y^2+y^4 &= (x^4-2x^2y^2+y^4)-4x^2y^2 \end{aligned}$$

x^2

$$\begin{aligned} &= (x^2-y^2)^2-\frac{1}{4}(4xy)^2 \\ &= \{(x+y)(x-y)\}^2-\frac{1}{4}\{(x+y)^2-(x-y)^2\}^2 \\ &= (ab)^2-\frac{1}{4}(a^2-b^2)^2; \\ \therefore 4(x^4-6x^2y^2+y^4) &= 4a^2b^2-(a^2-b^2)^2 \\ &= 6a^2b^2-a^4-b^4. \end{aligned}$$

EXAMPLES XXVIII. b.

Find the product of

1. $2x - 7y + 3z$ and $2x + 7y - 3z$.
2. $3x^2 - 4xy + 7y^2$ and $3x^2 + 4xy + 7y^2$.
3. $5x^2 + 5xy - 9y^2$ and $5x^2 - 5xy - 9y^2$.
4. $7x^2 - 8xy + 3y^2$ and $7x^2 + 8xy - 3y^2$.
5. $x^3 + 2x^2y + 2xy^2 + y^3$ and $x^3 - 2x^2y + 2xy^2 - y^3$.
6. $(x + y)^2 + 2(x + y) + 4$ and $(x + y)^2 - 2(x + y) + 4$.
7. $(1 + x + 2x^2)^2 - (1 - x - 2x^2)^2$ and $(1 + x - 2x^2)^2 - (1 - x + 2x^2)^2$.
8. $(a^2 + 3a - 1)^2 - (a^2 - 3a - 1)^2$ and $(a^2 + a + 1)^2 - (a^2 - a + 1)^2$.
9. $x^3 - 4x^2 + 8x - 8$ and $x^3 + 4x^2 + 8x + 8$.
10. $x^3 - 6ax^2 + 18a^2x - 27a^3$ and $x^3 + 6ax^2 + 18a^2x + 27a^3$.
11. $x - a - \frac{x^2}{a} - \frac{a^2}{x}$ and $x + a + \frac{x^2}{a} - \frac{a^2}{x}$.
12. $(2x^2 + 3x + 1)^2 - (2x^2 - 3x - 1)^2$ and $(x^2 + 6x - 2)^2 - (x^2 - 6x + 2)^2$.

Find the continued product of

13. $x^2 + ax + a^2$, $x^2 - ax + a^2$, $x^4 - a^2x^2 + a^4$.
14. $1 - x + x^2$, $1 + x + x^2$, $1 - x^2 + x^4$, $1 - x^4 + x^8$.
15. $(a - x)^3$, $(a + x)^3$, $(a^2 + x^2)^3$.
16. $(1 - x)^2$, $(1 + x)^2$, $(1 + x^2)^2$, $(1 + x^4)^2$.
17. $x^2 + 4x + 3$, $x^2 + x - 2$, $x^2 - 5x + 6$.
18. $x^2 + 2x - 3$, $x^2 - 5x + 6$, $x^2 + 3x + 2$.
19. $x + 2$, $x^2 + 2x + 4$, $x - 2$, $x^2 - 2x + 4$.
20. Multiply the square of $a + 3b$ by $a^2 - 6ab + 9b^2$.
21. Multiply $\frac{1}{2}(a - b)^2 + \frac{1}{2}(b - c)^2 + \frac{1}{2}(c - a)^2$ by $a + b + c$.
22. Divide $(4x + 3y - 2z)^2 - (3x - 2y + 3z)^2$ by $x + 5y - 5z$.
23. Divide $x^8 + 16a^4x^4 + 256a^8$ by $x^2 + 2ax + 4a^2$.
24. Divide $(3x + 4y - 2z)^2 - (2x + 3y - 4z)^2$ by $x + y + 2z$.
25. Divide the product of $x^2 + 7x + 10$ and $x + 3$ by $x^2 + 5x + 6$.
26. Divide $2x(x^2 - 1)(x + 2)$ by $x^2 + x - 2$.
27. Divide $5x(x - 11)(x^2 - x - 156)$ by $x^3 + x^2 - 132x$.

28. Divide $x^6 + 19x^3 - 216$ by $(x^2 - 3x + 9)(x - 2)$.
29. Divide $(5x^2 - 3x - 6)^2 - (2x^2 - 7x + 9)^2$ by the product of $3x - 5$ and $x + 3$.
30. Divide $a^9 - b^9$ by the product of $a^2 + ab + b^2$ and $a^6 + a^3b^3 + b^6$.
31. Divide $(x^3 - 3x^2y)^2 - (3xy^2 - y^3)^2$ by $(x - y)^3$.
32. Divide $(x^2 - yz)^3 + 8y^3z^3$ by $x^2 + yz$.
33. Divide $18xy + 1 + 27x^2 - 8y^3$ by $1 + 3x - 2y$.
34. Divide $(2x^2 + 3x - 1)^2 - (x^2 + 4x + 5)^2$ by the product of $3x + 4$ and $x + 2$.
35. Divide the product of $6a^2 - 23a + 20$ and $22a^2 - 81a + 14$ by $33a^2 - 50a + 8$.
36. Divide the product of $x^2 + (a - b)x - ab$ and $x^2 - (a - b)x - ab$ by $x^2 + (a + b)x + ab$.
37. Divide $a^3 - 8y^3 - 9x(3x^2 + 2ay)$ by $a - 3x - 2y$.
38. Divide $27 - 8x^3 - 64y^3 - 72xy$ by $3 - 2(x + 2y)$.
39. Shew that $(2x - 3y + 1)^3 - (1 - 3x + 2y)^3$ is divisible by $5(x - y)$.
40. Shew that the square of $x + 1$ exactly divides $(x^3 + x^2 + 4)^3 - (x^3 - 2x + 3)^3$.
41. Shew that $2b + 2d$ is a factor of the expression $(a + b + c + d)^3 - (a - b + c - d)^3$.
42. Shew that $(3x^2 - 7x + 2)^3 - (x^2 - 8x + 8)^3$ is divisible by $2x - 3$ and by $x + 2$.
43. Shew that $(7x^2 + 3x - 3)^3 + (5x^2 - 4x - 3)^3$ is divisible by $4x - 3$ and by $3x + 2$.
44. Shew that the sum of the cubes of $2x^2 - 5x - 9$ and $x^2 + 6x - 5$ is divisible by the product of $3x + 7$ and $x - 2$.
45. If $x + y = m$ and $x - y = n$, express $x^3 + y^3$ in terms of m and n .
46. If $x + y = m$ and $x - y = n$, shew that $16(x^4 - 7x^2y^2 + y^4) = (5m^2 - n^2)(5n^2 - m^2)$.
47. Find the value of $x^4 + x^2y^2 + y^4$ when $x + y = 2a$, $x - y = 2b$.
48. If $x + y = 2a$ and $x - y = 2b$ prove that $x^4 - 23x^2y^2 + y^4 = (7a^2 - 3b^2)(7b^2 - 3a^2)$.
49. Find the value of $x^4 - 47x^2y^2 + y^4$ in terms of p and q when $x + y = p$ and $x - y = q$.
50. Find the value of $x^4 - 2x^3y + 2xy^3 - y^4$ when $x = a + b$ and $y = a - b$.

CHAPTER XXIX.

MISCELLANEOUS THEOREMS AND EXAMPLES.

215. Examples upon the simple rules, *e.g.* Division, Highest Common Factor, Evolution, etc., frequently occur which cannot be neatly and concisely worked without a ready use of factors and compound expressions. These we have hitherto excluded as unsuitable for the student until he has gained confidence and power by practice. We propose in the present chapter to bring together a miscellaneous collection of examples, for the most part not new in principle, but requiring some skill for their solution. The chapter will be found useful as a revision of the earlier chapters.

Example. Divide

$$\begin{array}{r} ax^4 - (ap - b)x^3 + (aq - bp - c)x^2 + (bq + cp)x - cq \text{ by } ax^2 + bx - c. \\ ax^2 + bx - c \overline{) ax^4 - (ap - b)x^3 + (aq - bp - c)x^2 + (bq + cp)x - cq} \quad x^2 - px + q \\ \underline{ax^4 + bx^3 - cx^2} \\ -apx^3 + (aq - bp)x^2 + (bq + cp)x \\ \underline{-apx^3 - bpx^2} \\ aqx^2 + bq - cq \\ \underline{aqx^2 + bq - cq} \\ 0 \end{array}$$

Note. When the coefficients in divisor or dividend are compound quantities it is best to retain them in brackets throughout the work.

216. In the process of finding the highest common factor, by the rules explained in Chap. XVIII., every remainder that occurs in the course of the work contains the factor we are seeking. Hence when any one of the remainders admits of being resolved into factors, we may often shorten the work.

Example 1. Find the H.C.F. of $2x^3 - (4a - 3c)x^2 + 6(b - ac)x + 9bc$ and $2x^3 + (2a + 3c)x^2 + (3ac - 4b)x - 6bc$.

$$\begin{array}{r} 2x^3 - (4a - 3c)x^2 + 6(b - ac)x + 9bc \overline{) 2x^3 + (2a + 3c)x^2 + (3ac - 4b)x - 6bc} \\ \underline{2x^3 - (4a - 3c)x^2 + (6b - 6ac)x + 9bc} \\ 6ax^2 + (9ac - 10b)x - 15bc \end{array}$$

$$\begin{aligned} \text{Now the remainder} &= 6ax^2 + 9acx - 10bx - 15bc \\ &= 3ax(2x + 3c) - 5b(2x + 3c) \\ &= (2x + 3c)(3ax - 5b). \end{aligned}$$

Of these factors, $3ax - 5b$ may clearly be rejected; therefore if there is a common factor it must be $2x + 3c$. And by division, or by the method explained in Art. 152, we find that $2x + 3c$ is a factor of each expression.

Hence the H.C.F. is $2x + 3c$.

Example 2. Find the H.C.F. of $(a^2 - 2a)x^2 + 2(2a - 1)x - a^2 + 1$ and $(a^2 - a - 2)x^2 + (4a + 1)x - a^2 - a$.

Each of these expressions can be resolved into factors as explained in Art. 212, Ex. 4. Thus

$$\begin{aligned}(a^2 - 2a)x^2 + 2(2a - 1)x - a^2 + 1 &= a(a - 2)x^2 + 2(2a - 1)x - (a + 1)(a - 1) \\ &= \{(a - 2)x + (a + 1)\}\{ax - (a - 1)\}. \\ (a^2 - a - 2)x^2 + (4a + 1)x - a^2 - a &= (a - 2)(a + 1)x^2 + (4a + 1)x - a(a + 1) \\ &= \{(a - 2)x + (a + 1)\}\{(a + 1)x - a\}.\end{aligned}$$

Hence the H.C.F. is $(a - 2)x + a + 1$.

EXAMPLES XXIX. a.

Divide

- $x^3 + (a + b + c)x^2 + (bc + ca + ab)x + abc$ by $x^2 + (a + b)x + ab$.
- $x^3 - (5 + a)x^2 + (4 + 5a + b)x^2 - (4a + 5b)x + 4b$ by $x^2 - 5x + 4$.
- $x^3 - (a - b)x^2 - (ab + 2b^2)x + 2ab^2$ by $x - b$.
- $x^3 - (p^2 + 3q^2)x + 2p^2q - 2q^3$ by $x + p + q$.
- $x^3 - (3mn + n^2)x + m(m^2 - n^2)$ by $x + m + n$.
- $a(a - 1)x^2 + (2a^2 - 1)x + a(a + 1)$ by $(a - 1)x + a$.
- $x^4 + (a + b)x^3 + (a^2 + ab + b^2)x^2 + (a^3 + b^3)x + a^2b^2$ by $x^2 + ax + b^2$.
- $2l^2x^2 - 2(3m - 4n)(m - n)y^2 + lmxxy$ by $lx + 2(m - n)y$.
- $(a^2 + a - 2)x^2 - (2a + 1)xy - (a^2 + a)y^2$ by $(a - 1)x - ay$.
- $x^3 - (a - b - 2)x^2 - (ab + 2a - 2b)x - 2ab$ by $(x - a)(x + 2)$.
- $(x + 1)^5 + 4(x + 1)^6 + 6(x + 1)^4 + 4(x + 1)^2 + 1$ by $x^2 + 2x + 2$.
- $(m + 1)(bx + an)b^2x^2 - (n + 1)(mbx + a)a^2$ by $bx - a$.

Find the H.C.F. of

- $(m^2 - 3m + 2)x^2 + (2m^2 - 4m + 1)x + m(m - 1)$ and $m(m - 1)x^2 + (2m^2 - 1)x + m(m + 1)$.
- $mpx^3 + (mq - np)x^2 - (mr + nq)x + nr$ and $max^3 - (mc + na)x^2 - (mb - nc)x + nb$.
- $2ap^3 + (3a - 2b)p^2q + (a - 3b)pq^2 - bq^3$ and $3ap^3 - (a + 3b)p^2q + (2a + b)pq^2 - 2bq^3$.
- $acx^3 + (bc + ad)x^2 + (bd + ac)x + bc$ and $2acx^3 + (2bc - ad)x^2 - (3ac + bd)x - 3bc$.

Find the H.C.F. of

17. $2a^2x^3 - (4b+3)ax^2 + 2(3b-ac)x + 3c$ and
 $2a^2x^3 + (2b-3)ax^2 - (4ac+3b)x + 6c$.
 18. $2ax^3 + (4a^2-1)bx^2 - (2ab^2+3c)x - 6abc$ and
 $ax^3 - (3-2a^2)bx^2 + (2c-6ab^2)x + 4abc$.

Find the L.C.M. of

19. $x^4 - px^3 + (q-1)x^2 + px - q$ and $x^4 - qx^3 + (p-1)x^2 + qx - p$.
 20. $p(p+1)x^2 + x - p(p-1)$ and $p(p+2)x^2 + 2x - p^2 + 1$.
 21. $(a^2 - 5a + 6)x^2 + 2(a-1)x - a(a+1)$ and
 $a(a-3)x^2 + 12x - (a+1)(a+4)$.

217. We add some miscellaneous questions in Evolution.

The *fourth* root of an expression is obtained by extracting the square root of the square root of the expression.

Similarly by successive applications of the rule for finding the square root, we may find the *eighth*, *sixteenth* ... root. The *sixth* root of an expression is found by taking the cube root of the square root, or the square root of the cube root.

Similarly by combining the two processes for extraction of cube and square roots, certain other higher roots may be obtained.

Example 1. Find the fourth root of
 $8x^4 - 216x^2y + 216x^2y^2 - 96xy^3 + 16y^4$.

Extracting the square root by the rule we obtain $9x^2 - 12xy + 4y^2$; and *by inspection*, the square root of this is $3x - 2y$, which is the required fourth root.

Example 2. Find the sixth root of
 $\left(x^3 - \frac{1}{x^3}\right)^2 - 6\left(x - \frac{1}{x}\right)\left(x^2 - \frac{1}{x^3}\right) + 9\left(x - \frac{1}{x}\right)^2$.

By inspection, the square root of this is

$$\left(x^3 - \frac{1}{x^3}\right) - 3\left(x - \frac{1}{x}\right),$$

which may be written $x^3 - 3x + \frac{3}{x} - \frac{1}{x^3}$;

and the cube root of this is $x - \frac{1}{x}$,

which is the required sixth root.

218. In Chap. vi. we have given examples of inexact division. In a similar manner when an expression is not an exact square

or cube
many

Exan

Thus

*219
arithme
cube ro
square
of figur
may be

*220
when th
method,

Let
root alr
common
the root
Then

Now
represen
E.A.

or cube, we may perform the process of evolution, and obtain as many terms of the root as we please.

Example. To find four terms of the square root of $1+2x-2x^2$.

$$\begin{array}{r}
 1 + 2x - 2x^2 \left(1 + x - \frac{3}{2}x^2 + \frac{3}{2}x^3 \right. \\
 \hline
 2 + x \quad \begin{array}{l} 2x - 2x^2 \\ 2x + x^2 \end{array} \\
 \hline
 2 + 2x - \frac{3}{2}x^2 \quad \begin{array}{l} - 3x^2 \\ - 3x^2 - 3x^3 + \frac{9}{4}x^4 \end{array} \\
 \hline
 2 + 2x - 3x^2 + \frac{3}{2}x^3 \quad \begin{array}{l} 3x^3 - \frac{9}{4}x^4 \\ 3x^3 + 3x^4 - \frac{9}{2}x^5 + \frac{9}{4}x^6 \\ - \frac{21}{4}x^4 + \frac{9}{2}x^5 - \frac{9}{4}x^6 \end{array}
 \end{array}$$

Thus the required result is $1 + x - \frac{3}{2}x^2 + \frac{3}{2}x^3$.

*219. In Art. 124 we pointed out the similarity between the arithmetical and algebraical methods of extracting square and cube roots. We shall now shew that in extracting either the square or the cube root of any number, when a certain number of figures have been obtained by the common rule, that number may be nearly doubled by ordinary division.

*220. *If the square root of a number consists of $2n+1$ figures, when the first $n+1$ of these have been obtained by the ordinary method, the remaining n may be obtained by division.*

Let N denote the given number; a the part of the square root already found, that is the first $n+1$ figures found by the common rule, with n ciphers annexed; x the remaining part of the root.

Then

$$\begin{aligned}
 \sqrt{N} &= a + x; \\
 \therefore N &= a^2 + 2ax + x^2; \\
 \therefore \frac{N - a^2}{2a} &= x + \frac{x^2}{2a} \dots \dots \dots (1).
 \end{aligned}$$

Now $N - a^2$ is the remainder after $n+1$ figures of the root, represented by a , have been found; and $2a$ is the divisor at the

E.A.

P

same stage of the work. We see from (1) that $N - a^2$ divided by $2a$ gives x , the rest of the quotient required, increased by $\frac{x^3}{2a}$. We shall shew that $\frac{x^3}{2a}$ is a *proper fraction*, so that by neglecting the remainder arising from the division, we obtain x , the rest of the root.

For x contains n figures, and therefore x^2 contains $2n$ figures at most; also a is a number of $2n+1$ figures (the last n of which are ciphers) and thus $2a$ contains $2n+1$ figures at least; and therefore $\frac{x^3}{2a}$ is a proper fraction.

From the above investigation, by putting $n=1$, we see that *two* at least of the figures of a square root must have been obtained in order that the method of division, which is employed to obtain the next figure of the square root, may give that figure correctly.

Example. Find the square root of 290 to five places of decimals.

$$\begin{array}{r}
 290 (17 \cdot 02 \\
 \underline{1} \\
 27 \overline{) 190} \\
 \underline{189} \\
 3402 \overline{) 10000} \\
 \underline{6804} \\
 3196
 \end{array}$$

Here we have obtained four figures in the square root by the ordinary method. Three more may be obtained by division only, using 2×1702 , that is 3404, for divisor, and 3196 as remainder. Thus

$$\begin{array}{r}
 3404 \overline{) 31960 (938} \\
 \underline{30636} \\
 13240 \\
 \underline{10212} \\
 30280 \\
 \underline{27232} \\
 3048
 \end{array}$$

And therefore to five places of decimals $\sqrt{290} = 17 \cdot 02938$.

When the divisor consists of several digits, the method of contracted division may be employed with advantage.

Again, it may be noticed that in obtaining the second figure of the root, the division of 190 by 20 gives 9 for the next figure; this is too great, and the figure 7 has to be obtained tentatively. This is one of the modifications of the algebraical rule to which we referred in Art. 124.

* 22
when the
method.

Let
already
rule, w

Then

Now
represent
same st
by $3a^2$
 $\frac{x^3}{a} + \frac{x^3}{3a^2}$
so that
we obtain

By sup

and

hence

and is th

Find th

1. $x^4 - 2$
2. $16 - \frac{1}{2}$
3. $a^4 + 8$
4. $1 + 4a$
5. $1 + 8a$

divided by
ased by $\frac{x^3}{2a}$
neglecting

the rest of

s $2n$ figures
t n of which
least; and

we see that
ve been ob-
employed to
that figure

of decimals.

root by the
ivision only,
inder. Thus

38.
thod of con-

ond figure of
figure; this is
ely. This is
we referred

*221. If the cube root of a number consists of $2n+2$ figures, when the first $n+2$ of these have been obtained by the ordinary method, the remaining n may be obtained by division.

Let N denote the given number; a the part of the cube root already found, that is the first $n+2$ figures found by the common rule, with n ciphers annexed; x the remaining part of the root.

Then

$$\sqrt[3]{N} = a + x;$$

$$\therefore N = a^3 + 3a^2x + 3ax^2 + x^3;$$

$$\therefore \frac{N - a^3}{3a^2} = x + \frac{x^2}{a} + \frac{x^3}{3a^2} \dots\dots\dots(1)$$

Now $N - a^3$ is the remainder after $n+2$ figures of the root, represented by a , have been found; and $3a^2$ is the divisor at the same stage of the work. We see from (1) that $N - a^3$ divided by $3a^2$ gives x , the rest of the quotient required, increased by $\frac{x^2}{a} + \frac{x^3}{3a^2}$. We shall shew that this expression is a *proper fraction*, so that by neglecting the remainder arising from the division, we obtain x , the rest of the root.

By supposition, x is $< 10^n$, and a is $> 10^{2n+1}$;

$$\therefore \frac{x^2}{a} \text{ is } < \frac{10^{2n}}{10^{2n+1}}; \text{ that is, } < \frac{1}{10};$$

$$\text{and } \frac{x^3}{3a^2} \text{ is } < \frac{10^{3n}}{3 \times 10^{4n+2}}; \text{ that is, } < \frac{1}{3 \times 10^{n+1}};$$

$$\text{hence } \frac{x^2}{a} + \frac{x^3}{3a^2} \text{ is } < \frac{1}{10} + \frac{1}{3 \times 10^{n+1}},$$

and is therefore a proper fraction.

EXAMPLES XXIX. b.

Find the fourth roots of the following expressions:

1. $x^4 - 28x^3 + 294x^2 - 1372x + 2401$.

2. $16 - \frac{32}{m} + \frac{24}{m^2} - \frac{8}{m^3} + \frac{1}{m^4}$.

3. $a^4 + 8a^3x + 16a^2x^2 + 32ax^3 + 24a^2x^2$.

4. $1 + 4x + 2x^2 - 8x^3 - 5x^4 + 8x^5 + 2x^6 - 4x^7 + x^8$.

5. $1 + 8x + 20x^2 + 8x^3 - 26x^4 - 8x^5 + 20x^6 - 8x^7 + x^8$.

Find the sixth roots of the following expressions :

8. $1 + 6x + 15x^2 + 20x^3 + 15x^4 + 6x^5 + x^6$.
7. $x^6 - 12ax^5 + 240a^2x^4 - 192a^3x^3 + 60a^4x^2 - 160a^5x + 64a^6$.
8. $a^6 - 18a^5x + 135a^4x^2 - 540a^3x^3 + 1215a^2x^4 - 1458ax^5 + 729x^6$.

Find the eighth roots of the following expressions :

9. $x^8 - 8x^7y + 28x^6y^2 - 56x^5y^3 + 70x^4y^4 - 56x^3y^5 + 28x^2y^6 - 8xy^7 + y^8$.
10. $\{x^4 + 2(p-1)x^3 + (p^3 - 2p - 1)x^2 - 2(p-1)x + 1\}^4$.

Find to four terms the square root of

11. $1 + x$.
12. $1 - 2x$.
13. $4 + 2x$.
14. $1 - x - x^2$.
15. $a^2 - x$.
16. $x^3 + a^3$.
17. $a^4 - 3x^2$.
18. $9a^3 + 12ax$.

Find to three terms the cube root of

19. $x^3 - a^3$.
20. $8 + x$.
21. $\frac{1}{a^3} + 9x$.
22. $1 - 6x + 21x^2$.
23. $27x^6 - 27x^5 - 18x^4$.
24. $64 - 48x + 9x^2$.

Identities and Transformations.

***222.** DEFINITION. An **identity** is an algebraical statement which is true for all values of the letters involved in it.

Examples. $a^3 + b^3 = (a+b)(a^2 - ab + b^2)$.
 $x^3 + y^3 + z^3 - 3xyz = (x+y+z)(x^2 + y^2 + z^2 - yz - zx - xy)$.

***223.** An identity asserts that two expressions are always equal; and the proof of this equality is called "proving the identity." The method of procedure is to choose one of the expressions given, and to shew by successive transformations that it can be made to assume the form of the other.

Example 1. To prove that

$$bc(b-c) + ca(c-a) + ab(a-b) = -(b-c)(c-a)(a-b).$$

The first side = $bc(b-c) + c^2a - ca^2 + a^2b - ab^2$
 $= bc(b-c) + a^2(b-c) - a(b^2 - c^2)$
 $= (b-c)\{bc + a^2 - a(b+c)\}$
 $= (b-c)\{bc + a^2 - ab - ac\}$
 $= (b-c)\{a(a-b) - c(a-b)\}$
 $= (b-c)(a-b)(a-c)$
 $= -(b-c)(c-a)(a-b),$

changing the signs of the factor $a-c$, so as to preserve cyclic order.
 [Compare Art. 229, Example 3.]

The expression on the left-hand side can be readily put in the following forms:

$$a^2(b-c) + b^2(c-a) + c^2(a-b);$$

$$- \{ a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2) \}.$$

Hence we have the following results:

$$bc(b-c) + ca(c-a) + ab(a-b) = -(b-c)(c-a)(a-b);$$

$$a^2(b-c) + b^2(c-a) + c^2(a-b) = -(b-c)(c-a)(a-b);$$

$$a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2) = (b-c)(c-a)(a-b).$$

These identities are of such frequent occurrence that they should be carefully noticed and remembered.

Example 2. If $2s = a + b + c$ prove that

$$\frac{1}{s-a} + \frac{1}{s-b} + \frac{1}{s-c} - \frac{1}{s} = \frac{abc}{s(s-a)(s-b)(s-c)}.$$

$$\text{The first side} = \left(\frac{1}{s-a} + \frac{1}{s-b} \right) + \left(\frac{1}{s-c} - \frac{1}{s} \right)$$

$$= \frac{s-b+s-a}{(s-a)(s-b)} + \frac{s-s+c}{s(s-c)}$$

$$= \frac{2s-a-b}{(s-a)(s-b)} + \frac{c}{s(s-c)}$$

$$= \frac{c}{(s-a)(s-b)} + \frac{c}{s(s-c)}$$

$$= c \left\{ \frac{s(s-c) + (s-a)(s-b)}{s(s-a)(s-b)(s-c)} \right\}$$

$$= \frac{c \{ s^2 - cs + s^2 - as - bs + ab \}}{s(s-a)(s-b)(s-c)}$$

$$= \frac{c \{ 2s^2 - s(a+b+c) \}}{s(s-a)(s-b)(s-c)}$$

$$= \frac{abc}{s(s-a)(s-b)(s-c)}, \text{ for } s(a+b+c) = s \cdot 2s = 2s^2.$$

Note. Here $2s$ is a convenient abbreviation of $a+b+c$; and the reduction is much simplified by working in terms of s instead of substituting its value at once. In examples of this kind, as a rule, the student should avoid substituting as long as the work can be carried on in terms of the symbol of abbreviation.

Example 3. If $x^2 + u^2 = 2(xy + yz + zu - y^2 - z^2)$ prove that

$$x = y = z = u.$$

By transposing, we have

$$x^2 - 2xy + y^2 + y^2 - 2yz + z^2 + z^2 - 2zu + u^2 = 0,$$

or

$$(x-y)^2 + (y-z)^2 + (z-u)^2 = 0.$$

Now since the square of any quantity is always positive, each of the expressions $(x-y)^2$, $(y-z)^2$, $(z-u)^2$ is positive. Hence their sum cannot be zero unless each of them be separately equal to zero.

$$\therefore x-y=0, y-z=0, z-u=0;$$

or

$$x=y=z=u.$$

Note. The student should be careful to notice the difference between the conclusions to be drawn from the two statements

$$(x-a)^2 + (y-b)^2 = 0 \dots\dots\dots(1),$$

and

$$(x-a)(y-b) = 0 \dots\dots\dots(2).$$

From (1) we infer that *both* $x-a=0$ and $y-b=0$ *simultaneously*, while from (2) we infer that *either* $x-a=0$ or $y-b=0$.

*EXAMPLES XXIX. c.

Prove the following identities :

1. $b(x^3 + a^3) + ax(x^2 - a^2) + a^3(x + a) = (a+b)(x+a)(x^2 - ax + a^2).$
2. $(ax + by)^2 + (ay - bx)^2 + c^2x^2 + c^2y^2 = (x^2 + y^2)(a^2 + b^2 + c^2).$
3. $(x+y)^3 + 3(x+y)^2z + 3(x+y)z^2 + z^3$
 $= (x+z)^3 + 3(x+z)^2y + 3(x+z)y^2 + y^3.$
4. $(a+b+c)(ab+bc+ca) - abc = (a+b)(b+c)(c+a).$
5. $(a+b+c)^2 - a(b+c-a) - b(a+c-b) - c(a+b-c) = 2(a^2 + b^2 + c^2).$
6. $(x-y)^3 + (x+y)^3 + 3(x-y)^2(x+y) + 3(x+y)^2(x-y) = 8x^3.$
7. $x^2(y-z) + y^2(z-x) + z^2(x-y) + (y-z)(z-x)(x-y) = 0.$
8. $a^3(b-c) + b^3(c-a) + c^3(a-b) = -(b-c)(c-a)(a-b)(a+b+c).$
9. If $x+y+z=0$, prove that $x^3 + y^3 + z^3 = 3xyz.$
10. Prove that $(b-c)^3 + (c-a)^3 + (a-b)^3 = 3(b-c)(c-a)(a-b).$

If $2s = a+b+c$, shew that

11. $(s-a)^2 + (s-b)^2 + (s-c)^2 + s^2 = a^2 + b^2 + c^2.$
12. $(s-a)^3 + (s-b)^3 + (s-c)^3 + 3abc = s^3.$
13. $16s(s-a)(s-b)(s-c) = 2b^2c^2 + 2c^2a^2 + 2a^2b^2 - a^4 - b^4 - c^4.$
14. $2(s-a)(s-b)(s-c) + a(s-b)(s-c) + b(s-c)(s-a)$
 $+ c(s-a)(s-b) = abc.$

If $a+b+c=0$, shew that

15. $(2a-b)^3 + (2b-c)^3 + (2c-a)^3 = 3(2a-b)(2b-c)(2c-a).$
16. $\frac{a^2}{2a^3+bc} + \frac{b^2}{2b^3+ca} + \frac{c^2}{2c^3+ab} = 1.$
17. Prove that
 $(x+y+z)^3 + (x+y-z)^3 + (x-y+z)^3 + (x-y-z)^3 = 4x(x^2 + 3y^2 + 3z^2).$

18. If

19. If

20. Fi

21. Pr

22. Pr

23. If

24. If

25. If

26. If

27. Pr

is

h

28. If

a

*224

to illu
to cycli

18. If $a+b+c=s$, prove that
 $(s-3a)^3 + (s-3b)^3 + (s-3c)^3 - 3(s-3a)(s-3b)(s-3c) = 0$.
19. If $X=b+c-2a$, $Y=c+a-2b$, $Z=a+b-2c$, find the value of
 $X^3 + Y^3 + Z^3 - 3XYZ$.
20. Find the value of $a(a^2+bc)+b(b^2+ac)-c(c^2+ab)$ when $a=.7$,
 $b=.08$, $c=.78$.
21. Prove that $(a-b)^2 + (b-c)^2 + (c-a)^2$
 $= 2(c-b)(c-a) + 2(b-a)(b-c) + 2(a-b)(a-c)$.
22. Prove that $a^2(b^3-c^3) + b^2(c^3-a^3) + c^2(a^3-b^3)$
 $= (a-b)(b-c)(c-a)(ab+bc+ca)$
 $= a^2(b-c)^2 + b^2(c-a)^2 + c^2(a-b)^2$
 $= -[a^2b^2(a-b) + b^2c^2(b-c) + c^2a^2(c-a)]$.
23. If $(a+b)^2 + (b+c)^2 + (c+d)^2 = 4(ab+bc+cd)$, prove that
 $a=b=c=d$.
24. If $x=a+d$, $y=b+d$, $z=c+d$, prove that
 $x^2+y^2+z^2-yz-zx-xy = a^2+b^2+c^2-bc-ca-ab$.
25. If $a+b+c=0$, prove that
 $\frac{1}{b^2+c^2-a^2} + \frac{1}{c^2+a^2-b^2} + \frac{1}{a^2+b^2-c^2} = 0$.
26. If $a+b+c=0$, simplify
 $\frac{b+c}{bc}(b^2+c^2-a^2) + \frac{c+a}{ca}(c^2+a^2-b^2) + \frac{a+b}{ab}(a^2+b^2-c^2)$.
27. Prove that the equation
 $(x-a)^2 + (y-b)^2 + (a^2+b^2-1)(x^2+y^2-1) = 0$,
 is equivalent to the equation
 $(ax+by-1)^2 + (bx-ay)^2 = 0$;
 hence shew that the only possible values of x and y are
 $\frac{a}{a^2+b^2}, \frac{b}{a^2+b^2}$.
28. If $2(x^2+a^2-ax)(y^2+b^2-by) = x^2y^2+a^2b^2$, shew that
 $(x-a)^2(y-b)^2 + (bx-ay)^2 = 0$,
 and therefore that $x=a$, $y=b$ are the only possible solutions.

*224. We shall now give some further examples of fractions to illustrate the advantage of arranging expressions with regard to cyclic order. [Art. 172.]

Example. Find the value of

$$\frac{a}{(a-b)(a-c)(x-a)} + \frac{b}{(b-c)(b-a)(x-b)} + \frac{c}{(c-a)(c-b)(x-c)}.$$

Changing the sign of one factor in each denominator, so as to preserve cyclic order, we get for the lowest common denominator,

$$(a-b)(b-c)(c-a)(x-a)(x-b)(x-c).$$

The whole expression has for its numerator

$$- [a(b-c)(x-b)(x-c) + \dots + \dots]$$

$$\text{or} \quad - [a(b-c)\{x^2 - (b+c)x + bc\} + \dots + \dots].$$

Arrange it according to powers of x ; thus

$$\begin{aligned} \text{coefficient of } x^2 &= -\{a(b-c) + b(c-a) + c(a-b)\} \\ &= 0; \end{aligned}$$

$$\begin{aligned} \text{coefficient of } x &= \{a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2)\} \\ &= (b-c)(c-a)(a-b); \quad [\text{Art. 223.}] \end{aligned}$$

terms which do not contain x

$$\begin{aligned} &= -\{abc(b-c) + abc(c-a) + abc(a-b)\} \\ &= -abc\{b-c+c-a+a-b\} \\ &= 0. \end{aligned}$$

$$\begin{aligned} \text{Hence the expression} &= \frac{(b-c)(c-a)(a-b)x}{(b-c)(c-a)(a-b)(x-a)(x-b)(x-c)} \\ &= \frac{x}{(x-a)(x-b)(x-c)}. \end{aligned}$$

Note. In examples of this kind the work will be much facilitated if the student accustoms himself to readily writing down the following equivalents:

$$\begin{aligned} (b-c) + (c-a) + (a-b) &= 0. \\ a(b-c) + b(c-a) + c(a-b) &= 0. \\ a^2(b-c) + b^2(c-a) + c^2(a-b) &= -(a-b)(b-c)(c-a). \\ bc(b-c) + ca(c-a) + ab(a-b) &= -(a-b)(b-c)(c-a). \\ a(b^2 - c^2) + b(c^2 - a^2) + c(a^2 - b^2) &= (a-b)(b-c)(c-a). \end{aligned}$$

Some of the identities in XXIX. c. may also be remembered with advantage.

*EXAMPLES XXIX. d.

$$1. \quad \frac{a}{(a-b)(a-c)} + \frac{b}{(b-c)(b-a)} + \frac{c}{(c-a)(c-b)}.$$

$$2. \quad \frac{bc}{(a-b)(a-c)} + \frac{ca}{(b-c)(b-a)} + \frac{ab}{(c-a)(c-b)}.$$

3. $\frac{a^2}{(a-b)(a-c)} + \frac{b^2}{(b-c)(b-a)} + \frac{c^2}{(c-a)(c-b)}$
4. $\frac{a^3}{(a-b)(a-c)} + \frac{b^3}{(b-c)(b-a)} + \frac{c^3}{(c-a)(c-b)}$
5. $\frac{a(b+c)}{(a-b)(c-a)} + \frac{b(a+c)}{(a-b)(b-c)} + \frac{c(a+b)}{(c-a)(b-c)}$
6. $\frac{1}{a(a-b)(a-c)} + \frac{1}{b(b-c)(b-a)} + \frac{1}{c(c-a)(c-b)}$
7. $\frac{bc}{a(a^2-b^2)(a^2-c^2)} + \frac{ca}{b(b^2-c^2)(b^2-a^2)} + \frac{ab}{c(c^2-a^2)(c^2-b^2)}$
8. $\frac{(x-b)(x-c)}{(a-b)(a-c)} + \frac{(x-c)(x-a)}{(b-c)(b-a)} + \frac{(x-a)(x-b)}{(c-a)(c-b)}$
9. $\frac{bc(a+d)}{(a-b)(a-c)} + \frac{ca(b+d)}{(b-c)(b-a)} + \frac{ab(c+d)}{(c-a)(c-b)}$
10. $\frac{1}{(a-b)(a-c)(x-a)} + \frac{1}{(b-c)(b-a)(x-b)} + \frac{1}{(c-a)(c-b)(x-c)}$
11. $\frac{a^2}{(a-b)(a-c)(x+a)} + \frac{b^2}{(b-c)(b-a)(x+b)} + \frac{c^2}{(c-a)(c-b)(x+c)}$
12. $\frac{a^2(a+b)(a+c)}{(a-b)(a-c)} + \frac{b^2(b+c)(b+a)}{(b-c)(b-a)} + \frac{c^2(c+a)(c+b)}{(c-a)(c-b)}$
13. $\frac{a^3(b-c) + b^3(c-a) + c^3(a-b)}{(b-c)^3 + (c-a)^3 + (a-b)^3}$
14. $\frac{a^2(b-c) + b^2(c-a) + c^2(a-b) + 2(a-b)(b-c)(c-a)}{(b-c)^3 + (c-a)^3 + (a-b)^3}$
15. $\frac{a^3(b-c) + b^3(c-a) + c^3(a-b)}{a^2(b-c) + b^2(c-a) + c^2(a-b)}$
16. $\frac{a^2(b-c)^3 + b^2(c-a)^3 + c^2(a-b)^3}{(a-b)(b-c)(c-a)}$
17. $\frac{\frac{1}{a}(b-c) + \frac{1}{b}(c-a) + \frac{1}{c}(a-b)}{\frac{1}{a}\left(\frac{1}{b^2} - \frac{1}{c^2}\right) + \frac{1}{b}\left(\frac{1}{c^2} - \frac{1}{a^2}\right) + \frac{1}{c}\left(\frac{1}{a^2} - \frac{1}{b^2}\right)}$
18. $\frac{a^2\left(\frac{1}{c^2} - \frac{1}{b^2}\right) + b^2\left(\frac{1}{a^2} - \frac{1}{c^2}\right) + c^2\left(\frac{1}{b^2} - \frac{1}{a^2}\right)}{\frac{1}{bc}\left(\frac{1}{c} - \frac{1}{b}\right) + \frac{1}{ca}\left(\frac{1}{a} - \frac{1}{c}\right) + \frac{1}{ab}\left(\frac{1}{b} - \frac{1}{a}\right)}$

***225.** To find when $x^3 + px^2 + qx + r$ (1),
is divisible by $x^2 + ax + b$ (2).

Divide (1) by (2) in the ordinary way; thus

$$\begin{array}{r} x^3 + ax + b \overline{) x^3 + px^2 + qx + r} \quad | \quad x + (p-a) \\ \underline{x^3 + ax^2 + bx} \\ (p-a)x^2 + (q-b)x + r \\ \underline{(p-a)x^2 + a(p-a)x + b(p-a)} \\ \{(q-b) - a(p-a)\}x + r - b(p-a) \dots\dots\dots (3) \end{array}$$

Now if the remainder is zero the division is exact. This is the case when

$$\{(q-b) - a(p-a)\}x + r - b(p-a) = 0,$$

or
$$x = \frac{b(p-a) - r}{q-b-a(p-a)}$$

Hence when x has this value, (1) is divisible by (2).

But if in (3), $q-b-a(p-a)=0$,
and also $r-b(p-a)=0$,

the remainder is equal to zero *whatever value* x may have. Thus $x^3 + px^2 + qx + b$ is divisible by $x^2 + ax + b$ for *all* values of x , provided that

$$q-b-a(p-a)=0,$$

and
$$r-b(p-a)=0.$$

***226.** To find the condition that $x^2 + px + q$ may be a perfect square.

Using the ordinary rule for square root, we have

$$\begin{array}{r} x^2 + px + q \left(x + \frac{p}{2} \right. \\ \underline{x^2} \\ px + q \\ \underline{px + \frac{p^2}{4}} \\ q - \frac{p^2}{4} \end{array}$$

If therefore $x^2 + px + q$ be a perfect square, the remainder, $q - \frac{p^2}{4}$, must be zero.

Hence $q - \frac{p^2}{4} = 0$, or $p^2 = 4q$, is the condition required.

*227. To prove that $x^4 + px^3 + qx^2 + rx + s$ is a perfect square
if $\left(q - \frac{p^2}{4}\right)^2 = 4s$ and $r^2 = p^2s$.

The square root must clearly be a trinomial expression of the form $x^2 + lx + m$; if therefore we put

$$x^4 + px^3 + qx^2 + rx + s = (x^2 + lx + m)^2,$$

we have, on expanding the right-hand side

$$x^4 + px^3 + qx^2 + rx + s = x^4 + 2lx^3 + x^2(l^2 + 2m) + 2lmx + m^2.$$

Since this is to be true for all values of x , we may assume that the coefficients of the like powers of x are the same; hence

$$2l = p,$$

$$l^2 + 2m = q,$$

$$2lm = r,$$

$$m^2 = s.$$

From these equations, by eliminating the unknown quantities l and m , we shall obtain the necessary relations between p , q , r , and s .

Thus we have
$$q - \frac{p^2}{4} = 2m = 2\sqrt{s},$$

$$r = 2lm = p\sqrt{s};$$

$$\therefore \left(q - \frac{p^2}{4}\right)^2 = 4s \text{ and } r^2 = p^2s.$$

Note. The method of Art. 226 might have been used here. Also the method of the present article may be used to establish the results of Arts. 225 and 226.

*228. The proposition in the preceding article has been given to illustrate a useful method, which admits of very wide application. In the course of the proof we assume the truth of an important principle; namely,

If two rational integral expressions involving x are identically equal, the coefficients of like powers of x in the two expressions are equal.

[An expression is said to be *rational* when no term contains a square or other root, and it is said to be *integral with respect to x* when the powers of x are all positive integers.]

The demonstration of this principle belongs to a more advanced part of the subject, and could not be discussed completely here. [See *Higher Algebra*. Art. 311.]

The Remainder Theorem.

***229.** *If a rational integral algebraical expression*

$$x^n + p_1x^{n-1} + p_2x^{n-2} + p_3x^{n-3} + \dots + p_{n-1}x + p_n$$

be divided by $x - a$, the remainder will be

$$a^n + p_1a^{n-1} + p_2a^{n-2} + p_3a^{n-3} + \dots + p_{n-1}a + p_n.$$

Divide the given expression by $x - a$ till a remainder is obtained which does not involve x . Let Q be the quotient, and R the remainder; then

$$x^n + p_1x^{n-1} + p_2x^{n-2} + \dots + p_{n-1}x + p_n = Q(x - a) + R.$$

Since R does not contain x , it will remain unaltered whatever value we give to x .

Put $x = a$, then

$$a^n + p_1a^{n-1} + p_2a^{n-2} + \dots + p_{n-1}a + p_n = Q \times 0 + R,$$

$$\therefore R = a^n + p_1a^{n-1} + p_2a^{n-2} + \dots + p_{n-1}a + p_n;$$

which proves the proposition.

From this it appears that when an algebraical expression is divided by $x - a$, the remainder can be obtained at once by writing a in the place of x in the given expression.

Again, the remainder is zero when the given expression is exactly divisible by $x - a$; hence we deduce another important proposition, known as the **Factor Theorem**.

If a rational integral expression involving x become equal to 0 when a is written for x , it will contain $x - a$ as a factor.

Example 1. Resolve into factors $x^3 + 3x^2 - 13x - 15$.

By trial we find that this expression vanishes when $x = 3$; hence $x - 3$ is a factor.

$$\begin{aligned} \therefore x^3 + 3x^2 - 13x - 15 &= x^2(x - 3) + 6x(x - 3) + 5(x - 3) \\ &= (x - 3)(x^2 + 6x + 5) \\ &= (x - 3)(x + 1)(x + 5). \end{aligned}$$

Note. The only numerical values that need be substituted for x are the factors of the last term of the expression. Thus, in the present case, by making trial of -5 , we should have detected the factor $x + 5$.

Exo

 $x + 2$

that is

Or

 $x = -$

Exo

On

factor

and s

dimen

indep

partic

terms

Let

whenc

***23**

made

I.

By

the re

that is

II.

but no

By

the re

hence

odd;

In

 $x + y$

Example 2. The remainder when $x^4 - 2x^3 + x - 7$ is divided by $x + 2$ is

$$(-2)^4 - 2(-2)^3 + (-2) - 7;$$

that is,

$$16 + 16 - 2 - 7, \text{ or } 23.$$

Or the remainder may be found more shortly by substituting $x = -2$ in $\{(x-2)x\}x+1\}x-7$.

Example 3. Find the factors of $bc(b-c) + ca(c-a) + ab(a-b)$.

On trial, this expression vanishes when $b=c$; therefore $b-c$ is a factor. Similarly $c-a$, $a-b$ may be shewn to be factors.

$$\therefore bc(b-c) + ca(c-a) + ab(a-b) = M(b-c)(c-a)(a-b) \dots (1);$$

and since the left-hand member of this identity is only of three dimensions in a, b, c , the factor M must be some numerical quantity independent of a, b, c ; its value can therefore be found by giving particular values to a, b, c , or by equating the coefficients of like terms on each side.

Let $a=0, b=1, c=2$, then (1) becomes

$$2(-1) + 0 + 0 = M(-1) \times 2 \times (-1);$$

whence

$$M = -1.$$

$$\therefore bc(b-c) + ca(c-a) + ab(a-b) = -(b-c)(c-a)(a-b).$$

***230.** We shall now give general proofs of the statements made in Art. 55. We suppose n to be positive and integral.

I. To prove that $x^n - y^n$ is always divisible by $x - y$.

By the remainder theorem when $x^n - y^n$ is divided by $x - y$ the remainder is

$$y^n - y^n, \text{ or } 0,$$

that is, $x^n - y^n$ is always divisible by $x - y$.

II. To prove that $x^n + y^n$ is divisible by $x + y$ when n is odd, but not when n is even.

By the remainder theorem when $x^n + y^n$ is divided by $x + y$ the remainder is

$$(-y)^n + y^n.$$

$$(1) \text{ if } n \text{ is odd, } (-y)^n + y^n = -y^n + y^n = 0;$$

$$(2) \text{ if } n \text{ is even, } (-y)^n + y^n = y^n + y^n = 2y^n;$$

hence there is a remainder when n is even, but none when n is odd; which proves the proposition.

In like manner it may be proved that $x^n - y^n$ is divisible by $x + y$ when n is even; and $x^n + y^n$ is never divisible by $x - y$.

By going through a few steps of the division, the form of the quotient in each case is easily determined. The results of the present article may be conveniently stated as follows :

- (i) For all values of n ,

$$x^n - y^n = (x - y)(x^{n-1} + x^{n-2}y + x^{n-3}y^2 + \dots + y^{n-1}).$$
- (ii) When n is odd,

$$x^n + y^n = (x + y)(x^{n-1} - x^{n-2}y + x^{n-3}y^2 - \dots + y^{n-1}).$$
- (iii) When n is even,

$$x^n - y^n = (x + y)(x^{n-1} - x^{n-2}y + x^{n-3}y^2 - \dots - y^{n-1}).$$

*EXAMPLES XXIX. e.

Find the values of x which will make each of the following expressions a perfect square :

1. $x^4 + 6x^3 + 13x^2 + 13x - 1$.
2. $x^4 + 6x^3 + 11x^2 + 3x + 31$.
3. $x^4 - 2ax^3 + (a^2 + 2b)x^2 - 3abx + 2b^2$.
4. $4p^2x^4 - 4pqx^3 + (q^2 + 2p^2)x^2 - 5pqx + \frac{p^2}{2}$.
5. $\frac{a^2x^6}{9} - \frac{abx^4}{2} + \frac{2acx^3}{3} + \frac{9b^2x^2}{16} - \frac{5bcx}{2} + 6c^2$.
6. $x^4 + 2ax^3 + 3a^2x^2 + cx + d$.
7. Find the conditions that $x^4 - ax^3 + bx^2 - cx + 1$ may be a perfect square for all values of x .

Find the values of x which will make each of the following expressions a perfect cube :

8. $8x^3 - 36x^2 + 56x - 39$.
9. $\frac{x^6}{27} - \frac{a^2x^4}{3} + 4a^4x^2 - 28a^6$.
10. $m^3x^6 - 9m^2nx^4 + 39mn^2x^2 - 51n^3$.
11. Find the relation between b and c in order that

$$x^3 + 3ax^2 + bx + c$$
may be a perfect cube for all values of x .
12. Find the conditions that

$$x^6 + 3ax^5 + 3bx^4 + a(6b - 5a^2)x^3 + 3b(b - a^2)x^2 + 3cx + d$$
may be a perfect cube for all values of x .
13. What number must be added to $x^3 + 2x^2$ in order that the expression may be divisible by $x + 4$?
14. If $x + a$ be a common factor of $x^2 + px + q$ and $x^2 + lx + m$, shew that $a = \frac{m - q}{l - p}$.

Resolve into factors :

15. $x^3 - 6x^2 + 11x - 6$. 16. $x^3 - 5x^2 - 2x + 24$.
 17. $x^3 + 9x^2 + 26x + 24$. 18. $x^3 - x^2 - 41x + 105$.
 19. $x^3 - 39x + 70$. 20. $x^3 - 8x^2 - 31x - 22$.
 21. $6x^3 + 7x^2 - x - 2$. 22. $6x^3 + x^2 - 19x + 6$.

Write down the quotient in the following cases :

23. $\frac{x^2 + y^2}{x + y}$. 24. $\frac{x^3 - y^3}{x + y}$. 25. $\frac{x^6 - y^6}{x - y}$. 26. $\frac{x^9 - y^9}{x - y}$.

Find the square root of

27. $x^4 + (2a - 4)x^3 + (a^2 - 2a + 4)x^2 + (2a^2 - 4a)x + a^2$.
 28. $(a + 1)^2x^4 + (2a^2 + 2a)x^3 + (3a^2 - 4a - 6)x^2 + (2a^2 - 6a)x + a^2 - 6a + 9$.
 29. Find what values of m make $3mx^2 + (6m - 12)x + 8$ a perfect square.
 30. If $4x^4 + 12x^3y + Px^2y^2 + 6xy^3 + y^4$ is a perfect square, find P .

Without actual division shew that

31. $32x^{10} - 33x^5 + 1$ is divisible by $x - 1$.
 32. $3x^4 + 5x^3 - 13x^2 - 20x + 4$ $x^2 - 4$.
 33. $x^4 + 4x^3 - 5x^2 - 36x - 36$ $x^2 - x - 6$.

Without actual division find the remainder when

34. $x^5 - 5x^2 + 5$ is divided by $x - 5$.
 35. $x^3 - 7x^2a + 8xa^2 + 15a^3$ $x + 2a$.
 36. If $ax^2 - bx + c$ and $dx^3 - bx + c$ have a common factor, then
 $a^3 - abd + cd^2 = 0$.
 37. If n be any positive integer, prove that $5^{2n} - 1$ is always divisible by 24.
 38. Shew that $1 - x - x^n + x^{n+1}$ is exactly divisible by
 $1 - 2x + x^2$.
 39. If $x^3 + px + r$ and $3x^2 + p$ have a common factor, prove that
 $\frac{p^3}{27} + \frac{r^2}{4} = 0$.
 40. Shew that if $x^n + py^n + qz^n$ is exactly divisible by
 $x^2 - (ay + bz)x + aby$,
 then $\frac{p}{a^n} + \frac{q}{b^n} + 1 = 0$.

CHAPTER XXX.

THE THEORY OF INDICES.

231. HITHERTO all the definitions and rules with regard to indices have been based upon the supposition that they were positive integers; for instance

- (1) $a^{14} = a . a . a \dots$ to fourteen factors.
- (2) $a^{14} \times a^3 = a^{14+3} = a^{17}.$
- (3) $a^{14} \div a^3 = a^{14-3} = a^{11}.$
- (4) $(a^{14})^3 = a^{14 \times 3} = a^{42}.$

The object of the present chapter is twofold: first, to give *general* proofs which shall establish the laws of combination in the case of all positive integral indices; secondly, to explain how, in strict accordance with these laws, intelligible meanings may be given to symbols whose indices are fractional, zero, or negative.

We shall begin by proving, directly from the definition of a positive integral index, three important propositions.

232. DEFINITION. When m is a *positive integer*, a^m stands for the product of m factors each equal to a .

233. PROP. I. To prove that $a^m \times a^n = a^{m+n}$, when m and n are positive integers.

By definition, $a^m = a . a . a \dots$ to m factors;

$a^n = a . a . a \dots$ to n factors;

$$\begin{aligned} \therefore a^m \times a^n &= (a . a . a \dots \text{to } m \text{ factors}) \times (a . a . a \dots \text{to } n \text{ factors}) \\ &= a . a . a \dots \text{to } m+n \text{ factors} \\ &= a^{m+n}, \text{ by definition.} \end{aligned}$$

COR. If p is also a positive integer, then

$$a^m \times a^n \times a^p = a^{m+n+p};$$

and so for any number of factors.

CHAP.

234

are po

235

positiv

(a

the br

236

indices

intellig

and in

But

indices

have at

definiti

positio

or nega

Now

negativ

laws.

$\frac{p}{a^q}$, a^{-n} ,

the fun

which

symbols

in Prop

237.

integers.

Since

by repla

E.A.

234. PROP. II. To prove that $a^m \div a^n = a^{m-n}$, when m and n are positive integers, and $m > n$.

$$\begin{aligned} a^m \div a^n &= \frac{a^m}{a^n} = \frac{a \cdot a \cdot a \dots \text{to } m \text{ factors}}{a \cdot a \cdot a \dots \text{to } n \text{ factors}} \\ &= a \cdot a \cdot a \dots \text{to } m-n \text{ factors} \\ &= a^{m-n}. \end{aligned}$$

235. PROP. III. To prove that $(a^m)^n = a^{mn}$, when m and n are positive integers.

$$\begin{aligned} (a^m)^n &= a^m \cdot a^m \cdot a^m \dots \text{to } n \text{ factors} \\ &= (a \cdot a \cdot a \dots \text{to } m \text{ factors})(a \cdot a \cdot a \dots \text{to } m \text{ factors}) \dots \end{aligned}$$

the bracket being repeated n times,

$$\begin{aligned} &= a \cdot a \cdot a \dots \text{to } mn \text{ factors} \\ &= a^{mn}. \end{aligned}$$

236. These are the fundamental laws of combination of indices, and they are proved directly from a definition which is intelligible only on the supposition that the indices are *positive* and *integral*.

But it is found convenient to use fractional and negative indices, such as $a^{\frac{1}{2}}$, a^{-7} , $a^{\frac{p}{q}}$, more generally, $a^{\frac{p}{q}}$, a^{-n} ; and these have at present no intelligible meaning. For it is plain that the definition of a^m , [Art. 232], upon which we based the three propositions just proved, is no longer applicable when m is *fractional*, or *negative*.

Now it is important that all indices, whether positive or negative, integral or fractional, should be governed by the same laws. We therefore determine meanings for symbols such as $a^{\frac{p}{q}}$, a^{-n} , in the following way: we assume that they conform to the fundamental law, $a^m \times a^n = a^{m+n}$, and accept the meaning to which this assumption leads us. It will be found that the symbols so interpreted will also obey the other laws enunciated in Props. II. and III.

237. To find a meaning for $a^{\frac{p}{q}}$, p and q being positive integers.

Since $a^m \times a^n = a^{m+n}$ is to be true for all values of m and n , by replacing each of the indices m and n by $\frac{p}{q}$, we have

$$a^{\frac{p}{q}} \times a^{\frac{p}{q}} = a^{\frac{p}{q} + \frac{p}{q}} = a^{\frac{2p}{q}}.$$

E.A.

Q

Similarly, $a^{\frac{p}{q}} \times a^{\frac{p}{q}} \times a^{\frac{p}{q}} = a^{\frac{3p}{q}} = a^{\frac{2p}{q} + \frac{p}{q}} = a^{\frac{2p}{q}} \times a^{\frac{p}{q}}$.

Proceeding in this way for 4, 5, ... q factors, we have

$$a^{\frac{p}{q}} \times a^{\frac{p}{q}} \times a^{\frac{p}{q}} \dots \text{to } q \text{ factors} = a^{\frac{qp}{q}};$$

that is, $(a^{\frac{p}{q}})^q = a^p$.

Therefore, by taking the q^{th} root,

$$a^{\frac{p}{q}} = \sqrt[q]{a^p}.$$

or, in words, $a^{\frac{p}{q}}$ is equal to "the q^{th} root of a^p ."

Examples.

(1) $x^{\frac{5}{7}} = \sqrt[7]{x^5},$

(2) $a^{\frac{1}{3}} = \sqrt[3]{a}.$

(3) $4^{\frac{3}{2}} = \sqrt{4^3} = \sqrt{64} = 8.$

(4) $a^{\frac{3}{4}} \times a^{\frac{5}{4}} = a^{\frac{3}{4} + \frac{5}{4}} = a^{\frac{8}{4}}.$

(5) $k^{\frac{9}{8}} \times k^{\frac{3}{8}} = k^{\frac{9}{8} + \frac{3}{8}} = k^{\frac{12}{8}}.$

(6) $3a^{\frac{3}{4}}b^{\frac{1}{4}} \times 4a^{\frac{1}{4}}b^{\frac{3}{4}} = 12a^{\frac{3}{4} + \frac{1}{4}}b^{\frac{1}{4} + \frac{3}{4}} = 12a^1b^1.$

238. To find a meaning for a^0 .

Since $a^m \times a^n = a^{m+n}$ is to be true for *all* values of m and n , by replacing the index m by 0, we have

$$a^0 \times a^n = a^{0+n}$$

$$= a^n;$$

$$\therefore a^0 = \frac{a^n}{a^n}$$

$$= 1.$$

Hence *any quantity* with zero index is equivalent to 1

Example. $x^{b-c} \times x^{c-b} = x^{b-c+c-b} = x^0 = 1.$

239. To find a meaning for a^{-n} .

Since $a^m \times a^n = a^{m+n}$ is to be true for *all* values of m and n , by replacing the index m by $-n$, we have

$$a^{-n} \times a^n = a^{-n+n} = a^0.$$

But

$$a^0 = 1;$$

hence

$$a^{-n} = \frac{1}{a^n},$$

and

$$a^n = \frac{1}{a^{-n}}.$$

From this it follows that any *factor* may be transferred from the numerator to the denominator of an expression, or vice-versa, by merely changing the sign of the index.

Examples. (1) $x^{-3} = \frac{1}{x^3}.$

(2) $\frac{1}{y^{-\frac{1}{2}}} = y^{\frac{1}{2}} = \sqrt{y}.$

(3) $27^{-\frac{2}{3}} = \frac{1}{27^{\frac{2}{3}}} = \frac{1}{\sqrt[3]{(27)^2}} = \frac{1}{\sqrt[3]{3^6}} = \frac{1}{3^2} = \frac{1}{9}.$

240. To prove that $a^m \div a^n = a^{m-n}$ for all values of m and n .

$$a^m \div a^n = a^m \times \frac{1}{a^n}$$

$$= a^m \times a^{-n}$$

$= a^{m-n}$, by the fundamental law.

Examples. (1) $a^3 \div a^5 = a^{3-5} = a^{-2} = \frac{1}{a^2}.$

(2) $c \div c^{-\frac{2}{3}} = c^{1+\frac{2}{3}} = c^{\frac{5}{3}}.$

(3) $x^{a-b} \div x^{a-c} = x^{a-b-(a-c)} = x^{c-b}.$

241. The method of finding a meaning for a symbol, as explained in the preceding articles, deserves careful attention. The usual algebraical process is to make choice of symbols, give them meanings, and then prove the rules for their combination. Here the process is reversed; the symbols are given, and the law to which they are to conform, and from this the meanings of the symbols are determined.

242. The following examples will illustrate the different principles we have established.

Examples. (1) $\frac{3a^{-2}}{5x^{-1}y} = \frac{3x}{5a^2y}.$

(2) $\frac{2a^{\frac{1}{2}} \times a^{\frac{3}{4}} \times 6a^{-\frac{7}{4}}}{9a^{-\frac{1}{2}} \times a^{\frac{1}{4}}} = \frac{4}{3} a^{\frac{1}{2} + \frac{3}{4} - \frac{7}{4} - \frac{1}{2}} = \frac{4}{3} a^{-1} = \frac{4}{3a}.$

$$(3) \frac{\sqrt{x^3} \times \sqrt[3]{y^2}}{\sqrt[6]{y^{-2}} \times \sqrt[4]{x^6}} = \frac{x^{\frac{3}{2}} \times y^{\frac{2}{3}}}{y^{-\frac{1}{3}} \times x^{\frac{3}{2}}} = x^{\frac{3}{2} - \frac{3}{2}} y^{\frac{2}{3} + \frac{1}{3}} = x^0 y = y.$$

$$(4) 2\sqrt{a} + \frac{3}{a^{-\frac{1}{2}}} + a^{\frac{5}{2}} = 2a^{\frac{1}{2}} + 3a^{\frac{1}{2}} + a^{\frac{5}{2}} \\ = 5a^{\frac{1}{2}} + a^{\frac{5}{2}} = a^{\frac{1}{2}}(5 + a^2).$$

EXAMPLES XXX. a.

Express with positive indices :

- | | | | |
|--|---|---|---|
| 1. $2x^{-\frac{1}{2}}$ | 2. $3a^{-\frac{2}{3}}$ | 3. $4x^{-2}a^3$ | 4. $3 \div a^{-2}$ |
| 5. $\frac{1}{4a^{-2}}$ | 6. $\frac{1}{5x^{-\frac{1}{2}}}$ | 7. $\frac{3a^{-2}x^2}{5y^2c^{-4}}$ | 8. $\frac{x^ay^{-b}}{b^{-a}}$ |
| 9. $2x^{\frac{1}{2}} \times 3x^{-1}$ | 10. $1 \div 2a^{-\frac{1}{2}}$ | 11. $xy^2 \times x^{-1}$ | 12. $a^{-2}x^{-1} \div 3x$ |
| 13. $\frac{1}{\sqrt{x^3}}$ | 14. $\frac{1}{4\sqrt[5]{x^{-3}}}$ | 15. $\frac{2}{\sqrt{y^{-3}}}$ | 16. $\frac{\sqrt[4]{x^3}}{\sqrt{x^{-1}}}$ |
| 17. $a^{-2}x^{-\frac{1}{2}} \div a^{-3}$ | 18. $\sqrt[3]{a^{-1}} \div \sqrt[3]{a}$ | 19. $\sqrt[5]{a^{-3}} \div \sqrt[5]{a^7}$ | |

Express with radical signs and positive indices :

- | | | | |
|---|---|---|--|
| 20. $x^{\frac{3}{4}}$ | 21. $a^{-\frac{1}{2}}$ | 22. $5x^{-\frac{1}{2}}$ | 23. $2a^{-\frac{1}{2}}$ |
| 24. $\frac{1}{2a^{\frac{1}{2}}}$ | 25. $\frac{2}{b^{-\frac{3}{4}}}$ | 26. $\frac{c^{-\frac{1}{2}}}{2}$ | 27. $\frac{1}{x^{-\frac{1}{2}}}$ |
| 28. $a^{-\frac{1}{2}} \times 2a^{-\frac{1}{2}}$ | 29. $x^{-\frac{2}{3}} \div 2a^{-\frac{1}{2}}$ | 30. $7a^{-\frac{1}{2}} \times 3a^{-1}$ | |
| 31. $\frac{2a^{-2}}{a^{-\frac{1}{2}}}$ | 32. $\frac{a^{-\frac{1}{2}}}{3a}$ | 33. $\frac{4x^{-1}}{x^{-\frac{1}{2}}}$ | 34. $\frac{\sqrt[3]{x^{-c}}}{\sqrt[3]{x^2}}$ |
| 35. $\sqrt[3]{a^2} \times \sqrt[2]{a^3}$ | 36. $\sqrt[5]{a^{-x}} \div \sqrt[5]{a^{-2x}}$ | 37. $\sqrt[2a]{x} \times \sqrt[3]{x^2}$ | |
| 38. $\sqrt[2]{a} \div \sqrt[2]{x^3}$ | 39. $\sqrt[3]{a^3} \div \sqrt[3]{a^2}$ | 40. $\sqrt[4]{a^n} \times \sqrt[3]{a^n} \div \sqrt[12]{a^{5n}}$ | |

Find the value of

41. $16^{\frac{3}{4}}$. 42. $4^{-\frac{5}{2}}$. 43. $125^{\frac{2}{3}}$. 44. $8^{-\frac{2}{3}}$. 45. $36^{-\frac{1}{2}}$.
 46. $\frac{1}{25^{-2}}$. 47. $243^{\frac{2}{3}}$. 48. $\left(\frac{8}{27}\right)^{-\frac{1}{3}}$. 49. $\left(\frac{81}{16}\right)^{\frac{1}{4}}$. 50. $\left(\frac{32}{243}\right)^{-\frac{1}{5}}$.

243. To prove that $(a^m)^n = a^{mn}$ is universally true for all values of m and n .

CASE I. Let n be a positive integer.

Now, whatever be the value of m

$$\begin{aligned}(a^m)^n &= a^m \cdot a^m \cdot a^m \dots \text{to } n \text{ factors} \\ &= a^{m+m+m+\dots \text{to } n \text{ terms}} \\ &= a^{mn}.\end{aligned}$$

CASE II. Let m be unrestricted as before, and let n be a positive fraction. Replacing n by $\frac{p}{q}$, where p and q are positive integers, we have $(a^m)^n = (a^m)^{\frac{p}{q}}$.

Now the q^{th} power of $(a^m)^{\frac{p}{q}} = \{(a^m)^{\frac{p}{q}}\}^q$

$$= (a^m)^{\frac{p}{q} \cdot q}, \quad [\text{Case I.}]$$

$$= (a^m)^p \quad [\text{Case I.}]$$

$$= a^{mp}.$$

Hence by taking the q^{th} root of these equals,

$$\begin{aligned}(a^m)^{\frac{p}{q}} &= \sqrt[q]{a^{mp}} \\ &= a^{\frac{mp}{q}}.\end{aligned} \quad [\text{Art. 237.}]$$

CASE III. Let m be unrestricted as before, and let n be any negative quantity. Replacing n by $-r$, where r is positive, we have

$$(a^m)^n = (a^m)^{-r} = \frac{1}{(a^m)^r}, \quad [\text{Art. 239.}]$$

$$= \frac{1}{a^{mr}}, \quad [\text{Case II.}]$$

$$= a^{-mr} = a^{mn}.$$

Hence Prop. III., Art. 235, $(a^m)^n = a^{mn}$ has been shewn to be universally true.

- Examples.* (1) $(b^{\frac{2}{3}})^{\frac{3}{2}} = b^{\frac{2}{3} \times \frac{3}{2}} = b^1$.
 (2) $\{(x^{-2})^3\}^{-4} = (x^{-6})^{-4} = x^{24}$.
 (3) $\left(x^{\frac{1}{a-c}}\right)^{a^2-c^2} = x^{\frac{1}{a-c} \times (a^2-c^2)} = x^{a+c}$.

244. To prove that $(ab)^n = a^n b^n$, whatever be the value of n ; a and b being any quantities whatever.

CASE I. Let n be a *positive integer*.

Now $(ab)^n = ab \cdot ab \cdot ab \dots$ to n factors
 $= (a \cdot a \cdot a \dots$ to n factors $)(b \cdot b \cdot b \dots$ to n factors)
 $= a^n b^n$.

CASE II. Let n be a *positive fraction*. Replacing n by $\frac{p}{q}$, where p and q are *positive integers*, we have $(ab)^n = (ab)^{\frac{p}{q}}$.

Now the q^{th} power of $(ab)^{\frac{p}{q}} = \{(ab)^{\frac{p}{q}}\}^q$
 $= (ab)^p$, [Art. 243.]
 $= a^p b^p$
 $= (a^{\frac{p}{q}} b^{\frac{p}{q}})^q$. [Case I.]

Taking the q^{th} root, $(ab)^{\frac{p}{q}} = a^{\frac{p}{q}} b^{\frac{p}{q}}$.

CASE III. Let n have any *negative value*. Replacing n by $-r$, where r is positive,

$$\begin{aligned}(ab)^n &= (ab)^{-r} = \frac{1}{(ab)^r} \\ &= \frac{1}{a^r b^r} = a^{-r} b^{-r} \\ &= a^n b^n.\end{aligned}$$

Hence the proposition is proved universally.

The result we have just proved may be expressed in a verbal form by saying that the index of a product may be *distributed* over its *factors*.

Note. An index is not distributive over the *terms* of an expression. Thus $(a^{\frac{1}{2}} + b^{\frac{1}{2}})^2$ is not equal to $a + b$. Again $(a^2 + b^2)^{\frac{1}{2}}$ is equal to $\sqrt{a^2 + b^2}$, and cannot be further simplified.

Examples. (1) $(yz)^{a-c}(zx)^c(xy)^{-c} = y^{a-c}z^{a-c}z^c x^c x^{-c} y^{-c}$
 $= y^{a-2c}z^a.$

(2) $\{(a-b)^k\}^{-i} \times \{(a+b)^{-k}\}^l = (a-b)^{-ki} \times (a+b)^{-kl}$
 $= \{(a-b)(a+b)\}^{-kl}$
 $= (a^2 - b^2)^{-kl}.$

245. It should be observed that in the proof of Art. 244 the quantities a and b are *wholly unrestricted*, and may themselves involve indices.

Examples. (1) $(x^{\frac{1}{2}}y^{-\frac{1}{2}})^{\frac{4}{3}} \div (x^2y^{-1})^{-\frac{1}{2}} = x^{\frac{2}{3}}y^{-\frac{2}{3}} \div x^{-\frac{1}{2}}y^{\frac{1}{2}}$
 $= x^{\frac{4}{3}}y^{-1}.$

(2) $\left(\frac{\alpha^{\frac{2}{3}}\sqrt{b^{-1}}}{b^{\frac{2}{3}}\sqrt{a^{-2}}}\right)^6 \div \left(\frac{\sqrt{\alpha}\sqrt{b^{-4}}}{b\sqrt{a^{-2}}}\right)^6 = \left(\frac{\alpha^{\frac{2}{3}}b^{-\frac{1}{2}}}{ba^{-\frac{2}{3}}}\right)^6 \div \left(\frac{\sqrt{ab^{-2}}}{ba^{-1}}\right)^6$
 $= \left(\frac{\alpha^{\frac{2}{3}}b^{-\frac{1}{2}}}{ba^{-\frac{2}{3}}}\right)^6 \div \left(\frac{\alpha^{\frac{1}{2}}b^{-\frac{1}{2}}}{ab^{-\frac{1}{2}}}\right)^6$
 $= (a^{\frac{1}{3}})^6 = a^2.$

EXAMPLES XXX. b.

Simplify and express with positive indices :

1. $(\sqrt{\alpha^2b^3})^6.$
2. $(\sqrt[3]{x^{-4}y^3})^{-3}.$
3. $(x^ay^{-b})^3 \times (x^3y^2)^{-a}.$
4. $\left(\frac{16x^2}{y^{-2}}\right)^{-\frac{1}{2}}.$
5. $\left(\frac{27x^3}{8a^{-3}}\right)^{-\frac{2}{3}}.$
6. $\left(\frac{\alpha^{-\frac{1}{2}}}{4c^3}\right)^{-2}.$
7. $\left\{\sqrt[4]{\left(x^{-\frac{3}{2}}y^{\frac{1}{2}}\right)^3}\right\}^{-\frac{3}{2}}.$
8. $\sqrt[4]{x^2/x^{-1}}.$
9. $(4a^{-2} \div 9x^2)^{-\frac{1}{2}}.$
10. $(x \div 2/x)^n.$
11. $(x \times 2/x^{-\frac{1}{n}})^{\frac{n^2}{1-n}}.$
12. $(2/x^b \div 3/x)^{\frac{1}{1-a}}.$
13. $\sqrt{a^{-2}b} \times \sqrt[3]{ab^{-3}}.$
14. $\sqrt[2]{ab^{-1}c^{-2}} \times (a^{-1}b^{-2}c^{-4})^{-\frac{1}{3}}.$
15. $\sqrt[6]{a^{4b}x^6} \times (a^{\frac{2}{3}}x^{-1})^{-b}.$
16. $\sqrt[3]{x^{-1}}\sqrt{y^3} \div \sqrt{y^3/x}.$

Simplify and express with positive indices :

17. $(a^{-\frac{1}{2}}\sqrt{x})^{-3} \times \sqrt{x^{-2}}\sqrt{a^{-6}}$.
18. $\sqrt[n]{a^{n+k}b^{2n-k}} \div (a^{\frac{1}{n}}b^{-\frac{1}{n}})^k$.
19. $\sqrt[3]{(a+b)^6} \times (a+b)^{-\frac{2}{3}}$.
20. $\{(x-y)^{-3}\}^n \div \{(x+y)^n\}^3$.
21. $\left(\frac{a^{-2b}}{a^3b^{-4}}\right)^{-3} \div \left(\frac{ab^{-1}}{a^{-3}b^3}\right)^5$.
22. $\left\{\sqrt[3]{\frac{a}{b^{-1}}} \cdot \left(\frac{b^{\frac{1}{2}}}{a^{\frac{1}{3}}}\right)^2 \div \frac{a^{-\frac{1}{6}}}{b^{-\frac{1}{2}}}\right\}^6$.
23. $\left(a^{-\frac{1}{2}}x^{\frac{1}{3}}\sqrt{ax^{-\frac{1}{3}}\sqrt[4]{x^{\frac{2}{3}}}}\right)^{\frac{1}{2}}$.
24. $\sqrt[4]{(a+b)^6} \times (a^2 - b^2)^{-\frac{1}{2}}$.
25. $\left(\frac{a^{-3}}{b^{-\frac{1}{2}}c}\right)^{-2} \div \left(\frac{\sqrt{a^{-\frac{1}{2}}}}{a^2c^{-1}} \cdot \sqrt[6]{b^3}\right)^{-2}$.
26. $\left(\frac{a^{-\frac{3}{2}}x^{\frac{1}{2}}}{x^{-1}a}\right)^2 \div \sqrt[3]{\frac{a^{-1}}{x^{-3}}}$.
27. $\left(\sqrt[5]{\frac{a^{\frac{1}{2}}x^{-2}}{x^{\frac{1}{2}}a^{-2}}} \times \sqrt[3]{\frac{a\sqrt{x}}{x^{-1}\sqrt{a}}}\right)^{-4}$.
28. $\frac{\sqrt[3]{(a^3b^3 + a^6)}}{\sqrt[3]{(b^6 - a^3b^3)^{-1}}}$.
29. $(a^{n^2-1})^{\frac{n}{n+1}} + \frac{\sqrt[n]{a^{2n}}}{a}$.
30. $(x^{\frac{n}{n+1}})^{n^2-1} + \frac{\sqrt{x^{2n}}}{x}$.
31. $\left\{\frac{a^{p-q}}{\sqrt[2]{a^{q^2-p^2}}} \times a^{2(p-q)}\right\}^n$.
32. $(x^{\frac{a}{b}}y^{-1})^b \div \left(\frac{x^{a^2-b^2}}{y^{ab+b^3}}\right)^{\frac{1}{a+b}}$.
33. $\left(\frac{x^{-2}y^3}{x^3y^{-2}}\right)^{-\frac{1}{2}} \times \left(\frac{y^3x^{-3}}{x^3y^{-3}}\right)^{-1}$.
34. $\left(\frac{y^{-3}}{x^{\frac{2}{3}}z^{-1}}\right)^{-\frac{2}{3}} \times \left(\frac{y^{\frac{1}{3}}x^{-1}}{z^{-\frac{2}{4}}}\right)^{\frac{2}{3}}$.
35. $\frac{2^n \times (2^n-1)^n}{2^{n+1} \times 2^{n-1}} \times \frac{1}{4^{-n}}$.
36. $\frac{2^{n+1}}{(2^n)^{n-1}} \div \frac{4^{n+1}}{(2^n-1)^{n+1}}$.
37. $\frac{3 \cdot 2^n - 4 \cdot 2^{n-2}}{2^n - 2^{n-1}}$.
38. $\frac{3^{n+4} - 6 \cdot 3^{n+1}}{3^{n+2} \times 7}$.

246. Since the index-laws are universally true, all the ordinary operations of multiplication, division, involution and evolution are applicable to expressions which contain fractional and negative indices.

247. In Art. 121, we pointed out that the descending powers of x are

$$\dots x^3, x^2, x, 1, \frac{1}{x}, \frac{1}{x^2}, \frac{1}{x^3}, \dots$$

A reason for this may be seen if we write these terms in the form

$$\dots x^3, x^2, x^1, x^0, x^{-1}, x^{-2}, x^{-3}, \dots$$

Note.
of negat

Example 1. Multiply $3x^{-\frac{1}{3}} + x + 2x^{\frac{2}{3}}$ by $x^{\frac{1}{3}} - 2$.

Arrange in descending powers of x .

$$\begin{array}{r} x + 2x^{\frac{2}{3}} + 3x^{-\frac{1}{3}} \\ x^{\frac{1}{3}} - 2 \\ \hline x^{\frac{4}{3}} + 2x + 3 \\ - 2x - 4x^{\frac{2}{3}} - 6x^{-\frac{1}{3}} \\ \hline x^{\frac{4}{3}} - 4x^{\frac{2}{3}} + 3 - 6x^{-\frac{1}{3}}. \end{array}$$

Example 2. Divide $16a^{-3} - 6a^{-2} + 5a^{-1} + 6$ by $1 + 2a^{-1}$.

$$\begin{array}{r} 2a^{-1} + 1 \overline{) 16a^{-3} - 6a^{-2} + 5a^{-1} + 6} \\ \underline{16a^{-3} + 8a^{-2}} \\ - 14a^{-2} + 5a^{-1} \\ \underline{- 14a^{-2} - 7a^{-1}} \\ 12a^{-1} + 6 \\ \underline{12a^{-1} + 6} \\ 0 \end{array}$$

Example 3. Find the square root of

$$\frac{4x^2}{y} + \frac{\sqrt{x^3}}{y^{-\frac{1}{2}}} - 2x + \frac{y}{4} + x^3 - 4\sqrt{(x^2y^{-1})}.$$

Getting rid of the radical signs, and arranging in descending powers of x , we have

$$\begin{array}{r} x^3 - 4x^{\frac{5}{2}}y^{-\frac{1}{2}} + 4x^2y^{-1} + x^{\frac{3}{2}}y^{\frac{1}{2}} - 2x + \frac{y}{4} \left(x^{\frac{3}{2}} - 2xy^{-\frac{1}{2}} + \frac{y^{\frac{1}{2}}}{2} \right) \\ \hline 2x^{\frac{3}{2}} - 2xy^{-\frac{1}{2}} \left| \begin{array}{l} - 4x^{\frac{5}{2}}y^{-\frac{1}{2}} + 4x^2y^{-1} \\ - 4x^{\frac{3}{2}}y^{-\frac{1}{2}} + 4x^2y^{-1} \end{array} \right. \\ \hline 2x^{\frac{3}{2}} - 4xy^{-\frac{1}{2}} + \frac{y^{\frac{1}{2}}}{2} \left| \begin{array}{l} x^{\frac{3}{2}}y^{\frac{1}{2}} - 2x + \frac{y}{4} \\ x^{\frac{3}{2}}y^{\frac{1}{2}} - 2x + \frac{y}{4} \end{array} \right. \end{array}$$

Note. In this example it should be observed that the introduction of negative indices enables us to avoid the use of algebraical fractions.

EXAMPLES XXX. c.

1. Multiply $3x^{\frac{1}{2}} - 5 + 8x^{-\frac{1}{2}}$ by $4x^{\frac{1}{2}} + 3x^{-\frac{1}{2}}$.
2. Multiply $3a^{\frac{2}{3}} - 4a^{\frac{1}{3}} - a^{-\frac{1}{3}}$ by $3a^{\frac{1}{3}} + a^{-\frac{1}{3}} - 6a^{-\frac{2}{3}}$.
3. Find the product of $c^x + 2c^{-x} - 7$ and $5 - 3c^{-x} + 2c^x$.
4. Find the product of $5 + 2c^{2a} + 3x^{-2a}$ and $4x^a - 3x^{-a}$.
5. Divide $21x + x^{\frac{2}{3}} + x^{\frac{1}{3}} + 1$ by $3x^{\frac{1}{3}} + 1$.
6. Divide $15a - 3a^{\frac{1}{2}} - 2a^{-\frac{1}{2}} + 8a^{-1}$ by $5a^{\frac{1}{2}} + 4$.
7. Divide $16a^{-3} + 6a^{-2} + 5a^{-1} - 6$ by $2a^{-1} - 1$.
8. Divide $5b^{\frac{3}{2}} - 6b^{\frac{1}{2}} - 4b^{-\frac{3}{2}} - 4b^{-\frac{1}{2}} - 5$ by $b^{\frac{1}{2}} - 2b^{-\frac{1}{2}}$.
9. Divide $21a^{3x} + 20 - 27a^x - 23a^{2x}$ by $3a^x - 5$.
10. Divide $8c^{-n} - 8c^n + 5c^{3n} - 3c^{-3n}$ by $5c^n - 3c^{-n}$.

Find the square root of

11. $9x - 12x^{\frac{1}{2}} + 10 - 4x^{-\frac{1}{2}} + x^{-1}$.
12. $25a^{\frac{4}{3}} + 16 - 30a - 24a^{\frac{1}{3}} + 49a^{\frac{2}{3}}$.
13. $4x^n + 9x^{-n} + 28 - 24x^{-\frac{n}{2}} - 16x^{\frac{n}{2}}$.
14. $12a^x + 4 - 6a^{3x} + a^{4x} + 5a^{2x}$.
15. Multiply $a^{\frac{3}{2}} - 8a^{-\frac{3}{2}} + 4a^{-\frac{1}{2}} - 2a^{\frac{1}{2}}$ by $4a^{-\frac{3}{2}} + a^{\frac{1}{2}} + 4a^{-\frac{1}{2}}$.
16. Multiply $1 - 2\sqrt[3]{x} - 2x^{\frac{1}{2}}$ by $1 - \sqrt[3]{x}$.
17. Multiply $2\sqrt[3]{a^5} - a^{\frac{1}{3}} - \frac{3}{a}$ by $2a - 3\sqrt[3]{\frac{1}{a}} - a^{-\frac{2}{3}}$.
18. Divide $\sqrt[3]{x^3} + 2x^{\frac{1}{2}} - 16x^{-\frac{3}{2}} - \frac{32}{x}$ by $x^{\frac{1}{2}} + 4x^{-\frac{1}{2}} + \frac{4}{\sqrt{x}}$.
19. Divide $1 - \sqrt{a} - \frac{2}{a-1} + 2a^2$ by $1 - a^{\frac{1}{2}}$.
20. Divide $4\sqrt[3]{x^3} - 8x^{\frac{1}{2}} - 5 + \frac{10}{\sqrt[3]{x}} + 3x^{-\frac{3}{2}}$ by $2x^{\frac{1}{6}} - \sqrt[3]{x} - \frac{3}{\sqrt[3]{x}}$.

Find the square root of

$$21. \quad 9x^{-4} - 18x^{-3}\sqrt{y} + \frac{15y}{x^2} - 6\sqrt{\left(\frac{y^3}{x^2}\right)} + y^2.$$

$$22. \quad 4\sqrt{x^3} - 12\sqrt[4]{(x^3y)} + 25\sqrt{y} - 24\sqrt[4]{\left(\frac{y^3}{x^3}\right)} + 16x^{-\frac{3}{2}}y.$$

$$23. \quad 81\left(\frac{\sqrt[3]{x^4}}{y^2} + 1\right) + 36\frac{x^{\frac{1}{3}}}{\sqrt{y}}(x^{\frac{2}{3}}y^{-1} - 1) - 158\frac{\sqrt[3]{x^2}}{y}.$$

$$24. \quad \frac{x^{-2}}{16} + 1 + \frac{9}{\sqrt[3]{y^{-2}}} + \frac{1 - 3\sqrt[3]{y}}{2x} - 6\sqrt[3]{y}.$$

248. The following examples will illustrate the formulæ of earlier chapters when applied to expressions involving fractional and negative indices.

$$\begin{aligned} \text{Example 1. } (a^{\frac{h}{k}} - b^{\frac{p}{q}})(a^{-\frac{h}{k}} + b^{-\frac{p}{q}}) &= a^{\frac{h}{k} - \frac{h}{k}} - a^{-\frac{h}{k} \frac{p}{q}} + a^{\frac{h}{k} b^{-\frac{p}{q}}} - b^{\frac{p}{q} - \frac{p}{q}} \\ &= 1 - a^{-\frac{h}{k} \frac{p}{q}} + a^{\frac{h}{k} b^{-\frac{p}{q}}} - 1 \\ &= a^{\frac{h}{k} b^{-\frac{p}{q}}} - a^{-\frac{h}{k} \frac{p}{q}}. \end{aligned}$$

Example 2. Multiply $2x^{2p} - x^p + 3$ by $2x^{2p} + x^p - 3$.

$$\begin{aligned} \text{The product} &= \{2x^{2p} - (x^p - 3)\} \{2x^{2p} + (x^p - 3)\} \\ &= (2x^{2p})^2 - (x^p - 3)^2 \\ &= 4x^{4p} - x^{2p} + 6x^p - 9. \end{aligned}$$

Example 3. The square of $3x^{\frac{1}{2}} - 2 - x^{-\frac{1}{2}}$

$$\begin{aligned} &= 9x + 4 + x^{-1} - 2 \cdot 3x^{\frac{1}{2}} \cdot 2 - 2 \cdot 3x^{\frac{1}{2}} \cdot x^{-\frac{1}{2}} + 2 \cdot 2 \cdot x^{-\frac{1}{2}} \\ &= 9x + 4 + x^{-1} - 12x^{\frac{1}{2}} - 6 + 4x^{-\frac{1}{2}} \\ &= 9x - 12x^{\frac{1}{2}} - 2 + 4x^{-\frac{1}{2}} + x^{-1}, \end{aligned}$$

by collecting like terms and rearranging.

Example 4. Divide $a^{\frac{h}{n}} + a^{-\frac{h}{n}}$ by $a^{\frac{n}{2}} + a^{-\frac{n}{2}}$.

$$\begin{aligned} \text{The quotient} &= (a^{\frac{h}{n}} + a^{-\frac{h}{n}}) \div (a^{\frac{n}{2}} + a^{-\frac{n}{2}}) \\ &= \{(a^{\frac{h}{n}})^2 + (a^{-\frac{h}{n}})^2\} \div (a^{\frac{n}{2}} + a^{-\frac{n}{2}}) \\ &= (a^{\frac{h}{n}})^2 - a^{\frac{h}{n}} \cdot a^{-\frac{n}{2}} + (a^{-\frac{h}{n}})^2 \\ &= a^n - 1 + a^{-n}. \end{aligned}$$

EXAMPLES XXX. d.

Write down the value of

1. $(x^{\frac{1}{2}} - 7)(x^{\frac{1}{2}} + 3)$.
2. $(4x - 5x^{-1})(4x + 3x^{-1})$.
3. $(7x - 9y^{-1})(7x + 9y^{-1})$.
4. $(x^m - y^n)(x^{-m} + y^{-n})$.
5. $(a^x - 2a^{-x})^2$.
6. $(a^x + a^{\frac{1}{x}})^2$.
7. $(x^{\frac{2}{3}} - \frac{1}{2}x^{-a})^2$.
8. $(5x^ay^b - 3x^{-a}y^{-b})(4x^ay^b + 5x^{-a}y^{-b})$.
9. $(\frac{1}{3}a^{\frac{1}{3}} - a^{-\frac{1}{3}})^2$.
10. $(3x^ay^{-b} + 5x^{-a}y^b)(3x^ay^b - 5x^{-a}y^{-b})$.
11. $(a^x - \frac{1}{2} - a^{-x})^2$.
12. $(x^{\frac{1}{a}} - x^{-\frac{1}{a}} + x)^2$.
13. $\{(a+b)^{\frac{1}{2}} + (a-b)^{\frac{1}{2}}\}^2$.
14. $\{(a+b)^{\frac{1}{2}} - (a-b)^{-\frac{1}{2}}\}^2$.

Write down the quotient of

15. $x - 9a$ by $x^{\frac{1}{2}} + 3a^{\frac{1}{2}}$.
16. $x^{\frac{2}{3}} - 27$ by $x^{\frac{1}{3}} - 3$.
17. $a^{2x} - 16$ by $a^x - 4$.
18. $x^{3a} + 8$ by $x^a + 2$.
19. $c^{2x} - c^{-x}$ by $c^x - c^{-\frac{x}{2}}$.
20. $1 - 8a^{-3}$ by $1 - 2a^{-1}$.
21. $a^{4x} - x^6$ by $a^{2x} + x^3$.
22. $x^{-4} - 1$ by $x^{-1} + 1$.
23. $x^{\frac{2}{3}} - 1$ by $x^{\frac{1}{3}} - 1$.
24. $x^{5n} + 32$ by $x^n + 2$.

Find the value of

25. $(x + x^{\frac{1}{2}} - 4)(x + x^{\frac{1}{2}} + 4)$.
26. $(2x^{\frac{1}{3}} + 4 + 3x^{-\frac{1}{3}})(2x^{\frac{1}{3}} + 4 - 3x^{-\frac{1}{3}})$.
27. $(2 - x^{\frac{1}{2}} + x)(2 + x^{\frac{1}{2}} + x)$.
28. $(a^x + 7 + 3a^{-x})(a^x - 7 - 3a^{-x})$.
29. $\frac{a^{\frac{4}{3}} - 8a^{\frac{1}{3}}b}{a^{\frac{2}{3}} + 2\sqrt[3]{ab} + 4b^{\frac{2}{3}}}$.
30. $\frac{x - 7x^{\frac{1}{2}}}{x - 5\sqrt{x} - 14} \div \left(1 + \frac{2}{\sqrt{x}}\right)^{-1}$.
31. $\frac{x^{\frac{2}{3}} - 4\sqrt[3]{x^{-2}}}{\sqrt[3]{x^3 + 4} + 4x^{-\frac{1}{3}}}$.
32. $\frac{a^{\frac{2}{3}} + ab}{ab - b^3} - \frac{\sqrt{a}}{\sqrt{a} - b}$.

CHAPTER XXXI.

ELEMENTARY SURDS.

249. DEFINITION. If the root of a quantity cannot be exactly obtained the root is called a **surd**.

Thus $\sqrt{2}$, $\sqrt[3]{5}$, $\sqrt[5]{a^3}$, $\sqrt{a^2+b^2}$ are surds.

By reference to the preceding chapter it will be seen that these are only cases of fractional indices; for the above quantities might be written

$$2^{\frac{1}{2}}, 5^{\frac{1}{3}}, a^{\frac{3}{5}}, (a^2+b^2)^{\frac{1}{2}}.$$

Since surds may always be expressed as quantities with fractional indices they are subject to the same laws of combination as other algebraical symbols.

250. A quantity may be expressed in a surd form without really being a surd. Thus $\sqrt[3]{x^6}$ or $x^{\frac{2}{3}}$, though apparently a surd, can be expressed in the equivalent form x^2 .

251. A surd is sometimes called an **irrational quantity**; and quantities which are not surds are, for the sake of distinction, termed **rational quantities**.

252. In the case of numerical surds such as $\sqrt{2}$, $\sqrt[3]{5}$, ..., although the *exact* value can never be found, it can be determined to any degree of accuracy by carrying the process of evolution far enough.

Thus $\sqrt{5} = 2.236068.....$;

that is $\sqrt{5}$ lies between 2.23606 and 2.23607; and therefore the error in using either of these quantities instead of $\sqrt{5}$ is less than .00001. By taking the root to a greater number of decimal places we can approximate still nearer to the true value.

It thus appears that it will never be *absolutely necessary* to introduce surds into numerical work, which can always be carried on to a certain degree of accuracy; but we shall in the present chapter prove laws for combination of surd quantities which will enable us to work with symbols such as $\sqrt{2}$, $\sqrt[3]{5}$, $\sqrt[4]{a}$, ... with absolute accuracy so long as the symbols are kept in their surd form. Moreover it will be found that even where approximate numerical results are required, the work is considerably simpli-

fied and shortened by operating with surd symbols, and afterwards substituting numerical values, if necessary.

253. The *order* of a surd is indicated by the root symbol, or surd index. Thus $\sqrt[3]{x}$, $\sqrt[n]{a}$ are respectively surds of the third and n^{th} orders.

The surds of the most frequent occurrence are those of the second order; they are sometimes called **quadratic surds**. Thus $\sqrt{3}$, $\sqrt{a}$, $\sqrt{x+y}$ are quadratic surds.

254. It will frequently be found convenient to express a rational quantity in a surd form.

A rational quantity may be expressed in the form of a surd of *any required order* by raising it to the power whose root the surd expresses, and prefixing the radical sign. Thus

$$5 = \sqrt{25} = \sqrt[3]{125} = \sqrt[4]{625} = \sqrt[n]{5^n};$$

$$a+x = \sqrt{(a+x)^2} = \sqrt[3]{(a+x)^3} = \sqrt[n]{(a+x)^n}.$$

255. A surd of any order may be transformed into a surd of a different order.

Examples. (1) $\sqrt[3]{2} = 2^{\frac{1}{3}} = 2^{\frac{4}{12}} = \sqrt[12]{2^4}.$

(2) $\sqrt[n]{a} = a^{\frac{1}{n}} = a^{\frac{q}{nq}} = \sqrt[nq]{a^q}.$

256. Surds of different orders may be transformed into surds of the same order. This order may be *any* common multiple of each of the given orders, but it is usually most convenient to choose the *least* common multiple.

Example. Express $\sqrt[4]{a^3}$, $\sqrt[3]{b^2}$, $\sqrt[6]{c^5}$ as surds of the same lowest order.

The least common multiple of 4, 3, 6 is 12; and expressing the given surds as surds of the twelfth order they become $\sqrt[12]{a^9}$, $\sqrt[12]{b^8}$, $\sqrt[12]{c^{10}}$.

257. Surds of different orders may be arranged according to magnitude by transforming them into surds of the same order.

Example. Arrange $\sqrt{3}$, $\sqrt[3]{6}$, $\sqrt[4]{10}$ according to magnitude.

The least common multiple of 2, 3, 4 is 12; and, expressing the given surds as surds of the twelfth order, we have

$$\sqrt{3} = \sqrt[12]{3^6} = \sqrt[12]{729},$$

$$\sqrt[3]{6} = \sqrt[12]{6^4} = \sqrt[12]{1296},$$

$$\sqrt[4]{10} = \sqrt[12]{10^3} = \sqrt[12]{1000}.$$

Hence arranged in ascending order of magnitude the surds are

$$\sqrt{3}, \sqrt[4]{10}, \sqrt[3]{6}.$$

EXAMPLES XXXI. a.

Express as surds of the twelfth order with positive indices :

1. $x^{\frac{1}{2}}$.

2. $a^{-1} \div a^{-\frac{1}{2}}$.

3. $\sqrt[4]{ax^3} \times \sqrt[3]{a^{-1}x^{-2}}$.

4. $\frac{1}{a^{-\frac{1}{2}}}$.

5. $\frac{1}{\sqrt[8]{a^{-14}}}$.

6. $\sqrt[6]{\frac{1}{a^{-3}}}$.

Express as surds of the n^{th} order with positive indices :

7. $\sqrt[3]{x^2}$.

8. x^a .

9. $a^{\frac{1}{2}}$.

10. $\sqrt{a^{-\frac{1}{n}}}$.

11. $\sqrt[3]{x^ny^{\frac{1}{n}}}$.

12. $\frac{1}{a^{-1}}$.

13. $\frac{x^{-\frac{1}{2}}}{y^2}$.

14. $\frac{a^{\frac{1}{2}}}{a^{-n}}$.

Express as surds of the same lowest order :

15. $\sqrt{a}, \sqrt[9]{a^5}$.

16. $\sqrt[5]{a^3}, \sqrt{a}$.

17. $\sqrt[8]{x^2}, \sqrt[9]{x^6}, \sqrt[20]{x^5}$.

18. $\sqrt[10]{x^4}, \sqrt[12]{x^{10}}$.

19. $\sqrt[21]{a^5b^4}, \sqrt[7]{ab}$.

20. $\sqrt{ax^2}, \sqrt[39]{a^9x^6}$.

21. $\sqrt[4]{5}, \sqrt[3]{11}, \sqrt[9]{13}$.

22. $\sqrt[4]{8}, \sqrt{3}, \sqrt[8]{6}$.

23. $\sqrt[3]{2}, \sqrt[9]{8}, \sqrt[6]{4}$.

258. The root of any expression is equal to the product of the roots of the separate factors of the expression.

For

$$\sqrt[n]{ab} = (ab)^{\frac{1}{n}}$$

$$= a^{\frac{1}{n}} b^{\frac{1}{n}},$$

[Art. 244.]

$$= \sqrt[n]{a} \cdot \sqrt[n]{b}.$$

Similarly,

$$\sqrt[n]{abc} = \sqrt[n]{a} \cdot \sqrt[n]{b} \cdot \sqrt[n]{c};$$

and so for any number of factors.

Examples. (1) $\sqrt[4]{15} = \sqrt[4]{3} \cdot \sqrt[4]{5}$.

(2) $\sqrt[3]{a^6b} = \sqrt[3]{a^6} \cdot \sqrt[3]{b} = a^2 \sqrt[3]{b}$.

(3) $\sqrt{50} = \sqrt{25} \cdot \sqrt{2} = 5\sqrt{2}$.

Hence it appears that a surd may sometimes be expressed as the product of a rational quantity and a surd ; when so reduced the surd is said to be in its *simplest form*.Thus the simplest form of $\sqrt[4]{128}$ is $8\sqrt[4]{2}$.

Conversely, the coefficient of a surd may be brought under the radical sign by first reducing it to the form of a surd, and then multiplying the surds together.

Examples. (1) $7\sqrt{5} = \sqrt{49} \cdot \sqrt{5} = \sqrt{245}$.

(2) $a^{\frac{5}{2}}/b = \sqrt[5]{a^5} \cdot \sqrt[2]{b} = \sqrt[10]{a^5b}$.

When so reduced a surd is said to be an *entire surd*.

259. When surds have, or can be reduced to have, the same irrational factor, they are said to be *like*; otherwise, they are said to be *unlike*. Thus

$5\sqrt{3}$, $2\sqrt{3}$, $\frac{1}{5}\sqrt{3}$ are like surds.

But $3\sqrt{2}$ and $2\sqrt{3}$ are unlike surds.

Again, $3\sqrt{20}$, $4\sqrt{5}$, $\sqrt{\frac{1}{5}}$ are like surds;

for $3\sqrt{20} = 3\sqrt{4} \cdot \sqrt{5} = 3 \cdot 2\sqrt{5} = 6\sqrt{5}$;

and $\sqrt{\frac{1}{5}} = \sqrt{\frac{5}{25}} = \frac{1}{5}\sqrt{5}$.

260. In finding the sum of a number of like surds we reduce them to their simplest form, and prefix to their common irrational part the sum of the coefficients.

Example 1. The sum of $3\sqrt{20}$, $4\sqrt{5}$, $\frac{1}{5}\sqrt{5}$

$$= 6\sqrt{5} + 4\sqrt{5} + \frac{1}{5}\sqrt{5}$$

$$= \frac{51}{5}\sqrt{5}.$$

Example 2. The sum of $x^{\frac{3}{2}}\sqrt{8x^3a} + y^{\frac{3}{2}}\sqrt{-y^3a} - z^{\frac{3}{2}}\sqrt{x^3a}$

$$= x \cdot 2x^{\frac{3}{2}}\sqrt{a} + y(-y)^{\frac{3}{2}}\sqrt{a} - z \cdot z^{\frac{3}{2}}\sqrt{a}$$

$$= (2x^3 - y^2 - z^2)^{\frac{3}{2}}\sqrt{a}.$$

261. Unlike surds cannot be collected.

Thus the sum of $5\sqrt{2}$, $-2\sqrt{3}$ and $\sqrt{6}$ is $5\sqrt{2} - 2\sqrt{3} + \sqrt{6}$, and cannot be further simplified.

EXAMPLES XXXI b.

Express in the simplest form :

- | | | | |
|-----------------------|-------------------------|-------------------------|----------------------------|
| 1. $\sqrt{288}$. | 2. $\sqrt{147}$. | 3. $\sqrt[3]{256}$. | 4. $\sqrt[3]{432}$. |
| 5. $3\sqrt{150}$. | 6. $2\sqrt{720}$. | 7. $5\sqrt{245}$. | 8. $\sqrt[3]{1029}$. |
| 9. $\sqrt[3]{3125}$. | 10. $\sqrt[3]{-2187}$. | 11. $\sqrt[3]{96a^3}$. | 12. $\sqrt[3]{27a^3b^5}$. |

13. $\sqrt[3]{-108x^4y^3}$.

14. $\sqrt[n]{x^{3m}y^{2n+5}}$.

15. $\sqrt[n]{x^{a+py}y^{2p}}$.

16. $\sqrt{a^3+2a^2b+ab^2}$.

17. $\sqrt[3]{8x^4y-24x^3y^2+24x^2y^3-8xy^4}$.

Express as entire surds :

18. $11\sqrt{2}$.

19. $14\sqrt{5}$.

20. $6\sqrt[3]{4}$.

21. $5\sqrt[3]{6}$.

22. $\frac{4}{11}\sqrt{\frac{77}{8}}$.

23. $\frac{3ab}{2c}\sqrt{\frac{20c^2}{9a^2b}}$.

24. $\frac{3x}{y}\sqrt{\frac{a^2y^3}{x^2}}$.

25. $\frac{a}{x^2}\sqrt{\frac{3x^3}{a}}$.

26. $\frac{2a}{3x}\sqrt{\frac{27x^4}{a^2}}$.

27. $\frac{2a}{b}\sqrt[4]{\frac{b^4}{8a^3}}$.

28. $a\sqrt[n]{\frac{b^2}{a^{n-2}}}$.

29. $\frac{a}{b}\sqrt[p]{\frac{b^{p+1}}{a^{p-1}}}$.

30. $\frac{y}{x^n}\sqrt{\frac{x^{3n+1}}{y^3}}$.

31. $(x+y)\sqrt{\frac{x-y}{x+y}}$.

32. $\frac{ax}{a-x}\sqrt{\frac{a^2-x^2}{a^2x^2}}$.

Find the value of

33. $3\sqrt{45}-\sqrt{20}+7\sqrt{5}$.

34. $4\sqrt{63}+5\sqrt{7}-8\sqrt{28}$.

35. $\sqrt{44}-5\sqrt{176}+2\sqrt{99}$.

36. $2\sqrt{363}-5\sqrt{243}+\sqrt{192}$.

37. $2\sqrt[3]{189}+3\sqrt[3]{875}-7\sqrt[3]{56}$.

38. $5\sqrt[3]{81}-7\sqrt[3]{192}+4\sqrt[3]{648}$.

39. $3\sqrt[4]{162}-7\sqrt[4]{32}+\sqrt[4]{1250}$.

40. $5\sqrt[3]{-54}-2\sqrt[3]{-16}+4\sqrt[3]{686}$.

41. $4\sqrt{128}+4\sqrt{75}-5\sqrt{162}$.

42. $5\sqrt{24}-2\sqrt{54}-\sqrt{6}$.

43. $\sqrt{252}-\sqrt{294}-48\sqrt{\frac{1}{6}}$.

44. $3\sqrt{147}-\frac{7}{3}\sqrt{\frac{1}{3}}-\sqrt{\frac{1}{27}}$.

262. To multiply two surds of the same order: multiply separately the rational factors and the irrational factors.

For $a\sqrt[n]{x} \times b\sqrt[n]{y} = ax^{\frac{1}{n}} \times by^{\frac{1}{n}}$

$$= abx^{\frac{1}{n}}y^{\frac{1}{n}}$$

$$= ab(xy)^{\frac{1}{n}}$$

$$= ab\sqrt[n]{xy}.$$

Examples. (1) $5\sqrt{3} \times 3\sqrt{7} = 15\sqrt{21}$.

(2) $2\sqrt{x} \times 3\sqrt{x} = 6x$.

(3) $\sqrt[4]{a+b} \times \sqrt[4]{a-b} = \sqrt[4]{(a+b)(a-b)} = \sqrt[4]{a^2-b^2}$.

E.A.

R

263. If the surds are not in their simplest form, it will save labour to reduce them to this form before multiplication.

Example. The product of $5\sqrt{32}$, $\sqrt[4]{48}$, $2\sqrt{54}$

$$= 5 \cdot 4\sqrt{2} \times 4\sqrt{3} \times 2 \cdot 3\sqrt{6} = 480 \cdot \sqrt{2} \cdot \sqrt{3} \cdot \sqrt{6} = 480 \times 6 = 2880.$$

264. To multiply surds which are not of the same order: reduce them to equivalent surds of the same order, and proceed as before.

Example. Multiply $5\sqrt[3]{2}$ by $2\sqrt{5}$.

$$\text{The product} = 5\sqrt[3]{2^2} \times 2\sqrt[3]{5^3} = 10\sqrt[3]{2^2 \times 5^3} = 10\sqrt[3]{500}.$$

265. Suppose it is required to find the numerical value of the quotient when $\sqrt{5}$ is divided by $\sqrt{7}$.

At first sight it would seem that we must find the square root of 5, which is 2.236 ..., and then the square root of 7, which is 2.645 ..., and finally divide 2.236 ... by 2.645 ...; three troublesome operations.

But we may avoid much of this labour by multiplying both numerator and denominator by $\sqrt{7}$, so as to make the denominator a rational quantity. Thus

$$\frac{\sqrt{5}}{\sqrt{7}} = \frac{\sqrt{5}}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} = \frac{\sqrt{5 \times 7}}{7} = \frac{\sqrt{35}}{7}.$$

Now

$$\sqrt{35} = 5.916 \dots$$

$$\therefore \frac{\sqrt{5}}{\sqrt{7}} = \frac{5.916 \dots}{7} = .845 \dots$$

266. The great utility of this artifice in calculating the numerical value of surd fractions suggests its convenience in the case of *all* surd fractions, even where numerical values are not required. Thus it is usual to simplify $\frac{a\sqrt{b}}{\sqrt{c}}$ as follows:

$$\frac{a\sqrt{b}}{\sqrt{c}} = \frac{a\sqrt{b} \times \sqrt{c}}{\sqrt{c} \times \sqrt{c}} = \frac{a\sqrt{bc}}{c}.$$

The process by which surds are removed from the denominator of any fraction is known as **rationalising the denominator**. It is effected by multiplying both numerator and denominator by any factor which renders the denominator rational. We shall return to this point in Art. 270.

n, it will save
cation.

$\times 6 = 2880$.

same order;
and proceed as

rical value of

d the square
ot of 7, which
three trouble.

ultiplying both
e the denomi-

culating the
nience in the
values are not

ws :

the denomi-
enominator.
denominator
al. We shall

267. The quotient of one surd by another may be found by expressing the result as a fraction, and rationalising the denominator.

Example 1. Divide $4\sqrt{75}$ by $25\sqrt{56}$.

$$\begin{aligned}\text{The quotient} &= \frac{4\sqrt{75}}{25\sqrt{56}} = \frac{4 \times 5\sqrt{3}}{25 \times 2\sqrt{14}} = \frac{2\sqrt{3}}{5\sqrt{14}} \\ &= \frac{2\sqrt{3} \times \sqrt{14}}{5\sqrt{14} \times \sqrt{14}} = \frac{2\sqrt{42}}{5 \times 14} = \frac{\sqrt{42}}{35}.\end{aligned}$$

$$\text{Example 2. } \frac{\sqrt[3]{b}}{\sqrt[3]{c^2}} = \frac{\sqrt[3]{b} \times \sqrt[3]{c}}{\sqrt[3]{c^2} \times \sqrt[3]{c}} = \frac{\sqrt[3]{bc}}{\sqrt[3]{c^3}} = \frac{\sqrt[3]{bc}}{c}.$$

EXAMPLES XXXI. c.

Find the value of

1. $2\sqrt{14} \times \sqrt{21}$.
2. $3\sqrt{8} \times \sqrt{6}$.
3. $5\sqrt{a} \times 2\sqrt{3}$.
4. $2\sqrt{15} \times 3\sqrt{5}$.
5. $8\sqrt{12} \times 3\sqrt{24}$.
6. $\sqrt[3]{x+2} \times \sqrt[3]{x-2}$.
7. $21\sqrt{384} \div 8\sqrt{98}$.
8. $5\sqrt{27} \div 3\sqrt{24}$.
9. $-13\sqrt{125} \div 5\sqrt{65}$.
10. $\sqrt[3]{168} \times \sqrt[3]{147}$.
11. $5\sqrt[3]{128} \times 2\sqrt[3]{432}$.
12. $6\sqrt{14} \div 2\sqrt{21}$.
13. $a\sqrt{b^3} \times b^2\sqrt{a}$.
14. $\frac{3\sqrt{11}}{2\sqrt{98}} \div \frac{5}{7\sqrt{22}}$.
15. $\frac{3\sqrt{48}}{5\sqrt{112}} \div \frac{6\sqrt{84}}{\sqrt{392}}$.
16. $\frac{3}{x}\sqrt{\frac{a^3}{x}} \times \frac{4}{3}\sqrt{\frac{x^3}{2a^4}}$.
17. $\frac{3}{a-b}\sqrt{\frac{2x}{a-b}} \div \sqrt{\frac{18x^3}{(a-b)^5}}$.

Given $\sqrt{2} = 1.41421$, $\sqrt{3} = 1.73205$, $\sqrt{5} = 2.23607$, $\sqrt{6} = 2.44949$, $\sqrt{7} = 2.64575$: find to four places of decimals the numerical value of

18. $\frac{14}{\sqrt{2}}$.
19. $\frac{25}{\sqrt{5}}$.
20. $\frac{10}{\sqrt{7}}$.
21. $\frac{48}{\sqrt{6}}$.
22. $\frac{60}{\sqrt{5}}$.
23. $144 \div \sqrt{6}$.
24. $\sqrt{2} \div \sqrt{3}$.
25. $\frac{1}{2\sqrt{3}}$.
26. $\frac{1}{\sqrt{500}}$.
27. $\frac{4}{\sqrt{243}}$.
28. $\frac{25}{\sqrt{252}}$.
29. $\sqrt[3]{\frac{256}{1575}}$.

268. Hitherto we have confined our attention to **simple surds**, such as $\sqrt[3]{5}$, $\sqrt[3]{a}$, $\sqrt{x+y}$. An expression involving two or more simple surds is called a **compound surd**; thus $2\sqrt{a} - 3\sqrt{b}$; $\sqrt[3]{a} + \sqrt[3]{b}$ are compound surds.

269. The multiplication of compound surds is performed like the multiplication of compound algebraical expressions.

Example 1. Multiply $2\sqrt{x}-5$ by $3\sqrt{x}$.

$$\begin{aligned}\text{The product} &= 3\sqrt{x}(2\sqrt{x}-5) \\ &= 6x - 15\sqrt{x}.\end{aligned}$$

Example 2. Multiply $2\sqrt{5}+3\sqrt{x}$ by $\sqrt{5}-\sqrt{x}$.

$$\begin{aligned}\text{The product} &= (2\sqrt{5}+3\sqrt{x})(\sqrt{5}-\sqrt{x}) \\ &= 2\sqrt{5} \cdot \sqrt{5} + 3\sqrt{5} \cdot \sqrt{x} - 2\sqrt{5} \cdot \sqrt{x} - 3\sqrt{x} \cdot \sqrt{x} \\ &= 10 - 3x + \sqrt{5}x.\end{aligned}$$

Example 3. Find the square of $2\sqrt{x}+\sqrt{7-4x}$.

$$\begin{aligned}(2\sqrt{x}+\sqrt{7-4x})^2 &= (2\sqrt{x})^2 + (\sqrt{7-4x})^2 + 4\sqrt{x} \cdot \sqrt{7-4x} \\ &= 4x + 7 - 4x + 4\sqrt{7x-4x^2} \\ &= 7 + 4\sqrt{7x-4x^2}.\end{aligned}$$

EXAMPLES XXXI. d.

Find the value of

- | | |
|--|---|
| 1. $(3\sqrt{x}-5) \times 2\sqrt{x}$. | 2. $(\sqrt{x}-\sqrt{a}) \times 2\sqrt{x}$. |
| 3. $(\sqrt{a}+\sqrt{b}) \times \sqrt{ab}$. | 4. $(\sqrt{x+y}-1) \times \sqrt{x+y}$. |
| 5. $(2\sqrt{3}+3\sqrt{2})^2$. | 6. $(\sqrt{7}+5\sqrt{3})(2\sqrt{7}-4\sqrt{3})$. |
| 7. $(3\sqrt{5}-4\sqrt{2})(2\sqrt{5}+3\sqrt{2})$. | 8. $(3\sqrt{a}-2\sqrt{x})(2\sqrt{a}+3\sqrt{x})$. |
| 9. $(\sqrt{x}+\sqrt{x-1}) \times \sqrt{x-1}$. | 10. $(\sqrt{x+a}-\sqrt{x-a}) \times \sqrt{x+a}$. |
| 11. $(\sqrt{a+x}-2\sqrt{a})^2$. | 12. $(2\sqrt{a}-\sqrt{1+4a})^2$. |
| 13. $(\sqrt{a+x}-\sqrt{a-x})^2$. | 14. $(\sqrt{a+x}-2)(\sqrt{a+x}-1)$. |
| 15. $(\sqrt{2}+\sqrt{3}-\sqrt{5})(\sqrt{2}+\sqrt{3}+\sqrt{5})$. | |
| 16. $(\sqrt{5}+3\sqrt{2}+\sqrt{7})(\sqrt{5}+3\sqrt{2}-\sqrt{7})$. | |

Write down the square of

- | | |
|-----------------------------------|---|
| 17. $\sqrt{2x+a}-\sqrt{2x-a}$. | 18. $\sqrt{x^2-2y^2}+\sqrt{x^2+2y^2}$. |
| 19. $\sqrt{m+n}+\sqrt{m-n}$. | 20. $3\sqrt{a^2+b^2}-2\sqrt{a^2-b^2}$. |
| 21. $3x\sqrt{2}-3\sqrt{7-2x^2}$. | 22. $\sqrt{4x^2+1}-\sqrt{4x^2-1}$. |

is performed
expressions.

270. One case of the multiplication of compound surds deserves careful attention. For if we multiply together the sum and the difference of any two quadratic surds we obtain a rational product.

Examples. (1) $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) = (\sqrt{a})^2 - (\sqrt{b})^2 = a - b$.

(2) $(3\sqrt{5} + 4\sqrt{3})(3\sqrt{5} - 4\sqrt{3}) = (3\sqrt{5})^2 - (4\sqrt{3})^2 = 45 - 48 = -3$.

Similarly, $(4 - \sqrt{a+b})(4 + \sqrt{a+b}) = (4)^2 - (\sqrt{a+b})^2 = 16 - a - b$.

271. DEFINITION. When two binomial quadratic surds differ only in the sign which connects their terms they are said to be **conjugate**.

Thus $3\sqrt{7} + 5\sqrt{11}$ is conjugate to $3\sqrt{7} - 5\sqrt{11}$.

Similarly, $a - \sqrt{a^2 - x^2}$ is conjugate to $a + \sqrt{a^2 - x^2}$.

The product of two conjugate surds is rational. [Art. 270.]

Example. $(3\sqrt{a} + \sqrt{x-9a})(3\sqrt{a} - \sqrt{x-9a})$
 $= (3\sqrt{a})^2 - (\sqrt{x-9a})^2 = 9a - (x-9a) = 18a - x$.

272. The only case of the division of compound surds which we shall here consider is that in which the divisor is a binomial quadratic surd. If we express the division by means of a fraction, we can always rationalise the denominator by multiplying numerator and denominator by the surd which is conjugate to the divisor.

Example 1. Divide $4 + 3\sqrt{2}$ by $5 - 3\sqrt{2}$.

$$\begin{aligned} \text{The quotient} &= \frac{4 + 3\sqrt{2}}{5 - 3\sqrt{2}} = \frac{4 + 3\sqrt{2}}{5 - 3\sqrt{2}} \times \frac{5 + 3\sqrt{2}}{5 + 3\sqrt{2}} \\ &= \frac{20 + 18 + 12\sqrt{2} + 15\sqrt{2}}{25 - 18} = \frac{38 + 27\sqrt{2}}{7} \end{aligned}$$

Example 2. Rationalise the denominator of $\frac{b^3}{\sqrt{a^2 + b^2} + a}$.

$$\begin{aligned} \text{The expression} &= \frac{b^3}{\sqrt{a^2 + b^2} + a} \times \frac{\sqrt{a^2 + b^2} - a}{\sqrt{a^2 + b^2} - a} \\ &= \frac{b^3 \{ \sqrt{a^2 + b^2} - a \}}{(a^2 + b^2) - a^2} \\ &= \frac{b^3 \sqrt{a^2 + b^2} - ab^3}{b^2} \end{aligned}$$

Example 3. Divide $\frac{\sqrt{3}+\sqrt{2}}{2-\sqrt{3}}$ by $\frac{7+4\sqrt{3}}{\sqrt{3}-\sqrt{2}}$.

$$\begin{aligned}\text{The quotient} &= \frac{\sqrt{3}+\sqrt{2}}{2-\sqrt{3}} \times \frac{\sqrt{3}-\sqrt{2}}{7+4\sqrt{3}} \\ &= \frac{(\sqrt{3})^2 - (\sqrt{2})^2}{14 - 12 + 8\sqrt{3} - 7\sqrt{3}} \\ &= \frac{1}{2+\sqrt{3}} \\ &= 2-\sqrt{3}, \text{ on rationalising.}\end{aligned}$$

Example 4. Given $\sqrt{5} = 2.236068$, find the value of $\frac{87}{7-2\sqrt{5}}$.

Rationalising the denominator,

$$\frac{87}{7-2\sqrt{5}} = \frac{87(7+2\sqrt{5})}{49-20} = 3(7+2\sqrt{5}) = 34.416408.$$

It will be seen that by rationalising the denominator we have avoided the use of a divisor consisting of 7 figures.

EXAMPLES XXXI. e.

Find the value of

1. $(9\sqrt{2}-7)(9\sqrt{2}+7).$
2. $(3+5\sqrt{7})(3-5\sqrt{7}).$
3. $(5\sqrt{8}-2\sqrt{7})(5\sqrt{8}+2\sqrt{7}).$
4. $(2\sqrt{11}+5\sqrt{2})(2\sqrt{11}-5\sqrt{2}).$
5. $(\sqrt{a}+2\sqrt{b})(\sqrt{a}-2\sqrt{b}).$
6. $(3c-2\sqrt{x})(3c+2\sqrt{x}).$
7. $(\sqrt{a+x}-\sqrt{a})(\sqrt{a+x}+\sqrt{a}).$
8. $(\sqrt{2p+3q}-2\sqrt{q})(\sqrt{2p+3q}+2\sqrt{q}).$
9. $(\sqrt{a+x}+\sqrt{a-x})(\sqrt{a+x}-\sqrt{a-x}).$
10. $(5\sqrt{x^2-3y^2}+7a)(5\sqrt{x^2-3y^2}-7a).$
11. $29 \div (11+3\sqrt{7}).$
12. $17 \div (3\sqrt{7}+2\sqrt{3}).$
13. $(3\sqrt{2}-1) \div (3\sqrt{2}+1).$
14. $(2\sqrt{3}+7\sqrt{2}) \div (5\sqrt{3}-4\sqrt{2}).$
15. $(2x-\sqrt{xy}) \div (2\sqrt{xy}-y).$
16. $(3+\sqrt{5})(\sqrt{5}-2) \div (5-\sqrt{5}).$
17. $\frac{\sqrt{a}}{\sqrt{a}-\sqrt{x}} \div \frac{\sqrt{a}+\sqrt{x}}{\sqrt{x}}.$
18. $\frac{2\sqrt{15}+8}{5+\sqrt{15}} \div \frac{8\sqrt{3}-6\sqrt{5}}{5\sqrt{3}-3\sqrt{5}}.$

Rationalise the denominator of

19. $\frac{25\sqrt{3}-4\sqrt{2}}{7\sqrt{3}-5\sqrt{2}}.$
20. $\frac{10\sqrt{6}-2\sqrt{7}}{3\sqrt{6}+2\sqrt{7}}.$
21. $\frac{\sqrt{7}+\sqrt{2}}{9+2\sqrt{14}}.$

22. $\frac{2\sqrt{3}+3\sqrt{2}}{5+2\sqrt{6}}$

23. $\frac{y^2}{x+\sqrt{x^2-y^2}}$

24. $\frac{x^2}{\sqrt{x^2+a^2}+a}$

25. $\frac{\sqrt{1+x^2}-\sqrt{1-x^2}}{\sqrt{1+x^2}+\sqrt{1-x^2}}$

26. $\frac{2\sqrt{a+b}+3\sqrt{a-b}}{2\sqrt{a+b}-\sqrt{a-b}}$

27. $\frac{\sqrt{9+x^2}-3}{\sqrt{9+x^2}+3}$

28. $\frac{3+\sqrt{6}}{5\sqrt{3}-2\sqrt{12}-\sqrt{32}+\sqrt{50}}$

Given $\sqrt{2}=1.41421$, $\sqrt{3}=1.73205$, $\sqrt{5}=2.23607$: find to four places of decimals the value of

29. $\frac{1}{2+\sqrt{3}}$

30. $\frac{3+\sqrt{5}}{\sqrt{5}-2}$

31. $\frac{\sqrt{5}+\sqrt{3}}{4+\sqrt{15}}$

32. $\frac{\sqrt{5}-2}{9-4\sqrt{5}}$

33. $\frac{7\sqrt{5}+15}{\sqrt{5}-1} \times \frac{\sqrt{5}-2}{3+\sqrt{5}}$

34. $(2-\sqrt{3})(7-4\sqrt{3}) \div (3\sqrt{3}-5)$

273. *The square root of a rational quantity cannot be partly rational and partly a quadratic surd.*

If possible let
then by squaring,

$$\sqrt{n}=a+\sqrt{m};$$

$$n=a^2+m+2a\sqrt{m};$$

$$\therefore \sqrt{m}=\frac{n-a^2-m}{2a};$$

that is a surd is equal to a rational quantity; which is impossible.

274. *If $x+\sqrt{y}=a+\sqrt{b}$, then will $x=a$ and $y=b$.*

For if x is not equal to a , let $x=a+m$; then

$$a+m+\sqrt{y}=a+\sqrt{b};$$

$$\sqrt{b}=m+\sqrt{y};$$

that is,
which is impossible.

[Art. 273.]

Therefore

$$x=a,$$

and consequently,

$$y=b.$$

If therefore

$$x+\sqrt{y}=a+\sqrt{b},$$

we must also have

$$x-\sqrt{y}=a-\sqrt{b}.$$

275. It appears from the preceding article that in any equation of the form

$$X+\sqrt{Y}=A+\sqrt{B} \dots\dots\dots (1),$$

we may equate the rational parts on each side, and also the irrational parts; so that the equation (1) is really equivalent to two independent equations, $X=A$, and $Y=B$.

276. If $\sqrt{a+\sqrt{b}}=\sqrt{x}+\sqrt{y}$ then will $\sqrt{a-\sqrt{b}}=\sqrt{x}-\sqrt{y}$.

For by squaring, we obtain

$$a+\sqrt{b}=x+2\sqrt{xy}+y;$$

$$\therefore a=x+y, \sqrt{b}=2\sqrt{xy}. \quad [\text{Art. 275.}]$$

Hence

$$a-\sqrt{b}=x-2\sqrt{xy}+y,$$

and

$$\sqrt{a-\sqrt{b}}=\sqrt{x}-\sqrt{y}.$$

277. To find the square root of $a+\sqrt{b}$.

Suppose $\sqrt{a+\sqrt{b}}=\sqrt{x}+\sqrt{y}$;
then as in the last article,

$$x+y=a \dots\dots\dots(1),$$

$$2\sqrt{xy}=\sqrt{b} \dots\dots\dots(2).$$

$$\therefore (x-y)^2=(x+y)^2-4xy \\ =a^2-b, \quad \text{from (1) and (2).}$$

$$\therefore x-y=\sqrt{a^2-b}.$$

Combining this with (1) we find

$$x=\frac{a+\sqrt{a^2-b}}{2}, \text{ and } y=\frac{a-\sqrt{a^2-b}}{2}$$

$$\therefore \sqrt{a+\sqrt{b}}=\sqrt{\frac{a+\sqrt{a^2-b}}{2}}+\sqrt{\frac{a-\sqrt{a^2-b}}{2}}.$$

278. From the values just found for x and y , it appears that each of them is itself a compound surd unless a^2-b is a perfect square. Hence the method of Art. 277 for finding the square root of $a+\sqrt{b}$ is of no practical utility except when a^2-b is a perfect square.

Example. Find the square root of $16+2\sqrt{55}$.

Assume $\sqrt{16+2\sqrt{55}}=\sqrt{x}+\sqrt{y}$.

Then

$$16+2\sqrt{55}=x+2\sqrt{xy}+y.$$

$$\therefore x+y=16 \dots\dots\dots(1),$$

$$2\sqrt{xy}=2\sqrt{55} \dots\dots\dots(2).$$

$$\therefore (x-y)^2=(x+y)^2-4xy \\ =16^2-4 \times 55, \quad \text{by (1) and (2).} \\ =4 \times 9.$$

$$\therefore x-y=\pm 6 \dots\dots\dots(3).$$

From (1) and (3) we obtain

$$x=11, \text{ or } 5, \text{ and } y=5, \text{ or } 11.$$

That is, the required square root is $\sqrt{11}+\sqrt{5}$.

$\sqrt{x} - \sqrt{y}$.

[Art. 275.]

In the same way we may shew that

$$\sqrt{16 - 2\sqrt{55}} = \sqrt{11} - \sqrt{5}.$$

Note. Since every quantity has two square roots equal in magnitude but opposite in sign, strictly speaking we should have

$$\begin{aligned} \text{the square root of } 16 + 2\sqrt{55} &= \pm(\sqrt{11} + \sqrt{5}), \\ \dots\dots\dots 16 - 2\sqrt{55} &= \pm(\sqrt{11} - \sqrt{5}). \end{aligned}$$

However it is usually sufficient to take the positive value of the square root, so that in assuming $\sqrt{a} - \sqrt{b} = \sqrt{x} - \sqrt{y}$ it is understood that x is greater than y . With this proviso it will be unnecessary in any numerical example to use the double sign at the stage of work corresponding to equation (3) of the last example.

279. When the binomial whose square root we are seeking consists of *two* quadratic surds, we proceed as explained in the following example.

Example. Find the square root of $\sqrt{175} - \sqrt{147}$.

$$\text{Since } \sqrt{175} - \sqrt{147} = \sqrt{7}(\sqrt{25} - \sqrt{21}) = \sqrt{7}(5 - \sqrt{21}).$$

$$\therefore \sqrt{\sqrt{175} - \sqrt{147}} = \sqrt[4]{7} \cdot \sqrt{5 - \sqrt{21}}.$$

And, proceeding as in the last article,

$$\sqrt{5 - \sqrt{21}} = \sqrt{\frac{7}{2}} - \sqrt{\frac{3}{2}};$$

$$\therefore \sqrt{\sqrt{175} - \sqrt{147}} = \sqrt[4]{7} \left(\sqrt{\frac{7}{2}} - \sqrt{\frac{3}{2}} \right).$$

280. The square root of a binomial surd may often be found by inspection.

Example 1. Find the square root of $11 + 2\sqrt{30}$.

We have only to find two quantities whose sum is 11, and whose product is 30; thus

$$\begin{aligned} 11 + 2\sqrt{30} &= 6 + 5 + 2\sqrt{6 \times 5} \\ &= (\sqrt{6} + \sqrt{5})^2. \end{aligned}$$

$$\therefore \sqrt{11 + 2\sqrt{30}} = \sqrt{6} + \sqrt{5}.$$

Example 2. Find the square root of $53 - 12\sqrt{10}$.

First write the binomial so that the surd part has a coefficient 2; thus

$$53 - 12\sqrt{10} = 53 - 2\sqrt{360}.$$

We have now to find two quantities whose sum is 53 and whose product is 360; these are 45 and 8;

hence

$$\begin{aligned}
 53 - 12\sqrt{10} &= 45 + 8 - 2\sqrt{45 \times 8} \\
 &= (\sqrt{45} - \sqrt{8})^2; \\
 \therefore \sqrt{53 - 12\sqrt{10}} &= \sqrt{45} - \sqrt{8} \\
 &= 3\sqrt{5} - 2\sqrt{2}.
 \end{aligned}$$

EXAMPLES XXXI. f.

Find the square root of each of the following binomial surds :

- | | | |
|---------------------------------|--|-------------------------------|
| 1. $7 - 2\sqrt{10}$. | 2. $13 + 2\sqrt{30}$. | 3. $8 - 2\sqrt{7}$. |
| 4. $5 + 2\sqrt{6}$. | 5. $75 + 12\sqrt{21}$. | 6. $18 - 8\sqrt{5}$. |
| 7. $41 - 24\sqrt{2}$. | 8. $83 + 12\sqrt{35}$. | 9. $47 - 4\sqrt{33}$. |
| 10. $2\frac{1}{4} + \sqrt{5}$. | 11. $4\frac{1}{3} - \frac{4}{3}\sqrt{3}$. | 12. $16 + 5\sqrt{7}$. |
| 13. $\sqrt{27} + 2\sqrt{6}$. | 14. $\sqrt{32} - \sqrt{24}$. | 15. $3\sqrt{5} + \sqrt{40}$. |

Find the fourth roots of the following binomial surds :

- | | | |
|-------------------------|-------------------------|--|
| 16. $17 + 12\sqrt{2}$. | 17. $56 + 24\sqrt{5}$. | 18. $\frac{3}{2}\sqrt{5} + 3\frac{1}{2}$. |
| 19. $14 + 8\sqrt{3}$. | 20. $49 - 20\sqrt{6}$. | 21. $248 + 32\sqrt{60}$. |

Find, by inspection, the value of

- | | | |
|-------------------------------|--------------------------------|--------------------------------|
| 22. $\sqrt{3 - 2\sqrt{2}}$. | 23. $\sqrt{4 + 2\sqrt{3}}$. | 24. $\sqrt{6 - 2\sqrt{5}}$. |
| 25. $\sqrt{19 + 8\sqrt{3}}$. | 26. $\sqrt{8 + 2\sqrt{15}}$. | 27. $\sqrt{9 - 2\sqrt{14}}$. |
| 28. $\sqrt{11 + 4\sqrt{6}}$. | 29. $\sqrt{15 - 4\sqrt{14}}$. | 30. $\sqrt{29 + 6\sqrt{22}}$. |

Equations involving Surds.

281. Sometimes equations are proposed in which the unknown quantity appears under the radical sign. Such equations are very varied in character and often require special artifices for their solution. Here we shall only consider a few of the simpler cases, which can generally be solved by the following method. Bring to one side of the equation a single radical term by itself: on squaring both sides this radical will disappear. By repeating this process any remaining radicals can in turn be removed.

Example 1. Solve $2\sqrt{x} - \sqrt{4x-11} = 1$.
 Transposing $2\sqrt{x} - 1 = \sqrt{4x-11}$.
 Square both sides; then $4x - 4\sqrt{x} + 1 = 4x - 11$,
 $4\sqrt{x} = 12$,
 $\sqrt{x} = 3$;
 $\therefore x = 9$.

Example 2. Solve $2 + \sqrt[3]{x-5} = 13$.
 Transposing $\sqrt[3]{x-5} = 11$.
 Here we must *cube* both sides; thus $x - 5 = 1331$;
 whence $x = 1336$.

Example 3. Solve $\sqrt{x+5} + \sqrt{3x+4} = \sqrt{12x+1}$.
 Squaring both sides,
 $x + 5 + 3x + 4 + 2\sqrt{(x+5)(3x+4)} = 12x + 1$.
 Transposing and dividing by 2,
 $\sqrt{(x+5)(3x+4)} = 4x - 4$ (1).
 Squaring,
 $(x+5)(3x+4) = 16x^2 - 32x + 16$,
 or $13x^2 - 51x - 4 = 0$,
 $(x-4)(13x+1) = 0$;
 $\therefore x = 4$, or $-\frac{1}{13}$.

If we proceed to verify the solution by substituting these values in the original equation, it will be found that it is satisfied by $x=4$, but not by $x = -\frac{1}{13}$. But this latter value will be found on trial to satisfy the given equation if we alter the sign of the second radical; thus

$$\sqrt{x+5} - \sqrt{3x+4} = \sqrt{12x+1}.$$

On squaring this and reducing, we obtain

$$-\sqrt{(x+5)(3x+4)} = 4x - 4$$
(2);

and a comparison of (1) and (2) shews that in the next stage of the work the same quadratic equation is obtained in each case, the roots of which are 4 and $-\frac{1}{13}$, as already found.

From this it appears that when the solution of an equation requires that both sides should be squared, we cannot be certain without trial which of the values found for the unknown quantity will satisfy the original equation.

In order that all the values found by the solution of the equation may be applicable it will be necessary to take into account both signs of the radicals in the given equation.

EXAMPLES XXXI. g.

Solve the equations :

1. $\sqrt{x-5}=3$.
2. $\sqrt[3]{4x-7}=5$.
3. $7-\sqrt{x-4}=3$.
4. $13-\sqrt[3]{5x-4}=7$.
5. $\sqrt{5x-1}=2\sqrt{x+3}$.
6. $2\sqrt{3-7x}-3\sqrt{8x-12}=0$.
7. $2\sqrt[3]{5x-35}=5\sqrt[3]{2x-7}$.
8. $\sqrt{9x^2-11x-5}=3x-2$.
9. $\sqrt[3]{2x+11}=\sqrt{5}$.
10. $\sqrt{4x^2-7x+1}=2x-1\frac{1}{2}$.
11. $\sqrt{x+25}=1+\sqrt{x}$.
12. $\sqrt{8x+33}-3=2\sqrt{2x}$.
13. $\sqrt{x+3}+\sqrt{x}=5$.
14. $10-\sqrt{25+9x}=3\sqrt{x}$.
15. $\sqrt{x-4}+3=\sqrt{x+11}$.
16. $\sqrt{9x-8}=3\sqrt{x+4}-2$.
17. $\sqrt{4x+5}-\sqrt{x}=\sqrt{x+3}$.
18. $\sqrt{95x-29}-\sqrt{4x-11}=3\sqrt{x}$.
19. $\sqrt{8x+17}-\sqrt{2x}=\sqrt{2x+9}$.
20. $\sqrt{3x-11}+\sqrt{3x}=\sqrt{12x-23}$.
21. $\sqrt{12x-5}+\sqrt{3x-1}=\sqrt{27x-2}$.
22. $\sqrt{x+3}+\sqrt{x+8}-\sqrt{4x+21}=0$.
23. $\sqrt{x+2}+\sqrt{4x+1}-\sqrt{9x+7}=0$.
24. $\sqrt{x+4ab}=2a+\sqrt{x}$.
25. $\sqrt{x+\sqrt{4a+x}}=2\sqrt{b+x}$.
26. $\sqrt{a-x}+\sqrt{b+x}=\sqrt{a}+\sqrt{b}$.
27. $5\sqrt[3]{70x+29}=9\sqrt[3]{14x-15}$.
28. $\sqrt[3]{x^3-3x^2+7x-11}=x-1$.
29. $\sqrt[3]{8x^3+12x^2+12x-11}=2x+1$.
30. $\sqrt[3]{1+x}+\sqrt[3]{1-x}=\sqrt[3]{2}$.

282. When radicals appear in a fractional form in an equation, we must clear of fractions in the ordinary way, combining the irrational factors by the rules already explained in this chapter.

Example 1. Solve $\frac{6\sqrt{x-11}}{3\sqrt{x}} = \frac{2\sqrt{x+1}}{\sqrt{x+6}}$.

Multiplying across, we have

$$\begin{aligned}
 6x+25\sqrt{x}-66 &= 6x+3\sqrt{x}, \\
 \text{that is,} \quad 25\sqrt{x}-3\sqrt{x} &= 66, \\
 22\sqrt{x} &= 66, \\
 \sqrt{x} &= 3, \\
 x &= 9.
 \end{aligned}$$

the equation
account both

Example 2. Solve $\sqrt{9+2x} - \sqrt{2x} = \frac{5}{\sqrt{9+2x}}$.

Clearing of fractions, $9+2x - \sqrt{2x(9+2x)} = 5$,
 $4+2x = \sqrt{2x(9+2x)}$.

Squaring, $16+16x+4x^2 = 18+4x^2$,
 $16=2x$,
 $x=8$.

EXAMPLES XXXI. h.

Solve the equations :

1. $\frac{6\sqrt{x-21}}{3\sqrt{x-14}} = \frac{8\sqrt{x-11}}{4\sqrt{x-13}}$

2. $\frac{9\sqrt{x-23}}{3\sqrt{x-8}} = \frac{6\sqrt{x-17}}{2\sqrt{x-6}}$

3. $\frac{\sqrt{x+3}}{\sqrt{x-2}} = \frac{3\sqrt{x-5}}{3\sqrt{x-13}}$

4. $2 - \frac{\sqrt{x+3}}{\sqrt{x+2}} = \frac{\sqrt{x+9}}{\sqrt{x+7}}$

5. $\frac{2\sqrt{x-1}}{2\sqrt{x+3}} = \frac{\sqrt{x-2}}{\sqrt{x-3}}$

6. $\frac{6\sqrt{x-7}}{\sqrt{x-1}} - 5 = \frac{7\sqrt{x-26}}{7\sqrt{x-21}}$

7. $\frac{12\sqrt{x-11}}{4\sqrt{x-4\frac{2}{3}}} = \frac{6\sqrt{x+5}}{2\sqrt{x+3}}$

8. $\sqrt{1+x} + \sqrt{x} = \frac{2}{\sqrt{1+x}}$

9. $\sqrt{x-1} + \sqrt{x} = \frac{2}{\sqrt{x}}$

10. $\sqrt{x} - \sqrt{x-8} = \frac{2}{\sqrt{x-8}}$

11. $\sqrt{x+5} + \sqrt{x} = \frac{10}{\sqrt{x}}$

12. $2\sqrt{x} - \sqrt{4x-3} = \frac{1}{\sqrt{4x-3}}$

13. $3\sqrt{x} = \frac{8}{\sqrt{9x-32}} + \sqrt{9x-32}$

14. $\sqrt{x-7} = \frac{1}{\sqrt{x+7}}$

15. $(\sqrt{x+11})(\sqrt{x-11}) + 110 = 0$

16. $2\sqrt{x} = \frac{12-6\sqrt{x}}{2\sqrt{x-3}}$

17. $3\sqrt{x-1} = \frac{5}{3\sqrt{x+7}} + 6$

18. $\frac{x-1}{\sqrt{x-1}} = 3 + \frac{\sqrt{x+1}}{2}$

19. $\frac{1}{1-x} + \frac{1}{\sqrt{x+1}} + \frac{1}{\sqrt{x-1}} = 0$

20. $2 = \frac{\sqrt{2+x} + \sqrt{2-x}}{\sqrt{2+x} - \sqrt{2-x}}$

21. $\frac{2x-3}{\sqrt{x-2}+1} = 2\sqrt{x-2} - 1$


22. $\frac{2}{x-6+\sqrt{x}} + \frac{3}{\sqrt{x-2}} = \frac{4}{\sqrt{x+3}}$

[Further practice in surd equations will be found on pages 362, 363.]



MICROCOPY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482 - 0300 - Phone
(716) 288 - 5989 - Fax

CHAPTER XXXII.

RATIO, PROPORTION, AND VARIATION.

Ratio.

283. DEFINITION. **Ratio** is the relation which one quantity bears to another of the *same* kind, the comparison being made by considering what multiple, part, or parts, one quantity is of the other.

The ratio of A to B is usually written $A : B$. The quantities A and B are called the *terms* of the ratio. The first term is called the **antecedent**, the second term the **consequent**.

284. To find what multiple or part A is of B we divide A by B ; hence the ratio $A : B$ may be measured by the fraction $\frac{A}{B}$ and we shall usually find it convenient to adopt this notation.

In order to compare two quantities they must be expressed in terms of the same unit. Thus the ratio of £2 to 15s. is measured by the fraction $\frac{2 \times 20}{15}$ or $\frac{8}{3}$.

Note. Since a ratio expresses the *number* of times that one quantity contains another, *every ratio is an abstract quantity*.

285. By Art. 151, $\frac{a}{b} = \frac{ma}{mb}$;

and thus the ratio $a : b$ is equal to the ratio $ma : mb$; that is, *the value of a ratio remains unaltered if the antecedent and the consequent are multiplied or divided by the same quantity*.

286. Two or more ratios may be compared by reducing their equivalent fractions to a common denominator. Thus suppose $a : b$ and $x : y$ are two ratios. Now $\frac{a}{b} = \frac{ay}{by}$, and $\frac{x}{y} = \frac{bx}{by}$; hence the ratio $a : b$ is greater than, equal to, or less than the ratio $x : y$ according as ay is greater than, equal to, or less than bx .

287. The ratio of two fractions can be expressed as a ratio of two integers. Thus the ratio $\frac{a}{b} : \frac{c}{d}$ is measured by the fraction $\frac{\frac{a}{b}}{\frac{c}{d}}$, or $\frac{ad}{bc}$; and is therefore equivalent to the ratio $ad : bc$.

288. If either, or both, of the terms of a ratio be a surd quantity, then no two integers can be found which will *exactly* measure their ratio. Thus the ratio $\sqrt{2} : 1$ cannot be exactly expressed by any two integers.

289. DEFINITION. If the ratio of any two quantities can be expressed exactly by the ratio of two integers the quantities are said to be **commensurable**; otherwise, they are said to be **incommensurable**.

Although we cannot find two integers which will exactly measure the ratio of two incommensurable quantities, we can always find two integers whose ratio differs from that required by as small a quantity as we please.

Thus
$$\frac{\sqrt{5}}{4} = \frac{2.236067...}{4} = .559016...$$
and therefore
$$\frac{\sqrt{5}}{4} > \frac{559016}{1000000} \text{ and } < \frac{559017}{1000000},$$

and it is evident that by carrying the decimals further, any degree of approximation may be arrived at.

290. DEFINITION. Ratios are *compounded* by multiplying together the fractions which denote them; or by multiplying together the antecedents for a new antecedent, and the consequents for a new consequent.

Example. Find the ratio compounded of the three ratios

$$2a : 3b, 6ab : 5c^2, c : a.$$

$$\text{The required ratio} = \frac{2a}{3b} \times \frac{6ab}{5c^2} \times \frac{c}{a} = \frac{4a}{5c}.$$

291. DEFINITION. When the ratio $a : b$ is compounded with itself the resulting ratio is $a^2 : b^2$, and is called the **duplicate ratio** of $a : b$. Similarly $a^3 : b^3$ is called the **triplicate ratio** of $a : b$. Also $a^{\frac{1}{2}} : b^{\frac{1}{2}}$ is called the **subduplicate ratio** of $a : b$.

- Examples.* (1) The duplicate ratio of $2a : 3b$ is $4a^2 : 9b^2$.
 (2) The subduplicate ratio of $49 : 25$ is $7 : 5$.
 (3) The triplicate ratio of $2x : 1$ is $8x^3 : 1$.

292. DEFINITION. A ratio is said to be a ratio of *greater inequality*, of *less inequality*, or of *equality*, according as the antecedent is *greater than*, *less than*, or *equal to* the consequent.

293. If to each term of the ratio $8 : 3$ we add 4, a new ratio $12 : 7$ is obtained, and we see that it is less than the former because $\frac{12}{7}$ is clearly less than $\frac{8}{3}$.

This is a particular case of a more general proposition which we shall now prove.

A ratio of greater inequality is diminished, and a ratio of less inequality is increased, by adding the same quantity to both its terms.

Let $\frac{a}{b}$ be the ratio, and let $\frac{a+x}{b+x}$ be the new ratio formed by adding x to both its terms.

$$\begin{aligned}\text{Now} \quad \frac{a}{b} - \frac{a+x}{b+x} &= \frac{ax - bx}{b(b+x)} \\ &= \frac{x(a-b)}{b(b+x)};\end{aligned}$$

and $a-b$ is positive or negative according as a is greater or less than b .

$$\text{Hence if } a > b, \quad \frac{a}{b} > \frac{a+x}{b+x};$$

$$\text{and if } a < b, \quad \frac{a}{b} < \frac{a+x}{b+x},$$

which proves the proposition.

Similarly it can be proved that a ratio of *greater inequality* is *increased*, and a ratio of *less inequality* is *diminished*, by taking the same quantity from both its terms.

294. When two or more ratios are equal, many useful propositions may be proved by introducing a single symbol to denote each of the equal ratios.

The proof of the following important theorem will illustrate the method of procedure.

If

$$\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \dots,$$

$$\text{each of these ratios} = \left(\frac{pa^n + qc^n + re^n + \dots}{pb^n + qd^n + rf^n + \dots} \right)^{\frac{1}{n}},$$

where p, q, r, n are any quantities whatever.

Let

$$\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = \dots = k;$$

then

$$a = bk, \quad c = dk, \quad e = fk, \dots;$$

whence

$$pa^n = pb^n k^n, \quad qc^n = qd^n k^n, \quad re^n = rf^n k^n, \dots;$$

$$\therefore \frac{pa^n + qc^n + re^n + \dots}{pb^n + qd^n + rf^n + \dots} = \frac{pb^n k^n + qd^n k^n + rf^n k^n + \dots}{pb^n + qd^n + rf^n + \dots} = k^n;$$

$$\therefore \left(\frac{pa^n + qc^n + re^n + \dots}{pb^n + qd^n + rf^n + \dots} \right)^{\frac{1}{n}} = k = \frac{a}{b} = \frac{c}{d} = \dots$$

By giving different values to p, q, r, n many particular cases of this general proposition may be deduced; or they may be proved independently by using the same method. For instance,

$$\text{if } \frac{a}{b} = \frac{c}{d} = \frac{e}{f}, \text{ each of these ratios} = \frac{a+c+e}{b+d+f};$$

a result which will frequently be found useful.

Example 1. If $\frac{x}{y} = \frac{3}{4}$ find the value of $\frac{5x-3y}{7x+2y}$,

$$\frac{5x-3y}{7x+2y} = \frac{\frac{5x}{y} - 3}{\frac{7x}{y} + 2} = \frac{\frac{15}{4} - 3}{\frac{21}{4} + 2} = \frac{3}{29}.$$

Example 2. Two numbers are in the ratio of 5 : 8. If 9 be added to each they are in the ratio of 8 : 11. Find the numbers.

Let the numbers be denoted by $5x$ and $8x$.

Then

$$\frac{5x+9}{8x+9} = \frac{8}{11};$$

$$\therefore x = 3.$$

Hence the numbers are 15 and 24.

E.A.

S

Example 3. If $A : B$ be in the duplicate ratio of $A+x : B+x$, prove that $x^2 = AB$.

By the given condition, $\left(\frac{A+x}{B+x}\right)^2 = \frac{A}{B}$;

$$\begin{aligned}\therefore B(A+x)^2 &= A(B+x)^2, \\ A^2B + 2ABx + Bx^2 &= AB^2 + 2ABx + Ax^2, \\ x^2(A-B) &= AB(A-B); \\ \therefore x^2 &= AB,\end{aligned}$$

since $A-B$ is, by supposition, not zero.

EXAMPLES XXXII. a.

Find the ratio compounded of

1. The duplicate ratio of 4 : 3, and the ratio 27 : 8.
2. The ratio 32 : 27, and the triplicate ratio of 3 : 4.
3. The subduplicate ratio of 25 : 36, and the ratio 6 : 25.
4. The ratio 169 : 200, and the duplicate ratio of 15 : 26.
5. The triplicate ratio of $x : y$, and the ratio $2y^2 : 3x^2$.
6. The ratio $3a : 4b$, and the subduplicate ratio of $b^4 : a^4$.
7. If $x : y = 5 : 7$, find the value of $x+y : y-x$.
8. If $\frac{x}{y} = 3\frac{1}{3}$, find the value of $\frac{x-3y}{2x-5y}$.
9. If $b : a = 2 : 5$, find the value of $2a-3b : 3b-a$.
10. If $\frac{a}{b} = \frac{3}{4}$, and $\frac{x}{y} = \frac{5}{7}$, find the value of $\frac{3ax-by}{4by-7ax}$.
11. If $7x-4y : 3x+y = 5 : 13$, find the ratio $x : y$.
12. If $\frac{2a^3-3b^3}{a^3+b^3} = \frac{2}{41}$, find the ratio $a : b$.
13. If $2x : 3y$ be in the duplicate ratio of $2x-m : 3y-m$, prove that $m^2 = 6xy$.
14. If $P : Q$ be the subduplicate ratio of $P-x : Q-x$, prove that $x = \frac{PQ}{P+Q}$.
15. If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$, prove that each of these ratios is equal to

$$\sqrt[3]{\frac{2a^2c+3c^3e+4e^2c}{2b^2d+3d^3e+4f^2d}}$$

16. Two numbers are in the ratio of 3 : 4, and if 7 be subtracted from each the remainders are in the ratio of 2 : 3. Find them.
17. What number must be taken from each term of the ratio 27 : 35 that it may become 2 : 3?
18. What number must be added to each term of the ratio 37 : 29 that it may become 8 : 7?
19. If $\frac{p}{b-c} = \frac{q}{c-a} = \frac{r}{a-b}$, shew that $p+q+r=0$.
20. If $\frac{x}{b+c} = \frac{y}{c+a} = \frac{z}{a-b}$, shew that $x-y+z=0$.
21. If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$, shew that the square root of
$$\frac{a^6b - 2c^5e + 3a^4c^3e^2}{b^7 - 2d^6f + 3b^4cd^2e^2}$$
 is equal to $\frac{ace}{bdf}$.
22. Prove that the ratio $la+mc+ne : lb+md+nf$ will be equal to each of the ratios $a : b, c : d, e : f$, if these be all equal; and that it will be intermediate in value between the greatest and least of these ratios if they be not all equal.
23. If $\frac{bx-ay}{cy-az} = \frac{cx-az}{by-ax} = \frac{z+y}{x+z}$, then will each of these fractions be equal to $\frac{x}{y}$, unless $b+c=0$.
24. If $\frac{2x-3y}{3z+y} = \frac{z-y}{z-x} = \frac{x+3z}{2y-3x}$, prove that each of these ratios is equal to $\frac{x}{y}$; hence shew that either $x=y$, or $z=x+y$.

Proportion.

295. DEFINITION. When two ratios are equal, the four quantities composing them are said to be **proportionals**. Thus if $\frac{a}{b} = \frac{c}{d}$ then a, b, c, d are proportionals. This is expressed by saying that a is to b as c is to d , and the proportion is written

$$a : b :: c : d;$$

or

$$a : b = c : d.$$

The terms a and d are called the *extremes*, b and c the *means*.

296. *If four quantities are in proportion, the product of the extremes is equal to the product of the means.*

Let a, b, c, d be the proportionals.

Then by definition $\frac{a}{b} = \frac{c}{d}$;

whence

$$ad = bc.$$

Hence if any three terms of a proportion are given, the fourth may be found. Thus if a, c, d are given, then $b = \frac{ad}{c}$.

Conversely, if there are any four quantities, a, b, c, d , such that $ad = bc$, then a, b, c, d are proportionals; a and d being the extremes, b and c the means; or vice versa.

297. DEFINITION. Quantities are said to be in **continued proportion** when the first is to the second, as the second is to the third, as the third to the fourth; and so on. Thus $a, b, c, d, \dots$ are in continued proportion when

$$\frac{a}{b} = \frac{b}{c} = \frac{c}{d} = \dots$$

If three quantities a, b, c are in continued proportion, then

$$a : b = b : c;$$

$$\therefore ac = b^2.$$

[Art. 296.]

In this case b is said to be a **mean proportional** between a and c ; and c is said to be a **third proportional** to a and b .

298. *If three quantities are proportionals the first is to the third in the duplicate ratio of the first to the second.*

Let the three quantities be a, b, c ; then $\frac{a}{b} = \frac{b}{c}$.

Now

$$\frac{a}{c} = \frac{a}{b} \times \frac{b}{c} = \frac{a}{b} \times \frac{a}{b} = \frac{a^2}{b^2};$$

that is,

$$a : c = a^2 : b^2.$$

299. *If $a : b = c : d$ and $e : f = g : h$ then will*

$$ae : bf = cg : dh.$$

For

$$\frac{a}{b} = \frac{c}{d} \text{ and } \frac{e}{f} = \frac{g}{h}$$

$$\therefore \frac{ae}{bf} = \frac{cg}{dh} \text{ or } ae : bf = cg : dh.$$

Cor. If $a : b = c : d$,
 and $b : x = d : y$,
 then $a : x = c : y$.

This is the theorem known as *ex æquali* in Geometry.

300. If four quantities, a, b, c, d form a proportion, many other proportions may be deduced by the properties of fractions. The results of these operations are very useful, and some of them are often quoted by the annexed names borrowed from Geometry.

(1) If $a : b = c : d$, then $b : a = d : c$. [Invertendo.]

For $\frac{a}{b} = \frac{c}{d}$; therefore $1 \div \frac{a}{b} = 1 \div \frac{c}{d}$;

that is, $\frac{b}{a} = \frac{d}{c}$;

or $b : a = d : c$.

(2) If $a : b = c : d$, then $a : c = b : d$. [Alternando.]

For $ad = bc$; therefore $\frac{ad}{cd} = \frac{bc}{cd}$;

that is, $\frac{a}{c} = \frac{b}{d}$;

or $a : c = b : d$.

(3) If $a : b = c : d$, then $a + b : b = c + d : d$. [Componendo.]

For $\frac{a}{b} = \frac{c}{d}$; therefore $\frac{a}{b} + 1 = \frac{c}{d} + 1$;

that is, $\frac{a+b}{b} = \frac{c+d}{d}$;

or $a + b : b = c + d : d$.

(4) If $a : b = c : d$, then $a - b : b = c - d : d$. [Dividendo.]

For $\frac{a}{b} = \frac{c}{d}$; therefore $\frac{a}{b} - 1 = \frac{c}{d} - 1$;

that is, $\frac{a-b}{b} = \frac{c-d}{d}$;

or $a - b : b = c - d : d$.

(5) If $a : b = c : d$, then $a + b : a - b = c + d : c - d$.

For by (3)
$$\frac{a+b}{b} = \frac{c+d}{d};$$

and by (4)
$$\frac{a-b}{b} = \frac{c-d}{d};$$

$\therefore$ by division,
$$\frac{a+b}{a-b} = \frac{c+d}{c-d};$$

or
$$a+b : a-b = c+d : c-d.$$

Several other proportions may be proved in a similar way.

301. The results of the preceding article are the algebraical equivalents of some of the propositions in the fifth book of Euclid, and the student is advised to make himself familiar with them in their verbal form. For example, *dividendo* may be quoted as follows :

When there are four proportionals, the excess of the first above the second is to the second, as the excess of the third above the fourth is to the fourth.

302. We shall now compare the algebraical definition of proportion with that given in Euclid.

In algebraical symbols the definition of Euclid may be stated as follows :

Four magnitudes, a, b, c, d are in proportion when $pc \supseteq qd$ according as $pa \supseteq qb$, p and q being any positive integers whatever.

I. To deduce the geometrical definition of proportion from the algebraical definition.

Since $\frac{a}{b} = \frac{c}{d}$, by multiplying both sides by $\frac{p}{q}$, we obtain

$$\frac{pa}{qb} = \frac{pc}{qd};$$

hence, from the properties of fractions,

$$pc \supseteq qd \text{ according as } pa \supseteq qb,$$

which proves the proposition.

II. To deduce the algebraical definition of proportion from the geometrical definition.

Given that $pc \supseteq qd$ according as $pa \supseteq qb$, to prove

$$\frac{a}{b} = \frac{c}{d}$$

If $\frac{a}{b}$ is not equal to $\frac{c}{d}$, one of them must be the greater. Suppose $\frac{a}{b} > \frac{c}{d}$; then it will be possible to find some fraction $\frac{q}{p}$ which lies between them.

Hence

$$\frac{a}{b} > \frac{q}{p} \dots\dots\dots(1),$$

and

$$\frac{c}{d} < \frac{q}{p} \dots\dots\dots(2),$$

From (1)

$$pa > qb;$$

from (2)

$$pc < qd;$$

and these contradict the hypothesis.

Therefore $\frac{a}{b}$ and $\frac{c}{d}$ are not unequal; that is $\frac{a}{b} = \frac{c}{d}$; which proves the proposition.

Note. The student should observe that the geometrical definition of proportion deals with *concrete* magnitudes, such as lines or areas, represented geometrically but not referred to any common unit. So that Euclid's definition is applicable to incommensurable as well as to commensurable quantities; whereas the algebraical definition, strictly speaking, applies only to commensurable quantities, since it tacitly assumes that a is the same determinate multiple, part, or parts, of b that c is of d . But the proofs which have been given for commensurable quantities will still be true for incommensurables, since the ratio of two incommensurables can always be made to differ from the ratio of two integers by less than any assignable quantity. [See Art. 289.]

Example 1. If $a : b = c : d = e : f$,

shew that $2a^2 + 3c^2 - 5e^2 : 2b^2 + 3d^2 - 5f^2 = ae : bf$.

Let $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$; then $a = bk$, $c = dk$, $e = fk$;

$$\begin{aligned} \therefore \frac{2a^2 + 3c^2 - 5e^2}{2b^2 + 3d^2 - 5f^2} &= \frac{2b^2k^2 + 3d^2k^2 - 5f^2k^2}{2b^2 + 3d^2 - 5f^2} \\ &= k^2 = \frac{a}{b} \times \frac{e}{f} = \frac{ae}{bf}, \end{aligned}$$

or

$$2a^2 + 3c^2 - 5e^2 : 2b^2 + 3d^2 - 5f^2 = ae : bf.$$

Example 2. If

$(3a + 6b + c + 2d)(3a - 6b - c + 2d) = (3a - 6b + c - 2d)(3a + 6b - c - 2d)$,
 prove that a, b, c, d are in proportion.

We have
$$\frac{3a + 6b + c + 2d}{3a - 6b + c - 2d} = \frac{3a + 6b - c - 2d}{3a - 6b - c + 2d} \quad [\text{Art. 296.}]$$

Componendo and Dividendo,

$$\frac{2(3a + c)}{2(6b + 2d)} = \frac{2(3a - c)}{2(6b - 2d)}.$$

Alternando,

$$\frac{3a + c}{3a - c} = \frac{6b + 2d}{6b - 2d}.$$

Again, Componendo and Dividendo,

$$\frac{6a}{2c} = \frac{12b}{4d};$$

whence

$$a : b = c : d.$$

Example 3. Solve the equation $\frac{x^2 + x - 2}{x - 2} = \frac{4x^2 + 5x - 6}{5x - 6}$.

Dividendo,

$$\frac{x^2}{x - 2} = \frac{4x^2}{5x - 6};$$

whence, dividing by x^2 , which gives a solution $x = 0$, [Art. 201.]

$$\frac{1}{x - 2} = \frac{4}{5x - 6};$$

whence

$$x = -2,$$

and therefore the roots are 0, -2.

EXAMPLES XXXII. b.

Find a fourth proportional to

1. a, ab, c . 2. $a^2, 2ab, 3b^2$. 3. $x^3, xy, 5x^2y$.

Find a third proportional to

4. a^2b, ab . 5. $x^3, 2x^2$. 6. $3x, 6xy$. 7. $1, x$.

Find a mean proportional between

8. a^3, b^2 . 9. $2x^3, 8x$. 10. $12ax^2, 3a^3$. 11. $27a^2b^3, 3b$.

If a, b, c be three proportionals, shew that

12. $a : a + b = a - b : a - c$.

13. $(b^2 + bc + c^2)(ac - bc + c^2) = b^4 + ac^3 + c^4$.

If $a : b = c : d$, prove that

14. $ab + cd : ab - cd = a^2 + c^2 : a^2 - c^2$.
 15. $a^2 + ac + c^2 : a^2 - ac + c^2 = b^2 + bd + d^2 : b^2 - bd + d^2$.
 16. $a : b = \sqrt{3a^2 + 5c^2} : \sqrt{3b^2 + 5d^2}$.
 17. $\frac{a}{p} + \frac{b}{q} : a = \frac{c}{p} + \frac{d}{q} : c$.
 18. $\frac{b}{a} + \frac{a}{b} : \frac{ab}{a^2 + b^2} = \frac{d}{c} + \frac{c}{d} : \frac{cd}{c^2 + d^2}$.

Solve the equations:

19. $3x - 1 : 6x - 7 = 7x - 10 : 9x + 10$.
 20. $x - 12 : y + 3 = 2x - 19 : 5y - 13 = 5 : 14$.
 21. $\frac{x^2 - 2x + 3}{2x - 3} = \frac{x^2 - 3x + 5}{3x - 5}$. 22. $\frac{2x - 1}{x^2 + 2x - 1} = \frac{x + 4}{x^2 + x + 4}$.
 23. If $(a + b - 3c - 3d)(2a - 2b - c + d) = (2a + 2b - c - d)(a - b - 3c + 3d)$

prove that a, b, c, d are proportionals.

24. If a, b, c, d are in continued proportion, prove that
 $a : d = a^3 + b^3 + c^3 : b^3 + c^3 + d^3$.
 25. If b is a mean proportional between a and c , shew that $4a^3 - 9b^2$ is to $4b^3 - 9c^2$ in the duplicate ratio of a to b .
 26. If a, b, c, d are in continued proportion, prove that $b + c$ is a mean proportional between $a + b$ and $c + d$.
 27. If $a + b : b + c = c + d : d + a$,
 prove that $a = c$, or $a + b + c + d = 0$.

Variation.

303. DEFINITION. One quantity A is said to **vary directly** as another B , when the two quantities depend upon each other in such a manner that if B is changed, A is changed in the same ratio.

Note. The word *directly* is often omitted, and A is said to vary as B .

304. For instance: if a train moving at a uniform rate travels 40 miles in 60 minutes, it will travel 20 miles in 30 minutes, 80 miles in 120 minutes, and so on; the distance in each case being increased or diminished in the same ratio as the time. This is expressed by saying that when the velocity is uniform the distance is proportional to the time, or more briefly, the distance varies as the time.

Again, if we refer to the general formula of Art. 84, we find that $\frac{s}{t} = v$ is a relation connecting the space described by a body which moves for a time t with *uniform* velocity v . That is, if $s_1, s_2, s_3 \dots$ be spaces described in times $t_1, t_2, t_3 \dots$ respectively, we have

$$\frac{s_1}{t_1} = \frac{s_2}{t_2} = \frac{s_3}{t_3} = \dots = v.$$

From this it appears that the ratio of any value of s to the corresponding value of t is *constant*, that is, remains the same whatever numerical values s and t may have.

This is an instance of *direct variation*, and s is said to *vary* as t .

305. The symbol $\propto$ is used to denote variation; so that $A \propto B$ is read " A varies as B ."

306. If A varies as B , then A is equal to B multiplied by some constant quantity.

For suppose that $a_1, a_2, a_3 \dots, b_1, b_2, b_3 \dots$ are corresponding values of A and B .

Then, by definition, $\frac{A}{a_1} = \frac{B}{b_1}$; $\frac{A}{a_2} = \frac{B}{b_2}$; $\frac{A}{a_3} = \frac{B}{b_3}$; and so on.

$$\therefore \frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3} = \dots, \text{ each being equal to } \frac{A}{B}.$$

Hence $\frac{\text{any value of } A}{\text{the corresponding value of } B}$ is always the same;

that is, $\frac{A}{B} = m$, where m is constant.

$$\therefore A = mB.$$

307. DEFINITION. One quantity A is said to **vary inversely** as another B when A varies *directly* as the reciprocal of B . [See Art. 107].

Thus if A varies inversely as B , $A = \frac{m}{B}$, where m is constant.

The following is an illustration of inverse variation: If 6 men do a certain work in 8 hours, 12 men would do the same work in 4 hours, 2 men in 24 hours; and so on. Thus it appears that when the number of men is increased the time is proportionately decreased; and vice versa.

308. DEFINITION. One quantity is said to **vary jointly** as a number of others when it varies directly as their product.

Thus A varies jointly as B and C when $A = mBC$, where m is constant. For instance, the interest on a sum of money varies jointly as the principal, the time, and the rate per cent.

309. DEFINITION. A is said to vary directly as B and inversely as C when A varies as $\frac{B}{C}$.

310. If A varies as B when C is constant, and A varies as C when B is constant, then will A vary as BC when both B and C vary.

The variation of A depends partly on that of B and partly on that of C . Suppose these latter variations to take place separately, each in its turn producing its own effect on A ; also let a, b, c be certain simultaneous values of A, B, C .

1. Let C be constant while B changes to b ; then A must undergo a partial change and will assume some intermediate value a' , where

$$\frac{A}{a'} = \frac{B}{b} \dots\dots\dots(1).$$

2. Let B be constant, that is, let it retain its value b , while C changes to c ; then A must complete its change and pass from its intermediate value a' to its final value a , where

$$\frac{a'}{a} = \frac{C}{c} \dots\dots\dots(2).$$

$$\text{From (1) and (2)} \quad \frac{A}{a'} \times \frac{a'}{a} = \frac{B}{b} \times \frac{C}{c};$$

that is,

$$A = \frac{a}{bc} \cdot BC.$$

or

$$A \text{ varies as } BC.$$

311. The following are illustrations of the theorem proved in the last article.

The amount of work done by a given number of men varies directly as the number of days they work, and the amount of work done in a given time varies directly as the number of men; therefore when the number of days and the number of men are both variable, the amount of work will vary as the product of the number of men and the number of days.

Again, in Geometry the area of a triangle varies directly as its base when the height is constant, and directly as the height when the base is constant; and when both the height and base are variable, the area varies as the product of the numbers representing the height and the base.

Example 1. If $A \propto B$, and $C \propto D$, then will $AC \propto BD$.

For, by supposition, $A = mB$, $C = nD$, where m and n are constants. Therefore $AC = mnBD$; and as mn is constant, $AC \propto BD$.

Example 2. If x varies inversely as $y^2 - 1$, and is equal to 24 when $y = 10$; find x when $y = 5$.

By supposition, $x = \frac{m}{y^2 - 1}$, where m is constant.

Putting $x = 24$, $y = 10$, we obtain $24 = \frac{m}{99}$,

whence

$$m = 24 \times 99.$$

$$\therefore x = \frac{24 \times 99}{y^2 - 1};$$

hence, putting $y = 5$, we obtain $x = 99$.

Example 3. The volume of a pyramid varies jointly as its height and the area of its base; and when the area of the base is 60 square feet and the height 14 feet the volume is 280 cubic feet. What is the area of the base of a pyramid whose volume is 390 cubic feet and whose height is 26 feet?

Let V denote the volume, A the area of the base, and h the height;

then $V = mA h$, where m is constant.

Substituting the given values of V , A , h we have

$$280 = m \times 60 \times 14;$$

$$\therefore m = \frac{280}{60 \times 14} = \frac{1}{3}.$$

$$\therefore V = \frac{1}{3} A h.$$

Also when

$$V = 390, \quad h = 26;$$

$$\therefore 390 = \frac{1}{3} A \times 26;$$

$$\therefore A = 45.$$

Hence the area of the base is 45 square feet.

EXAMPLES XXXII. c.

1. If $x \propto y$, and $y=7$ when $x=18$, find x when $y=21$.
2. If $x \propto y$, and $y=3$ when $x=2$, find y when $x=18$.
3. A varies jointly as B and C ; and $A=6$ when $B=3$, $C=2$ find A when $B=5$, $C=7$.
4. A varies jointly as B and C ; and $A=9$ when $B=5$, $C=7$: find B when $A=54$, $C=10$.
5. If $x \propto \frac{1}{y}$, and $y=4$ when $x=15$, find y when $x=6$.
6. If $y \propto \frac{1}{x}$, and $y=1$ when $x=1$, find x when $y=5$.
7. A varies as B directly, and as C inversely; and $A=10$ when $B=15$, $C=6$; find A when $B=8$, $C=2$.
8. If x varies as y directly, and as z inversely, and $x=14$ when $y=10$, $z=14$; find z when $x=49$, $y=45$.
9. If $x \propto \frac{1}{y}$, and $y \propto \frac{1}{z}$, prove that $z \propto x$.
10. If $a \propto b$, prove that $a^n \propto b^n$.
11. If $x \propto z$ and $y \propto z$, prove that $x^2 - y^2 \propto z^2$.
12. If $3a+7b \propto 3a+13b$, and when $a=5$, $b=3$, find the equation between a and b .
13. If $5x-y \propto 10x-11y$, and when $x=7$, $y=5$, find the equation between x and y .
14. If the cube of x varies as the square of y , and if $x=3$ when $y=5$, find the equation between x and y .
15. If the square root of a varies as the cube root of b , and if $a=4$ when $b=8$, find the equation between a and b .
16. If y varies inversely as the square of x , and if $y=8$ when $x=3$, find x when $y=2$.
17. If $x \propto y+a$, where a is constant, and $x=15$ when $y=1$, and $x=35$ when $y=5$; find x when $y=2$.
18. If $a+b \propto a-b$, prove that $a^2+b^2 \propto ab$; and if $a \propto b$, prove that $a^2-b^2 \propto ab$.
19. If y be the sum of three quantities which vary as x , x^2 , x^3 respectively, and when $x=1$, $y=4$, when $x=2$, $y=8$, and when $x=3$, $y=18$, express y in terms of x .

20. Given that the area of a circle varies as the square of its radius, and that the area of a circle is 154 square feet when the radius is 7 feet; find the area of a circle whose radius is 10 feet 6 inches.
21. The area of a circle varies as the square of its diameter; prove that the area of a circle whose diameter is $2\frac{1}{2}$ inches is equal to the sum of the areas of two circles whose diameters are $1\frac{1}{2}$, and 2 inches respectively.
22. The pressure of wind on a plane surface varies jointly as the area of the surface, and the square of the wind's velocity. The pressure on a square foot is 1 lb. when the wind is moving at the rate of 15 miles per hour; find the velocity of the wind when the pressure on a square yard is 16 lbs.
23. The value of a silver coin varies directly as the square of its diameter, while its thickness remains the same; it also varies directly as its thickness while its diameter remains the same. Two silver coins have their diameters in the ratio of 4 : 3. Find the ratio of their thicknesses if the value of the first be four times that of the second.
24. The volume of a circular cylinder varies as the square of the radius of the base when the height is the same, and as the height when the base is the same. The volume is 88 cubic feet when the height is 7 feet, and the radius of the base is 2 feet; what will be the height of a cylinder on a base of radius 9 feet, when the volume is 396 cubic feet?
25. The altitude of a triangle varies directly as its area and inversely as its base. A triangle, 2 square yards in area, standing on a base of $13\frac{1}{2}$ feet, has an altitude of $2\frac{2}{3}$ feet: find the altitude of a triangle whose base is 1 foot 4 inches, and whose area is 2 square feet 96 inches.
26. The expenses of a school are partly constant and partly vary as the number of boys. The expenses were £1000 for 150 boys and £840 for 120 boys; what will the expenses be when there are 330 boys.

the square of its
square feet when
whose radius is

diameter; prove
inches is equal
diameters are $1\frac{1}{2}$,

s jointly as the
wind's velocity.
on the wind is
the velocity of
s 16 lbs.

the square of its
; it also varies
ains the same.
ratio of 4 : 3.
of the first be

the square of the
me, and as the
me is 88 cubic
of the base is
r on a base of
t?

s its area and
yards in area,
de of $2\frac{2}{3}$ feet;
foot 4 inches,

partly vary as
00 for 150 boys
nses be when

CHAPTER XXXIII.

ARITHMETICAL PROGRESSION.

312. DEFINITION. Quantities are said to be in **Arithmetical Progression** when they increase or decrease by a *common difference*.

Thus each of the following series forms an Arithmetical Progression :

3, 7, 11, 15, ...

8, 2, -4, -10, ...

$a, a+d, a+2d, a+3d, \dots$

The common difference is found by subtracting *any* term of the series from that which *follows* it. In the first of the above examples the common difference is 4; in the second it is -6; in the third it is d .

313. If we examine the series

$a, a+d, a+2d, a+3d, \dots$

we notice that *in any term the coefficient of d is always less by one than the number of the term in the series.*

Thus the

3rd term is $a+2d$;

6th term is $a+5d$;

20th term is $a+19d$;

and, generally, the p^{th} term is $a+(p-1)d$.

If n be the number of terms, and if l denote the last, or n^{th} term, we have

$$l = a + (n-1)d.$$

314. To find the sum of a number of terms in Arithmetical Progression.

Let a denote the first term, d the common difference, and n the number of terms. Also let l denote the last term, and s the required sum; then

$$s = a + (a+d) + (a+2d) + \dots + (l-2d) + (l-d) + l;$$

and, by writing the series in the reverse order,

$$s = l + (l-d) + (l-2d) + \dots + (a+2d) + (a+d) + a.$$

Adding together these two series,

$$\begin{aligned} 2s &= (a+l) + (a+l) + (a+l) + \dots \text{to } n \text{ terms} \\ &= n(a+l), \end{aligned}$$

$$\therefore s = \frac{n}{2}(a+l) \dots\dots\dots(1);$$

and $l = a + (n-1)d \dots\dots\dots(2),$

$$\therefore s = \frac{n}{2}\{2a + (n-1)d\} \dots\dots\dots(3).$$

315. In the last article we have three useful formulæ (1), (2), (3); in each of these any one of the letters may denote the unknown quantity when the three others are known. [See Art. 82, Chap. ix.] For instance, in (1) if we substitute given values for s, n, l , we obtain an equation for finding a ; and similarly in the other formulæ. But it is necessary to guard against a too mechanical use of these general formulæ, and it will often be found better to solve simple questions by a mental rather than by an actual reference to the requisite formula.

Example 1. Find the 20th and 35th terms of the series

$$38, 36, 34, \dots\dots.$$

Here the common difference is $36 - 38$, or -2 .

$$\begin{aligned} \therefore \text{the } 20^{\text{th}} \text{ term} &= 38 + 19(-2) \\ &= 0; \end{aligned}$$

$$\begin{aligned} \text{and the } 35^{\text{th}} \text{ term} &= 38 + 34(-2) \\ &= -30. \end{aligned}$$

Example 2. Find the sum of the series $5\frac{1}{2}, 6\frac{3}{4}, 8, \dots\dots$ to 17 terms.

Here the common difference is $1\frac{1}{4}$; hence from (3)

$$\begin{aligned} \text{The sum} &= \frac{17}{2} \left\{ 2 \times 5\frac{1}{2} + 16 \times 1\frac{1}{4} \right\} \\ &= \frac{17}{2} (11 + 20) \\ &= \frac{17 \times 31}{2} \\ &= 263\frac{1}{2}. \end{aligned}$$

Example 3. The first term of a series is 5, the last 45, and the sum 400: find the number of terms, and the common difference.

If n be the number of terms, then from (1)

$$400 = \frac{n}{2} (5 + 45);$$

whence

$$n = 16.$$

If d be the common difference,

$$45 = \text{the } 16^{\text{th}} \text{ term}$$

$$= 5 + 15d;$$

whence

$$d = 2\frac{2}{3}.$$

EXAMPLES XXXIII. a.

1. Find the 27th and 41st terms in the series 5, 11, 17, ...
2. Find the 13th and 109th terms in the series 71, 70, 69, ...
3. Find the 17th and 54th terms in the series 10, $11\frac{1}{2}$, 13, ...
4. Find the 20th and 13th terms in the series -3, -2, -1, ...
5. Find the 90th and 16th terms in the series -4, $2\frac{5}{8}$, 9, ...
6. Find the 37th and 89th terms in the series -2.8, 0, 2.8, ...

Find the last term in the following series :

- | | |
|---|--|
| 7. 5, 7, 9, ... to 20 terms. | 8. 7, 3, -1, ... to 15 terms. |
| 9. $13\frac{1}{2}$, 9, $4\frac{1}{2}$, ... to 13 terms. | 10. .8, 1.2, 1.8, ... to 12 terms. |
| 11. 2.7, 3.4, 4.1, ... to 11 terms. | 12. x , $2x$, $3x$, ... to 25 terms. |
| 13. $a-d$, $a+d$, $a+3d$, ... to 30 terms. | |
| 14. $2a-b$, $4a-3b$, $6a-5b$, ... to 40 terms. | |

Find the last term and sum of the following series :

- | | |
|--|--|
| 15. 14, 64, 114, ... to 20 terms. | 16. 1, 1.2, 1.4, ... to 12 terms. |
| 17. 9, 5, 1, ... to 100 terms. | 18. $\frac{1}{4}$, $-\frac{1}{4}$, $-\frac{3}{4}$, ... to 21 terms. |
| 19. $3\frac{1}{2}$, 1, $-1\frac{1}{2}$, ... to 19 terms. | 20. 64, 96, 128, ... to 16 terms. |

Find the sum of the following series :

- | | |
|--|--|
| 21. 5, 9, 13, ... to 19 terms. | 22. 12, 9, 6, ... to 23 terms. |
| 23. $4, 5\frac{1}{2}, 6\frac{1}{2}$, ... to 37 terms. | 24. $10\frac{1}{2}$, 9, $7\frac{1}{2}$, ... to 94 terms. |
| 25. -3, 1, 5, ... to 17 terms. | 26. $10, 9\frac{2}{3}, 9\frac{1}{3}$, ... to 21 terms. |
| 27. $p, 3p, 5p$, ... to p terms. | 28. $3a, a, -a$, ... to a terms. |
| 29. $a, 0, -a$, ... to a terms. | 30. $-3q, -q, q$, ... to p terms. |

E.A.

T

Find the number of terms and the common difference when

31. The first term is 3, the last term 90, and the sum 1395.
32. The first term is 79, the last term 7, and the sum 1075.
33. The sum is 24, the first term 9, the last term -6 .
34. The sum is 714, the first term 1, the last term $58\frac{1}{2}$.
35. The last term is -16 , the sum -133 , the first term -3 .
36. The first term is -75 , the sum -740 , the last term 1.
37. The first term is a , the last $13a$, and the sum $49a$.
38. The sum is $-320x$, the first term $3x$, the last term $-35x$.

316. If *any two* terms of an Arithmetical Progression be given, the series can be completely determined; for the data furnish *two* simultaneous equations, the solution of which will give the first term, and the common difference.

Example. Find the series whose 7th and 51st terms are -3 and -355 respectively.

If a be the first term, and d the common difference,

$$-3 = \text{the 7}^{\text{th}} \text{ term}$$

$$= a + 6d;$$

and

$$-355 = \text{the 51}^{\text{st}} \text{ term}$$

$$= a + 50d;$$

whence, by subtraction, $-352 = 44d$;

$\therefore d = -8$; and consequently $a = 45$.

Hence the series is 45, 37, 29, ...

317. DEFINITION. When three quantities are in Arithmetical Progression the middle one is said to be the **arithmetic mean** of the other two.

Thus a is the arithmetic mean between $a-d$ and $a+d$.

318. To find the arithmetic mean between two given quantities.

Let a and b be the two quantities; A the arithmetic mean. Then since a, A, b are in A.P. we must have

$$b - A = A - a,$$

each being equal to the common difference;

whence

$$A = \frac{a+b}{2}.$$

ference when

m 1395.

m 1075.

6.

81.

term - 3.

term 1.

9a.

term - 35x.

Progression be

; for the data

of which will

ms are - 3 and

e,

n Arithmetical

the arithmetic

and $a + d$.

iven quantities.

ithmetic mean.

33. Between two given quantities it is always possible to insert any number of terms such that the whole series thus formed shall be in A.P.; and by an extension of the definition in Art. 317, the terms thus inserted are called the *arithmetic means*.

Example. Insert 20 arithmetic means between 4 and 67.

Including the extremes the number of terms will be 22; so that we have to find a series of 22 terms in A.P., of which 4 is the first and 67 the last.

Let d be the common difference;

then

$$67 = \text{the } 22^{\text{nd}} \text{ term}$$

$$= 4 + 21d;$$

whence $d = 3$, and the series is 4, 7, 10, ... 61, 64, 67;

and the required means are 7, 10, 13, ... 58, 61, 64.

320. To insert a given number of arithmetic means between two given quantities.

Let a and b be the given quantities, n the number of means.

Including the extremes the number of terms will be $n + 2$; so that we have to find a series of $n + 2$ terms in A.P., of which a is the first, and b is the last.

Let d be the common difference;

then

$$b = \text{the } (n + 2)^{\text{th}} \text{ term}$$

$$= a + (n + 1)d;$$

whence

$$d = \frac{b - a}{n + 1};$$

and the required means are

$$a + \frac{b - a}{n + 1}, a + \frac{2(b - a)}{n + 1}, \dots, a + \frac{n(b - a)}{n + 1}.$$

Example 1. Find the 30th term of an A.P. of which the first term is 17, and the 100th term - 16.

Let d be the common difference;

then

$$-16 = \text{the } 100^{\text{th}} \text{ term}$$

$$= 17 + 99d;$$

$$\therefore d = -\frac{1}{3}.$$

The 30th term

$$= 17 + 29\left(-\frac{1}{3}\right)$$

$$= 7\frac{1}{3}.$$

Example 2. The sum of three numbers in A.P. is 33, and their product is 792; find them.

Let a be the *middle* number, d the common difference; then the three numbers are $a-d$, a , $a+d$.

Hence $a-d+a+a+d=33$;
whence $a=11$, and the three numbers are $11-d$, 11 , $11+d$.

$$\therefore 11(11+d)(11-d)=792,$$

$$121-d^2=72,$$

$$d=\pm 7;$$

and the numbers are 4, 11, 18.

Example 3. How many terms of the series 24, 20, 16, ... must be taken that the sum may be 72?

Let the number of terms be n ; then, since the common difference is 20-24, or -4, we have from (3), Art. 314,

$$72=\frac{n}{2}\{2 \times 24+(n-1)(-4)\}$$

$$=24n-2n(n-1);$$

$$\text{whence } n^2-13n+36=0,$$

$$\text{or } (n-4)(n-9)=0;$$

$$\therefore n=4 \text{ or } 9.$$

Both these values satisfy the conditions of the question; for if we write down the first 9 terms, we get 24, 20, 16, 12, 8, 4, 0, -4, -8; and, as the last five terms destroy each other, the sum of 9 terms is the same as that of 4 terms.

Example 4. An A.P. consists of 21 terms; the sum of the three terms in the middle is 129, and of the last three is 237; find the series.

Let a be the first term, and d the common difference. Then

$$237=\text{the sum of the last three terms}$$

$$=a+20d+a+19d+a+18d$$

$$=3a+57d;$$

$$\text{whence } a+19d=79 \dots\dots\dots(1).$$

Again, the three middle terms are the 10th, 11th, 12th;

$$\text{hence } 129=\text{the sum of the three middle terms}$$

$$=a+9d+a+10d+a+11d$$

$$=3a+30d;$$

$$\text{whence } a+10d=43 \dots\dots\dots(2).$$

From (1) and (2), we obtain $d=4$, $a=3$.

Hence the series is 3, 7, 11, ... 83.

EXAMPLES XXXIII. b.

Find the series in which

1. The 27th term is 186, and the 45th term 312.
2. The 5th term is 1, and the 31st term - 77.
3. The 15th term is - 25, and the 23rd term - 41.
4. The 9th term is - 11, and the 102nd term - 150 $\frac{1}{2}$.
5. The 15th term is 25, and the 29th term 46.
6. The 16th term is 214, and the 51st term 739.
7. The 3rd and 7th terms of an A.P. are 7 and 19; find the 15th term.
8. The 54th and 4th terms are - 125 and 0; find the 42nd term.
9. The 31st and 2nd terms are $\frac{1}{2}$ and $7\frac{3}{4}$; find the 59th term.
10. Insert 15 arithmetic means between 71 and 23.
11. Insert 17 arithmetic means between 93 and 69.
12. Insert 14 arithmetic means between $-7\frac{1}{8}$ and $-2\frac{1}{8}$.
13. Insert 16 arithmetic means between 7.2 and - 6.4.
14. Insert 36 arithmetic means between $8\frac{1}{3}$ and $2\frac{1}{3}$.

How many terms must be taken of

15. The series 42, 39, 36, ... to make 315?
16. The series - 16, - 15, - 14, ... to make - 100?
17. The series $15\frac{2}{3}$, $15\frac{1}{3}$, 15, ... to make 129?
18. The series 20, $18\frac{3}{4}$, $17\frac{1}{2}$, ... to make $162\frac{1}{2}$?
19. The series $-10\frac{1}{2}$, - 9, $-7\frac{1}{2}$, ... to make - 42?
20. The series $-6\frac{4}{5}$, $-6\frac{3}{5}$, - 6, ... to make $-52\frac{4}{5}$.
21. The sum of three numbers in A.P. is 39, and their product is 2184; find them.
22. The sum of three numbers in A.P. is 12, and the sum of their squares is 66; find them.
23. The sum of five numbers in A.P. is 75, and the product of the greatest and least is 161; find them.
24. The sum of five numbers in A.P. is 40, and the sum of their squares is 410; find them.
25. The 12th, 85th and last terms of an A.P. are 38, 257, 395 respectively; find the number of terms.

[Examples for revision will be found in Miscellaneous Examples V., page 307.]

CHAPTER XXXIV.

GEOMETRICAL PROGRESSION.

321. DEFINITION. Quantities are said to be in **Geometrical Progression** when they increase or decrease by a *constant factor*.

Thus each of the following series forms a Geometrical Progression :

$$3, 6, 12, 24, \dots\dots\dots$$

$$1, -\frac{1}{3}, \frac{1}{9}, -\frac{1}{27}, \dots\dots\dots$$

$$a, ar, ar^2, ar^3, \dots\dots\dots$$

The constant factor is also called the *common ratio*, and it is found by dividing *any* term by that which immediately *precedes* it. In the first of the above examples the common ratio is 2; in the second it is $-\frac{1}{3}$; in the third it is r .

322. If we examine the series

$$a, ar, ar^2, ar^3, ar^4, \dots\dots$$

we notice that *in any term the index of r is always less by one than the number of the term in the series.*

Thus the 3rd term is ar^2 ;
 the 6th term is ar^5 ;
 the 20th term is ar^{19} ;

and, generally, the p^{th} term is ar^{p-1} .

If n be the number of terms, and if l denote the last, or n^{th} term, we have

$$l = ar^{n-1}.$$

Example. Find the 8th term of the series $-\frac{1}{3}, \frac{1}{2}, -\frac{3}{4}, \dots\dots$

The common ratio is $\frac{1}{2} \div \left(-\frac{1}{3}\right)$, or $-\frac{3}{2}$;

$$\begin{aligned} \therefore \text{the 8}^{\text{th}} \text{ term} &= -\frac{1}{3} \times \left(-\frac{3}{2}\right)^7 \\ &= -\frac{1}{3} \times -\frac{2187}{128} \\ &= \frac{729}{128} \end{aligned}$$

323. DEFINITION. When three quantities are in Geometrical Progression the middle one is called the **geometric mean** between the other two.

To find the geometric mean between two given quantities.

Let a and b be the two quantities; G the geometric mean. Then since a, G, b are in G.P.,

$$\frac{b}{G} = \frac{G}{a}$$

each being equal to the common ratio ;

$$\therefore G^2 = ab ;$$

whence

$$G = \sqrt{ab}.$$

324. To insert a given number of geometric means between two given quantities.

Let a and b be the given quantities, n the number of means.

In all there will be $n+2$ terms ; so that we have to find a series of $n+2$ terms in G.P., of which a is the first and b the last.

Let r be the common ratio ;

then

$$\begin{aligned} b &= \text{the } (n+2)^{\text{th}} \text{ term} \\ &= ar^{n+1} ; \end{aligned}$$

$$\therefore r^{n+1} = \frac{b}{a} ;$$

$$\therefore r = \left(\frac{b}{a} \right)^{\frac{1}{n+1}} \dots \dots \dots (1).$$

Hence the required means are $ar, ar^2, \dots, ar^n$, where r has the value found in (1).

Example. Insert 4 geometric means between 160 and 5

We have to find 6 terms in G.P. of which 160 is the first, and 5 the sixth.

Let r be the common ratio ;

$$\begin{aligned} \text{then } 5 &= \text{the sixth term} \\ &= 160r^5 ; \end{aligned}$$

$$\therefore r^5 = \frac{1}{32} ;$$

whence, *by trial*,

$$r = \frac{1}{2} ;$$

and the means are 80, 40, 20, 10.

325. To find the sum of a number of terms in Geometrical Progression.

Let a be the first term, r the common ratio, n the number of terms, and s the sum required. Then

$$s = a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1};$$

multiplying every term by r , we have

$$rs = ar + ar^2 + \dots + ar^{n-1} + ar^n.$$

Hence by subtraction,

$$rs - s = ar^n - a;$$

$$\therefore (r-1)s = a(r^n - 1);$$

$$\therefore s = \frac{a(r^n - 1)}{r - 1} \dots\dots\dots(1).$$

Changing the signs in numerator and denominator [Art. 170.]

$$s = \frac{a(1 - r^n)}{1 - r} \dots\dots\dots(2).$$

Note. It will be found convenient to remember both forms given above for s , using (2) in all cases except when r is positive and greater than 1.

Since $ar^{n-1} = l$, the formula (1) may be written

$$s = \frac{rl - a}{r - 1};$$

a form which is sometimes useful.

Example 1. Sum the series 81, 54, 36, ... to 9 terms.

The common ratio $= \frac{54}{81} = \frac{2}{3}$, which is less than 1;

hence the sum

$$\begin{aligned} &= \frac{81 \left\{ 1 - \left(\frac{2}{3} \right)^9 \right\}}{1 - \frac{2}{3}} \\ &= 243 \left\{ 1 - \left(\frac{2}{3} \right)^9 \right\} \\ &= 243 - \frac{512}{81} \\ &= 236 \frac{55}{81} \end{aligned}$$

Example 2. Sum the series $\frac{2}{3}, -1, \frac{3}{2}, \dots$ to 7 terms.

The common ratio $= -\frac{3}{2}$; hence by formula (2)

$$\begin{aligned} \text{the sum} &= \frac{\frac{2}{3} \left\{ 1 - \left(-\frac{3}{2} \right)^7 \right\}}{1 + \frac{3}{2}} \\ &= \frac{\frac{2}{3} \left\{ 1 + \frac{2187}{128} \right\}}{\frac{5}{2}} \\ &= \frac{2}{3} \times \frac{2315}{128} \times \frac{2}{5} \\ &= \frac{463}{96}. \end{aligned}$$

EXAMPLES XXXIV. a.

1. Find the 5th and 8th terms of the series 3, 6, 12, ...
2. Find the 10th and 16th terms of the series 256, 128, 64, ...
3. Find the 7th and 11th terms of the series 64, -32, 16, ...
4. Find the 8th and 12th terms of the series 81, -27, 9, ...
5. Find the 14th and 7th terms of the series $\frac{1}{64}, \frac{1}{32}, \frac{1}{16}, \dots$
6. Find the 4th and 8th terms of the series .008, .04, .2, ...

Find the last term in the following series :

7. 2, 4, 8, ... to 9 terms.
8. 2, -6, 18, ... to 8 terms.
9. 2, 3, $4\frac{1}{2}$, ... to 6 terms.
10. 3, -3², 3³, ... to 2n terms.
11. $x, x^3, x^9, \dots$ to p terms.
12. $x, 1, \frac{1}{x}, \dots$ to 30 terms.
13. Insert 3 geometric means between 486 and 6.
14. Insert 4 geometric means between $\frac{1}{8}$ and 128.
15. Insert 6 geometric means between 56 and $-\frac{7}{16}$.
16. Insert 5 geometric means between $\frac{32}{81}$ and $4\frac{1}{2}$.

Find the last term and the sum of the following series :

17. 3, 6, 12, ... to 8 terms. 18. 6, -18, 54, ... to 6 terms.
 19. 64, 32, 16, ... to 10 terms. 20. 8·1, 2·7, ·9, ... to 7 terms.
 21. $\frac{1}{72}, \frac{1}{24}, \frac{1}{8}, \dots$ to 8 terms. 22. $4\frac{1}{2}, 1\frac{1}{2}, \frac{1}{2}, \dots$ to 9 terms.

Find the sum of the series

23. 3, -1, $\frac{1}{3}, \dots$ to 6 terms. 24. $\frac{1}{2}, \frac{1}{3}, \frac{2}{9}, \dots$ to 7 terms.
 25. $-\frac{2}{5}, \frac{1}{2}, -\frac{5}{8}, \dots$ to 6 terms. 26. $1, -\frac{1}{2}, \frac{1}{4}, \dots$ to 12 terms.
 27. 9, -6, 4, ... to 7 terms. 28. $\frac{2}{3}, -\frac{1}{6}, \frac{1}{24}, \dots$ to 8 terms.
 29. 1, 3, $3^2, \dots$ to p terms. 30. 2, -4, 8, ... to $2p$ terms.
 31. $\frac{1}{\sqrt{3}}, 1, \frac{3}{\sqrt{3}}, \dots$ to 8 terms. 32. $\sqrt{a}, \sqrt{a^3}, \sqrt{a^5}, \dots$ to a terms.
 33. $\frac{1}{\sqrt{2}}, -2, \frac{8}{\sqrt{2}}, \dots$ to 7 terms. 34. $\sqrt{2}, \sqrt{6}, 3\sqrt{2}, \dots$ to 12 terms.

326. Consider the series $1, \frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \dots$

$$\begin{aligned} \text{The sum to } n \text{ terms} &= \frac{1 - \frac{1}{2^n}}{1 - \frac{1}{2}} \\ &= 2\left(1 - \frac{1}{2^n}\right) \\ &= 2 - \frac{1}{2^{n-1}} \end{aligned}$$

From this result it appears that however many terms be taken the sum of the above series is always less than 2. Also we see that, by making n sufficiently large, we can make the fraction $\frac{1}{2^{n-1}}$ as small as we please. Thus by taking a sufficient number of terms the sum can be made to differ by as little as we please from 2.

In the next article a more general case is discussed.

g series :

to 6 terms,

to 7 terms,

to 9 terms,

to 7 terms.

to 12 terms.

to 8 terms.

to 2p terms.

... to a terms.

... to 12 terms.

327. From Art. 325 we have $s = \frac{a(1-r^n)}{1-r}$

$$= \frac{a}{1-r} - \frac{ar^n}{1-r}$$

Suppose r is a proper fraction ; then the greater the value of n the smaller is the value of r^n , and consequently of $\frac{ar^n}{1-r}$; and therefore by making n sufficiently large, we can make the sum of n terms of the series differ from $\frac{a}{1-r}$ by as small a quantity as we please.

This result is usually stated thus : *the sum of an infinite number of terms of a decreasing Geometrical Progression is $\frac{a}{1-r}$;*

or more briefly, *the sum to infinity is $\frac{a}{1-r}$;*

328. Recurring decimals furnish a good illustration of infinite Geometrical Progressions.

Example. Find the value of $\cdot 4\dot{2}\dot{3}$.

$$\cdot 4\dot{2}\dot{3} = \cdot 4232323 \dots\dots$$

$$= \frac{4}{10} + \frac{23}{1000} + \frac{23}{100000} + \dots\dots$$

$$= \frac{4}{10} + \frac{23}{10^3} + \frac{23}{10^5} + \dots\dots$$

$$= \frac{4}{10} + \frac{23}{10^3} \left(1 + \frac{1}{10^2} + \frac{1}{10^4} + \dots\dots \right)$$

$$= \frac{4}{10} + \frac{23}{10^3} \cdot \frac{1}{1 - \frac{1}{10^2}}$$

$$= \frac{4}{10} + \frac{23}{10^3} \cdot \frac{100}{99}$$

$$= \frac{4}{10} + \frac{23}{990}$$

$$= \frac{419}{990}$$

which agrees with the value found by the usual arithmetical rule.

EXAMPLES XXXIV. b.

Sum to infinity the following series :

- | | |
|--|--|
| 1. 9, 6, 4, ... | 2. 12, 6, 3, ... |
| 3. $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots$ | 4. $\frac{1}{2}, -\frac{1}{4}, \frac{1}{8}, \dots$ |
| 5. $\frac{1}{3}, \frac{2}{9}, \frac{4}{27}, \dots$ | 6. $\frac{8}{5}, -1, \frac{5}{8}, \dots$ |
| 7. .9, .03, .001, ... | 8. .8, -.4, .2, ... |

Find by the method of Art. 328, the value of

9. .3 10. .16. 11. .24. 12. .375. 13. .037.

Find the series in which

- The 10th term is 320 and the 6th term 20.
- The 5th term is $\frac{27}{16}$ and the 9th term is $\frac{1}{3}$.
- The 7th term is 625 and the 4th term -5.
- The 3rd term is $\frac{9}{16}$ and the 6th term $-4\frac{1}{2}$.
- Divide 183 into three parts in G.P. such that the sum of the first and third is $2\frac{1}{20}$ times the second.
- Show that the product of any odd number of consecutive terms of a G.P. will be equal to the n^{th} power of the middle term, n being the number of terms.
- The first two terms of an infinite G.P. are together equal to 1, and every term is twice the sum of all the terms which follow. Find the series.

Sum the following series :

- $y^2 + 2b, y^4 + 4b, y^6 + 6b, \dots$ to n terms.
- $\frac{3+2\sqrt{2}}{3-2\sqrt{2}}, 1, \frac{3-2\sqrt{2}}{3+2\sqrt{2}}, \dots$ to infinity.
- $\sqrt{\frac{3}{2}}, \frac{1}{3}\sqrt{2}, \frac{2}{9}\sqrt{\frac{2}{3}}, \dots$ to infinity.
- $2n - \frac{1}{2}, 4n + \frac{1}{6}, 6n - \frac{1}{18}, \dots$ to $2n$ terms.

[Examples for revision will be found in Miscellaneous Examples V., p. 307.]

CHAPTER XXXV.

HARMONICAL PROGRESSION.

329. DEFINITION. Three quantities a, b, c are said to be in **Harmonical Progression** when $\frac{a}{c} = \frac{a-b}{b-c}$.

Any number of quantities are said to be in **Harmonical Progression** when every three consecutive terms are in **Harmonical Progression**.

330. *The reciprocals of quantities in Harmonical Progression are in Arithmetical Progression.*

By definition, if a, b, c are in Harmonical Progression,

$$\frac{a}{c} = \frac{a-b}{b-c};$$

$$\therefore a(b-c) = c(a-b),$$

dividing every term by abc ,

$$\frac{1}{c} - \frac{1}{b} = \frac{1}{b} - \frac{1}{a},$$

which proves the proposition.

331. Harmonical properties are chiefly interesting because of their importance in Geometry and in the Theory of Sound: in Algebra the proposition just proved is the only one of any importance. There is no general formula for the sum of any number of quantities in Harmonical Progression. Questions in H.P. are generally solved by inverting the terms, and making use of the properties of the corresponding A.P.

Example. The 12th term of a H.P. is $\frac{1}{5}$, and the 19th term is $\frac{3}{22}$; find the series.

Let a be the first term, d the common difference of the corresponding A.P.; then

$$\begin{aligned} 5 &= \text{the 12th term} \\ &= a + 11d; \end{aligned}$$

and

$$\begin{aligned} \frac{22}{3} &= \text{the 19th term} \\ &= a + 18d; \end{aligned}$$

whence

$$d = \frac{1}{3}, \quad a = \frac{4}{3}.$$

Hence the Arithmetical Progression is $\frac{4}{3}, \frac{5}{3}, 2, \frac{7}{3}, \dots$;

and the Harmonical Progression is $\frac{3}{4}, \frac{3}{5}, \frac{1}{2}, \frac{3}{7}, \dots$.

332. To find the harmonic mean between two given quantities.

Let a, b be the two quantities, H their harmonic mean; then $\frac{1}{a}, \frac{1}{H}, \frac{1}{b}$ are in A.P.

$$\begin{aligned} \therefore \frac{1}{H} - \frac{1}{a} &= \frac{1}{b} - \frac{1}{H}, \\ \frac{2}{H} &= \frac{1}{a} + \frac{1}{b}, \\ H &= \frac{2ab}{a+b}. \end{aligned}$$

333. If A, G, H be the arithmetic, geometric, and harmonic means between a and b , we have proved

$$A = \frac{a+b}{2} \dots \dots \dots (1).$$

$$G = \sqrt{ab} \dots \dots \dots (2).$$

$$H = \frac{2ab}{a+b} \dots \dots \dots (3).$$

Therefore

$$\begin{aligned} AH &= \frac{a+b}{2} \cdot \frac{2ab}{a+b} \\ &= ab \\ &= G^2; \end{aligned}$$

that is, G is the geometric mean between A and H .

19th term is $\frac{3}{22}$:

the correspond.

..... ;

ven quantities.

monic mean ;

and harmonic

.....(1).

.....(2).

.....(3).

334. Miscellaneous questions in the Progressions afford scope for much skill and ingenuity, the solution being often very neatly effected by some special artifice. The student will find the following hints useful :

1. If the same quantity be added to, or subtracted from, all the terms of an A.P., the resulting terms will form an A.P. with the same common difference as before. [Art. 312.]

2. If all the terms of an A.P. be multiplied or divided by the same quantity, the resulting terms form an A.P., but with a new common difference. [Art. 312.]

3. If all the terms of a G.P. be multiplied or divided by the same quantity, the resulting terms form a G.P. with the same common ratio as before. [Art. 322.]

4. If $a, b, c, d, \dots$ be in G.P., they are also in *continued proportion*, since, by definition,

$$\frac{a}{b} = \frac{b}{c} = \frac{c}{d} = \dots = \frac{1}{r}$$

Conversely, a series of quantities in continued proportion may be represented by $x, ar, ar^2, \dots$.

Example 1. Find three quantities in G.P. such that their product is 343, and their sum 30 $\frac{1}{2}$.

Let $\frac{a}{r}, a, ar$ be the three quantities ;

then we have $\frac{a}{r} \times a \times ar = 343 \dots\dots\dots(1),$

and $a\left(\frac{1}{r} + 1 + r\right) = \frac{91}{3} \dots\dots\dots(2).$

From (1) $a^3 = 343,$

$$a = 7 ;$$

$\therefore$ from (2) $7(1 + r + r^2) = \frac{91}{3}r.$

Whence we obtain $r = 3, \text{ or } \frac{1}{3},$

and the numbers are $\frac{7}{3}, 7, 21.$

Example 2. If a, b, c be in H.P., prove that $\frac{a}{b+c}, \frac{b}{c+a}, \frac{c}{a+b}$ are also in H.P.

Since $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P.,

$$\frac{a+b+c}{a}, \frac{a+b+c}{b}, \frac{a+b+c}{c} \text{ are in A.P. ;}$$

$$\therefore 1 + \frac{b+c}{a}, 1 + \frac{a+c}{b}, 1 + \frac{a+b}{c} \text{ are in A.P. ;}$$

$$\therefore \frac{b+c}{a}, \frac{a+c}{b}, \frac{a+b}{c} \text{ are in A.P. ;}$$

$$\therefore \frac{a}{b+c}, \frac{b}{c+a}, \frac{c}{a+b} \text{ are in H.P.}$$

Example 3. The n^{th} term of an A.P. is $\frac{n}{5} + 2$, find the sum of 49 terms.

Let a be the first term, and l the last ; then by putting $n=1$, and $n=49$ respectively, we obtain

$$a = \frac{1}{5} + 2, \quad l = \frac{49}{5} + 2 ;$$

$$\begin{aligned} \therefore s &= \frac{n}{2}(a+l) = \frac{49}{2} \left(\frac{50}{5} + 4 \right) \\ &= \frac{49}{2} \times 14 = 343. \end{aligned}$$

Example 4. If a, b, c, d, e be in G.P., prove that $b+d$ is the geometric mean between $a+c$ and $c+e$.

Since a, b, c, d, e are in continued proportion,

$$\frac{a}{b} = \frac{b}{c} = \frac{c}{d} = \frac{d}{e} ;$$

$\therefore$ each ratio

$$= \frac{a+c}{b+d} = \frac{b+d}{c+e}. \quad [\text{Art. 294.}]$$

Whence

$$(b+d)^2 = (a+c)(c+e).$$

EXAMPLES XXXV.

1. Find the 6th term of the series $4, 2, 1\frac{1}{3}, \dots$
2. Find the 21st term of the series $2\frac{1}{2}, 1\frac{1}{3}, 1\frac{2}{5}, \dots$
3. Find the 8th term of the series $1\frac{1}{3}, 1\frac{1}{17}, 2\frac{2}{13}, \dots$
4. Find the n^{th} term of the series $3, 1\frac{1}{2}, 1, \dots$

$$\frac{a}{c}, \frac{b}{c+a}, \frac{c}{a+b}$$

P.;

P.;

P.;

P.

and the sum of

tting $n=1$, andat $b+d$ is the

[Art. 204.]

Find the series in which

5. The 15th term is $\frac{1}{25}$, and the 23rd term is $\frac{1}{41}$.
 6. The 2nd term is 2, and the 31st term is $\frac{4}{31}$.
 7. The 39th term is $\frac{1}{11}$, and the 54th term is $\frac{1}{26}$.

Find the harmonic mean between

8. 2 and 4. 9. 1 and 13. 10. $\frac{1}{4}$ and $\frac{1}{10}$.
 11. $\frac{1}{a}$ and $\frac{1}{b}$. 12. $\frac{1}{x+y}$ and $\frac{1}{x-y}$. 13. $x+y$ and $x-y$.
 14. Insert two harmonic means between 4 and 12.
 15. Insert three harmonic means between $2\frac{2}{5}$ and 12.
 16. Insert four harmonic means between 1 and 6.
 17. If G be the geometric mean between two quantities A and B , shew that the ratio of the arithmetic and harmonic means of A and G is equal to the ratio of the arithmetic and harmonic means of G and B .
 18. To each of three consecutive terms of a G.P. the second of the three is added. Shew that the three resulting quantities are in H.P.

Sum the following series :

19. $1 + 1\frac{3}{4} + 3\frac{1}{16} + \dots$ to 6 terms.
 20. $1 + 1\frac{3}{4} + 2\frac{1}{2} + \dots$ to 6 terms.
 21. $(2a+x) + 3a + (4a-x) + \dots$ to p terms.
 22. $1\frac{4}{5} - 1\frac{1}{5} + \frac{4}{5} - \dots$ to 8 terms.
 23. $1\frac{4}{5} + 1\frac{1}{5} + \frac{3}{5} + \dots$ to 12 terms.
 24. If $x-a$, $y-a$, and $z-a$ be in G.P., prove that $2(y-a)$ is the harmonic mean between $y-x$ and $y-z$.
 25. If a, b, c, d be in A.P., a, e, f, d in G.P., a, g, h, d in H.P. respectively; prove that $ad=ef=bh=cg$.
 26. If a^2, b^2, c^2 be in A.P., prove that $b+c, c+a, a+b$ are in H.P.

E.A.

U

27. If a, b, c be in A.P., and α, β, γ in H.P., shew that

$$\frac{\alpha+c}{b\beta} = \frac{\alpha+\gamma}{a\gamma}.$$

28. If a be the arithmetic mean between b and c , and b the geometric mean between a and c , prove that c will be the harmonic mean between a and b .

29. If $\frac{a+b}{2}, b, \frac{b+c}{2}$ be in H.P., then a, b, c are in G.P.

30. If a, b, c, d, e be in G.P., prove that $c(a+2c+e)=(b+d)^2$.

31. If $a, b, c, d \dots$ be a series of quantities in G.P., shew that the reciprocals of $a^2-b^2, b^2-c^2, c^2-d^2, \dots$ are also in G.P.; and find the sum of n terms of this latter series in terms of a and b .

32. If a, b, c be in A.P., and b, c, d in H.P., then $a, \frac{c^2}{d}, c$ are in H.P., and $b, \frac{ad}{b}, d$ are also in H.P.

33. If g be the geometric and a the arithmetic mean between m and n , and if k^2 be the arithmetic mean between m^2 and n^2 , prove that a^2 is the arithmetic mean between g^2 and k^2 .

34. If a, b, c, d be in G.P., prove that $(b-c)^2 = ac + bd - 2ad$.

35. If a, b, c, d be in G.P., prove that

$$(\alpha+d)(\alpha-b)^2 : \alpha(\alpha-c)(\alpha-d) = a-b+c : a+b+c.$$

36. If a, b, c be in H.P., prove that

$$\frac{1}{a} + \frac{1}{b+c}, \frac{1}{b} + \frac{1}{c+a}, \frac{1}{c} + \frac{1}{a+b}$$

are also in H.P.

MISCELLANEOUS EXAMPLES V.

(Chiefly on Chapters XXXII.—XXXV.)

1. Simplify $\frac{(a^2b^3)^{\frac{1}{2}}b^{-2}c^{\frac{1}{2}}}{a^{\frac{1}{2}}b^{-\frac{1}{2}}c^{\frac{1}{2}}}$, and find its value when $a=2$, $b=3$, and $c=432$.

2. Shew that the ratio $x+y : x-y$ is increased by subtracting y from each term.

3. If $\frac{a}{b} = \frac{c}{d}$, shew that

$$(1) \frac{2a+3b}{3a-7b} = \frac{2c+3d}{3c-7d}; \quad (2) \frac{a^2-c^2}{b^2-d^2} = \frac{(a+2c)(a+3c)}{(b+2d)(b+3d)}.$$

4. If $\frac{x}{b-c} = \frac{y}{c-a} = \frac{z}{a-b}$, prove that

$$(1) x+y+z=0; \quad (2) (b+c)x + (c+a)y + (a+b)z = 0.$$

5. If $\frac{x}{1} = \frac{y}{2} = \frac{z}{3}$, prove that $\sqrt{5x^2+8y^2+7z^2}=5y$.

6. If y is the sum of two numbers, of which the first varies directly and the second inversely as x , and if $y=7$, when $x=2$, and $y=-1$, when $x=1$, shew that

$$y = 5x - \frac{6}{x}.$$

7. Simplify $\sqrt{45} + \sqrt{8} - \sqrt{80} + \sqrt{18} + \sqrt{7} - \sqrt{40}$.

8. If $3x+10$ has to $9x+4$ the duplicate ratio of 5 to 7, find x .

9. If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$, prove that each ratio is equal to

$$(1) \sqrt[3]{\frac{4ac^2-3ce^3+2ace}{4bd^2-3cf^3+2bdf}}; \quad (2) \sqrt[5]{\frac{6a^2c^2e-c^4ef+7ac^5}{6b^2d^2f-d^4f^3+7ad^5}}.$$

10. The sides of a triangle are as $1 : 1\frac{1}{2} : 1\frac{3}{4}$, and the perimeter is 221 yards; find the sides.

11. If $3a + 5b : 3a - 5b = 3c + 5d : 3c - 5d$,
prove that $a : b = c : d$.

12. Reduce to their simplest forms:

$$(1) \frac{x^{a+b}}{x^{a-b}} + \frac{x^{a-b}}{x^{a+b}}; \quad (2) \frac{(a+b)^{\frac{3}{2}}}{(a-b)^{\frac{3}{2}}} \times \sqrt{a^2 - b^2}.$$

13. When $x = -\frac{3a}{4}$, find the value of

$$\frac{x^2 + ax + a^2}{x^3 - a^3} - \frac{x^2 - ax + a^2}{x^3 + a^3}$$

14. Simplify

$$(1) \left\{ \frac{a^{-1}b^3}{\sqrt[3]{a}} \right\}^{\frac{4}{3}} \div \sqrt[6]{\frac{a^2\sqrt{b}}{b^2}}; \quad (2) \frac{2^{n+4} - 2 \times 2^n}{2^{n+2} \times 4}.$$

15. Find the ratio compounded of the ratios

$$\frac{x-y}{a+b} : \frac{x^3-y^3}{a^3-b^3} \text{ and } \frac{x^2+xy+y^2}{a^2-b^2} : \frac{x^2-y^2}{(a+b)^2}.$$

16. If a, b, c be three proportionals, prove that

$$(1) a(a+b) : b(b-a) = b(b+c) : c(c-b);$$

$$(2) (a+b+c)(b^2-bc+c^2) = c(a^2+b^2+c^2).$$

17. If $a : b : c = xy : x^2 : yz$, prove that $x : y : z = ab : a^2 : bc$.

18. If $p : q$ be the duplicate ratio of $p-r : q-r$, prove that r is a mean proportional between p and q .

19. If $a : b = c : d$, prove that

$$(1) a+c : a+b+c+d = a : a+b;$$

$$(2) (a-b) - (c-d) = \frac{(a-b)(b-d)}{b}.$$

20. Shew that any ratio is made more nearly equal to unity by adding the same quantity to each of its terms.

21. If x varies as $y+z$, and z varies as x ; and if $x=2$ when $y=4$, find the value of y when $x=1$.

22. If $2x+3y : 2x-3y = 2a^2+3b^2 : 2a^2-3b^2$, then x has to y the duplicate ratio of a to b .

23. Find an A.P. of seven terms whose sum is 28 and common difference 3.

24. The sum of 10 terms of an A.P. is 145, and the sum of its fourth and ninth terms is five times the third term; determine the series.

25. Find the value of

$$\sqrt{19+4\sqrt{21}} + \sqrt{7} - \sqrt{12} - \sqrt{29-2\sqrt{28}}.$$

26. Sum to 10 terms each of the series

$$(1) \quad 5+10+15+20+\dots;$$

$$(2) \quad 5-10+20-40+\dots.$$

27. If $\frac{p}{bz-cy} = \frac{-q}{cx+az} = \frac{-r}{ay+bx}$, shew that

$$ap+bq-cr=0, \text{ and } xp-yq+rz=0.$$

28. The sum of five numbers in arithmetical progression is 10, and the sum of their squares is 60; find the numbers.

29. Find the sum of n terms of the progression

$$3+2\frac{1}{2}+2\frac{1}{2}+\dots.$$

30. Find the ninth term of the harmonic series whose first and third terms are 3 and 2 respectively.

31. Simplify $(a-b)^{\frac{1}{3}} \cdot \frac{\sqrt[3]{a^2+2ab+b^2}}{\sqrt[3]{a^2-b^2} \times (a+b)^{-\frac{2}{3}}}$.

32. Sum to n terms

$$\frac{3}{2} + \frac{9}{2} + \frac{15}{2} + \frac{21}{2} + \dots,$$

and find five consecutive terms of this progression whose sum is $187\frac{1}{2}$.

33. The 8th term of an Arithmetical Progression is double the 13th term; shew that the 2nd term is double the 10th term.

34. Sum the following series :

(1) $(a-2x)+2(a+x)+3(a+2x)+\dots$ to 18 terms.

(2) $3\frac{7}{8}-5\frac{2}{7}+8\frac{8}{9}-\dots$ to 7 terms.

35. Shew that the sum of $2n$ terms of the series

$$1-\frac{1}{3}-\frac{1}{9}+\frac{1}{27}+\frac{1}{81}-\frac{1}{243}-\frac{1}{729}+\frac{1}{2187}+\dots$$

is

$$\frac{3}{5}\{1-(-1)^n 3^{-2n}\}.$$

36. If $\frac{1}{b-a}, \frac{1}{2b}, \frac{1}{b-c}$ are in A.P., then a, b, c are in G.P.

37. The last term of an A.P. is ten times the first, and the last but one is equal to the sum of the 4th and 5th. Find the number of the terms, and shew that the common difference is equal to the first term.

38. Sum to $2n$ terms each of the series

(1) $1-3+9-27+\dots$;

(2) $1-3+5-7+\dots$,

and write down the last term of each series.

39. Find two numbers whose arithmetic mean exceeds their geometric mean by 2, and whose harmonic mean is one-fifth of the larger number.

40. Find an infinite geometrical progression, whose first term is 1, and in which each term is twice the sum of all the terms that follow it.

41. The arithmetic mean between two numbers is to the geometric mean as 5 to 4, and the difference of their geometric and harmonic means is $\frac{4}{5}$: find the numbers.

42. If x, y, z be in G.P., prove that

$$x^3 y^2 z^2 (x^{-3} + y^{-3} + z^{-3}) = x^3 + y^3 + z^3.$$

18 terms.

es

.....

re in G.P.

first, and the last
find the number of
s equal to the first

can exceeds their
is one-fifth of the

hose first term is
all the terms that

is to the geometric
tric and harmonic

CHAPTER XXXVI.

THE THEORY OF QUADRATIC EQUATIONS.

335. IN Chapter xxv. it was shewn that after suitable reduction every quadratic equation may be written in the form

$$ax^2 + bx + c = 0 \dots\dots\dots(1),$$

and that the solution of the equation is

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \dots\dots\dots(2).$$

We shall now prove some important propositions connected with the roots and coefficients of all equations of which (1) is the type.

336. *A quadratic equation cannot have more than two roots.*

For, if possible, let the equation $ax^2 + bx + c = 0$ have three different roots α, β, γ . Then since each of these values must satisfy the equation, we have

$$a\alpha^2 + b\alpha + c = 0 \dots\dots\dots(1),$$

$$a\beta^2 + b\beta + c = 0 \dots\dots\dots(2),$$

$$a\gamma^2 + b\gamma + c = 0 \dots\dots\dots(3)$$

From (1) and (2), by subtraction,

$$a(\alpha^2 - \beta^2) + b(\alpha - \beta) = 0;$$

divide out by $\alpha - \beta$ which, by hypothesis, is not zero; then

$$a(\alpha + \beta) + b = 0.$$

Similarly from (2) and (3)

$$a(\beta + \gamma) + b = 0;$$

$\therefore$ by subtraction

$$a(\alpha - \gamma) = 0;$$

which is impossible, since, by hypothesis, α is not zero, and α is not equal to γ . Hence there cannot be three different roots.

337. The terms 'unreal', 'imaginary', and 'impossible' are all used in the same sense : namely, to denote expressions which involve the square root of a negative quantity, such as

$$\sqrt{-1}, \sqrt{-3}, \sqrt{-a}.$$

It is important that the student should clearly distinguish between the terms *real* and *rational*, *imaginary* and *irrational*. Thus $\sqrt{25}$ or 5, $3\frac{1}{2}$, $-\frac{5}{8}$ are rational and real; $\sqrt{7}$ is irrational but real; while $\sqrt{-7}$ is irrational and also imaginary.

338. In Art. 335 let the two roots in (2) be denoted by α and β , so that

$$\alpha = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, \quad \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a};$$

then we have the following results :

(1) If $b^2 - 4ac$, the quantity under the radical, is positive, α and β are real and unequal.

(2) If $b^2 - 4ac$ is zero, α and β are real and equal, each reducing in this case to $-\frac{b}{2a}$.

(3) If $b^2 - 4ac$ is negative, α and β are imaginary and unequal.

(4) If $b^2 - 4ac$ is a perfect square, α and β are rational and unequal.

By applying these tests the nature of the roots of any quadratic may be determined without solving the equation.

Example 1. Shew that the equation $2x^2 - 6x + 7 = 0$ cannot be satisfied by any real values of x .

Here $a=2$, $b=-6$, $c=7$; so that

$$b^2 - 4ac = (-6)^2 - 4 \cdot 2 \cdot 7 = -20.$$

Therefore the roots are imaginary.

Example 2. For what value of k will the equation $3x^2 - 6x + k = 0$ have equal roots?

The condition for equal roots gives

$$(-6)^2 - 4 \cdot 3 \cdot k = 0,$$

whence

$$k = 3.$$

Example 3. Shew that the roots of the equation

$$x^2 - 2px + p^2 - q^2 + 2qr - r^2 = 0$$

are rational.

The roots will be rational provided $(-2p)^2 - 4(p^2 - q^2 + 2qr - r^2)$ is a perfect square. But this expression reduces to $4(q^2 - 2qr + r^2)$, or $4(q-r)^2$. Hence the roots are rational.

'impossible' are expressions which such as

early distinguish y and *irrational*. $\sqrt{7}$ is *irrational* *imaginary*.

be denoted by a

$\sqrt{b^2 - 4ac}$;

al, is positive, a

and equal, each

ary and unequal. are rational and

the roots of any equation.

$+7=0$ cannot be

on $3x^2 - 6x + k = 0$

$$339. \text{ Since } a = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, \quad \beta = \frac{-b - \sqrt{b^2 - 4ac}}{2a},$$

we have by addition

$$\begin{aligned} a + \beta &= \frac{-b + \sqrt{b^2 - 4ac} - b - \sqrt{b^2 - 4ac}}{2a}, \\ &= -\frac{2b}{2a} = -\frac{b}{a} \dots\dots\dots(1); \end{aligned}$$

and by multiplication we have

$$\begin{aligned} a\beta &= \frac{(-b + \sqrt{b^2 - 4ac})(-b - \sqrt{b^2 - 4ac})}{4a^2} \\ &= \frac{(-b)^2 - (b^2 - 4ac)}{4a^2} \\ &= \frac{4ac}{4a^2} = \frac{c}{a} \dots\dots\dots(2). \end{aligned}$$

By writing the equation in the form

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0,$$

these results may also be expressed as follows :

In a quadratic equation *where the coefficient of the first term is unity*,

(i) the sum of the roots is equal to the coefficient of x with its sign changed ;

(ii) the product of the roots is equal to the third term.

Note. In any equation the term which does not contain the unknown quantity is frequently called *the absolute term*.

$$340. \text{ Since } -\frac{b}{a} = a + \beta, \text{ and } \frac{c}{a} = a\beta,$$

the equation $x^2 + \frac{b}{a}x + \frac{c}{a}$ may be written

$$x^2 - (a + \beta)x + a\beta = 0 \dots\dots\dots(1).$$

Hence any quadratic may also be expressed in the form

$$x^2 - (\text{sum of roots})x + \text{product of roots} = 0 \dots\dots\dots(2).$$

Again, from (1) we have

$$(x - a)(x - \beta) = 0 \dots\dots\dots(3).$$

We may now easily form an equation with given roots.

Example 1. Form the equation whose roots are 3 and -2.

The equation is $(x-3)(x+2)=0$,

or $x^2 - x - 6 = 0$.

Example 2. Form the equation whose roots are $\frac{3}{7}$ and $-\frac{4}{5}$.

The equation is $\left(x - \frac{3}{7}\right)\left(x + \frac{4}{5}\right) = 0$;

that is, $(7x-3)(5x+4)=0$,

or $35x^2 + 13x - 12 = 0$.

When the roots are irrational it is easier to use the following method :

Example 3. Form the equation whose roots are $2+\sqrt{3}$ and $2-\sqrt{3}$.

We have $\begin{aligned} \text{sum of roots} &= 4, \\ \text{product of roots} &= 1; \end{aligned}$

$\therefore$ the equation is $x^2 - 4x + 1 = 0$,
by using formula (2) of the present article.

341. The results of Art. 339 are most important, and they are generally sufficient to solve problems connected with the roots of quadratics. In such questions *the roots should never be considered singly*, but use should be made of the relations obtained by writing down the sum of the roots, and their product, in terms of the coefficients of the equation.

Example 1. If α and β are the roots of $x^2 - px + q = 0$, find the value of (1) $\alpha^2 + \beta^2$, (2) $\alpha^3 + \beta^3$.

We have

$$\alpha + \beta = p,$$

$$\alpha\beta = q.$$

$$\therefore \alpha^2 + \beta^2 = (\alpha + \beta)^2 - 2\alpha\beta \\ = p^2 - 2q.$$

Again,

$$\alpha^3 + \beta^3 = (\alpha + \beta)(\alpha^2 + \beta^2 - \alpha\beta) \\ = p\{(\alpha + \beta)^2 - 3\alpha\beta\} \\ = p(p^2 - 3q).$$

Example 2. If α, β are the roots of the equation $lx^2 + mx + n = 0$, find the equation whose roots are $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$.

We have

$$\text{sum of roots} = \frac{\alpha}{\beta} + \frac{\beta}{\alpha} = \frac{\alpha^2 + \beta^2}{\alpha\beta},$$

$$\text{product of roots} = \frac{\alpha}{\beta} \cdot \frac{\beta}{\alpha} = 1;$$

3 and -2.

∴ by Art. 340 the required equation is

$$x^2 - \left(\frac{a^2 + \beta^2}{a\beta} \right) x + 1 = 0,$$

 $\frac{3}{7}$ and $-\frac{4}{5}$.

$$\text{or} \quad a\beta x^2 - (a^2 + \beta^2)x + a\beta = 0.$$

$$\text{As in the last example } a^2 + \beta^2 = \frac{m^2 - 2nl}{l^2}, \text{ and } a\beta = \frac{n}{l}.$$

se the following

$$\therefore \text{ the equation is } \frac{n}{l}x^2 - \frac{m^2 - 2nl}{l^2}x + \frac{n}{l} = 0,$$

 $2 + \sqrt{3}$ and $2 - \sqrt{3}$.

$$\text{or} \quad nlx^2 - (m^2 - 2nl)x + nl = 0.$$

Example 3. Find the condition that the roots of the equation $ax^2 + bx + c = 0$ should be (1) equal in magnitude and opposite in sign, (2) reciprocals.

The roots will be equal in magnitude and opposite in sign if their sum is zero; therefore $-\frac{b}{a} = 0$, or $b = 0$.

rtant, and they
ected with the
should never be
ne relations ob-
d their product,

Again, the roots will be reciprocals when their product is unity; therefore $\frac{c}{a} = 1$, or $c = a$.

Example 4. Find the relation which must subsist between the coefficients of the equation $px^2 + qx + r = 0$, when one root is three times the other.

 $+q=0$, find the

$$\text{We have} \quad a + \beta = -\frac{q}{p}, \quad a\beta = \frac{r}{p};$$

but since $a = 3\beta$, we obtain by substitution

$$4\beta = -\frac{q}{p}, \quad 3\beta^2 = \frac{r}{p}$$

From the first of these equations $\beta^2 = \frac{q^2}{16p^2}$, and from the second $\beta^2 = \frac{r}{3p}$.

 $lx^2 + mx + n = 0$,

$$\therefore \frac{q^2}{16p^2} = \frac{r}{3p},$$

or

$$3q^2 = 16pr,$$

which is the required condition.

342. The following example illustrates a useful application of the results proved in Art. 338.

Example. If x is a real quantity, prove that the expression $\frac{x^2+2x-11}{2(x-3)}$ can have all numerical values except such as lie between 2 and 6.

Let the given expression be represented by y , so that

$$\frac{x^2+2x-11}{2(x-3)} = y;$$

then multiplying up and transposing, we have

$$x^2+2x(1-y)+6y-11=0.$$

This is a quadratic equation, and if x is to have real values $4(1-y)^2-4(6y-11)$ must be positive; or simplifying and dividing by 4, $y^2-8y+12$ must be positive; that is, $(y-6)(y-2)$ must be positive. Hence the factors of this product must be both positive, or both negative. In the former case y is greater than 6; in the latter y is less than 2. Therefore y cannot lie between 2 and 6, but may have any other value.

In this example it will be noticed that the expression $y^2-8y+12$ is positive so long as y does not lie between the roots of the corresponding quadratic equation $y^2-8y+12=0$.

This is a particular case of the general proposition investigated in the next article.

343. For all real values of x the expression ax^2+bx+c has the same sign as a , except when the roots of the equation $ax^2+bx+c=0$ are real and unequal, and x lies between them.

CASE I. Suppose that the roots of the equation

$$ax^2+bx+c=0$$

are real; denote them by α and β , and let α be the greater.

$$\begin{aligned} \text{Then } ax^2+bx+c &= a\left(x^2+\frac{b}{a}x+\frac{c}{a}\right) \\ &= a\{x^2-(\alpha+\beta)x+\alpha\beta\} \quad [\text{Art. 339.}] \\ &= a(x-\alpha)(x-\beta). \end{aligned}$$

Now if x is greater than α or less than β , the factors $x-\alpha$, $x-\beta$ are either both positive or both negative; therefore the expression $(x-\alpha)(x-\beta)$ is positive, and ax^2+bx+c has the same sign as a . But if x lies between α and β , the expression

$$(x-\alpha)(x-\beta)$$

is negative, and the sign of ax^2+bx+c is opposite to that of a .

CASE II. If a and β are equal, then

$$ax^2 + bx + c = a(x - a)^2,$$

and $(x - a)^2$ is positive for all real values of x ; hence $ax^2 + bx + c$ has the same sign as a .

CASE III. Suppose that the equation $ax^2 + bx + c = 0$ has imaginary roots; then

$$\begin{aligned} ax^2 + bx + c &= a \left\{ x^2 + \frac{b}{a}x + \frac{c}{a} \right\} \\ &= a \left\{ \left(x + \frac{b}{2a} \right)^2 + \frac{4ac - b^2}{4a^2} \right\}; \end{aligned}$$

but since $b^2 - 4ac$ is negative [Art. 338], the expression

$$\left(x + \frac{b}{2a} \right)^2 + \frac{4ac - b^2}{4a^2}$$

is positive for all real values of x ; therefore $ax^2 + bx + c$ has the same sign as a .

EXAMPLES XXXVI.

Find (without actual solution) the nature of the roots of the following equations:

1. $x^2 + x - 870 = 0$.
2. $8 + 6x = 5x^2$.
3. $\frac{1}{2}x^2 = 14 - 3x^2$.
4. $x^2 + 7 = 4x$.
5. $2x = x^2 + 5$.
6. $(x + 2)^2 = 4x + 15$.

Form the equations whose roots are

7. 5, -3.
8. -9, -11.
9. $a + b$, $a - b$.
10. $\frac{3}{2}$, $\frac{5}{6}$.
11. $\frac{2}{3}a$, $-\frac{4}{5}a$.
12. 0, $\frac{7}{8}$.

13. If the equation $x^2 + 2(1 + k)x + k^2 = 0$ has equal roots, what is the value of k ?

14. Prove that the equation $3mx^2 - (2m + 3n)x + 2n = 0$ has rational roots.

15. Without solving the equation $3x^2 - 4x - 1 = 0$, find the sum, the difference, and the sum of the squares of the roots.

16. Shew that the roots of $a(x^2 - 1) = (b - c)x$ are always real.

Form the equations whose roots are

17. $3+\sqrt{5}$, $3-\sqrt{5}$. 18. $-2+\sqrt{3}$, $-2-\sqrt{3}$. 19. $-\frac{a}{5}$, $\frac{b}{6}$.
 20. $\frac{1}{2}(4\pm\sqrt{7})$. 21. $\frac{a+b}{a-b}$, $\frac{a-b}{a+b}$. 22. $\frac{a}{2b}$, $\frac{b}{2a}$.

If α , β are the roots of the equation $px^2+qx+r=0$, find the values of

23. $\alpha^2+\beta^2$. 24. $(\alpha-\beta)^2$. 25. $\alpha^2\beta+\alpha\beta^2$.
 26. $\alpha^4+\beta^4$. 27. $\alpha^5\beta^2+\alpha^2\beta^5$. 28. $\frac{\alpha^2}{\beta}+\frac{\beta^2}{\alpha}$.
 29. If α , β are the roots of $x^2-px+q=0$, and α^3 , β^3 the roots of $x^2-Px+Q=0$, find P and Q in terms of p and q .
 30. If α , β are the roots of $x^2-ax+b=0$, find the equation whose roots are $\frac{\alpha}{\beta^2}$, $\frac{\beta}{\alpha^2}$.

31. Find the condition that one root of the equation
 $ax^2+bx+c=0$

may be double the other.

32. Form an equation whose roots shall be the cubes of the roots of the equation $2x(x-a)=a^2$.

33. Prove that the roots of the equation

$$(a+b)x^2-(a+b+c)x+\frac{c}{2}=0$$

are always real.

34. Shew that $(a+b+c)x^2-2(a+b)x+(a+b-c)=0$ has rational roots.

35. Form an equation whose roots shall be the arithmetic and harmonic means between the roots of $x^2-px+q=0$.

36. In the equation $px^2+qx+r=0$ the roots are in the ratio of l to m , prove that

$$(l^2+m^2)pr+lm(2pr-q^2)=0.$$

37. Shew that if x is real the expression $\frac{x^2-15}{2x-8}$ cannot lie between 3 and 5.

38. If x is real, prove that $\frac{3x^2+2}{x^2-2x-1}$ can have all values except such as lie between 2 and $-\frac{3}{2}$.

$$19. -\frac{a}{5}, \frac{b}{6}.$$

$$22. \frac{a}{2b}, \frac{b}{2a}.$$

$x+r=0$, find the

$$25. a^2\beta + a\beta^2.$$

28. $\frac{a^2}{\beta} + \frac{\beta^2}{a}$.
 β^3 the roots of
 and q .
 equation whose

on
 bes of the roots

arithmetic and
 $q=0$.
 in the ratio of

not lie between

l values except

CHAPTER XXXVII.

PERMUTATIONS AND COMBINATIONS.

344. EACH of the *arrangements* which can be made by taking some or all of a number of things is called a **permutation**.

Each of the *groups* or *selections* which can be made by taking some or all of a number of things is called a **combination**.

Thus the *permutations* which can be made by taking the letters a, b, c, d two at a time are twelve in number: namely,

$$ab, ac, ad, bc, bd, cd, \\ ba, ca, da, cb, db, dc;$$

each of these presenting a different *arrangement* of two letters.

The *combinations* which can be made by taking the letters a, b, c, d two at a time are six in number: namely,

$$ab, ac, ad, bc, bd, cd;$$

each of these presenting a different *selection* of two letters.

From this it appears that in forming *combinations* we are only concerned with the number of things each selection contains; whereas in forming *permutations* we have also to consider the order of the things which make up each arrangement; for instance, if from four letters a, b, c, d we make a selection of three, such as abc , this single combination admits of being arranged in the following ways:

$$abc, acb, bca, bac, cab, cba,$$

and so gives rise to six different permutations.

345. Before discussing the general propositions of this chapter the following important principle should be carefully noticed.

If one operation can be performed in m ways, and (when it has been performed in any one of these ways) a second operation can then be performed in n ways; the number of ways of performing the two operations will be $m \times n$.

If the first operation be performed in *any one way*, we can associate with this any of the n ways of performing the second operation : and thus we shall have n ways of performing the two operations without considering more than *one way* of performing the first ; and so, corresponding to *each* of the m ways of performing the first operation, we shall have n ways of performing the two ; hence altogether the number of ways in which the two operations can be performed is represented by the product $m \times n$.

Example. There are 10 steamers plying between Liverpool and Dublin ; in how many ways can a man go from Liverpool to Dublin and return by a different steamer ?

There are *ten* ways of making the first passage ; and with each of these there is a choice of *nine* ways of returning (since the man is not to come back by the same steamer) ; hence the number of ways of making the two journeys is 10×9 , or 90.

This principle may easily be extended to the case in which there are more than two operations each of which can be performed in a given number of ways.

346. *To find the number of permutations of n dissimilar things taken r at a time.*

This is the same thing as finding the number of ways in which we can fill up r places when we have n different things at our disposal.

The first place may be filled up in n ways, for any one of the n things may be taken ; when it has been filled up in any one of these ways, the second place can then be filled up in $n - 1$ ways ; and since each way of filling up the first place can be associated with each way of filling up the second, the number of ways in which the first two places can be filled up is given by the product $n(n - 1)$. And when the first two places have been filled up in any way, the third place can be filled up in $n - 2$ ways. And reasoning as before, the number of ways in which three places can be filled up is $n(n - 1)(n - 2)$.

Proceeding thus, and noticing that a new factor is introduced with each new place filled up, and that at any stage the number of factors is the same as the number of places filled up, we shall have the number of ways in which r places can be filled up equal to

$$n(n - 1)(n - 2) \dots \dots \text{to } r \text{ factors ;}$$

and the r^{th} factor is $n - (r - 1)$, or $n - r + 1$.

one way, we can forming the second performing the two way of performing the m ways of performing of performing in which the two by the product

een Liverpool and Liverpool to Dublin

and with each of (since the man is the number of ways

the case in which which can be per-

dissimilar things

number of ways in different things at

or any one of the up in any one of up in $n-1$ ways; can be associated number of ways in n by the product been filled up in -2 ways. And which three places

or is introduced age the number led up, we shall can be filled up

Therefore the number of permutations of n things taken r at a time is

$$n(n-1)(n-2) \dots (n-r+1).$$

COR. The number of permutations of n things taken all at a time is

$$n(n-1)(n-2) \dots \text{to } n \text{ factors,}$$

$$\text{or } n(n-1)(n-2) \dots 3.2.1.$$

It is usual to denote this product by the symbol $[n]$, which is read "factorial n ." Also the symbol $n!$ is sometimes used for $[n]$.

347. We shall in future denote the number of permutations of n things taken r at a time by the symbol ${}^nP_r$, so that

$${}^nP_r = n(n-1)(n-2) \dots (n-r+1);$$

$$\text{also } {}^nP_n = [n].$$

In working numerical examples it is useful to notice that the suffix in the symbol ${}^nP_r$ always denotes the number of factors in the formula we are using.

Example 1. Four persons enter a railway carriage in which there are six seats; in how many ways can they take their places?

The first person may seat himself in 6 ways; and then the second person in 5; the third in 4; and the fourth in 3; and since each of these ways may be associated with each of the others, the required answer is $6 \times 5 \times 4 \times 3$, or 360.

Example 2. How many different numbers can be formed by using six out of the nine digits 1, 2, 3, ... 9?

Here we have 9 different things and we have to find the number of permutations of them taken 6 at a time;

$$\begin{aligned} \therefore \text{the required result} &= {}^9P_6 \\ &= 9 \times 8 \times 7 \times 6 \times 5 \times 4 \\ &= 60480. \end{aligned}$$

348. To find the number of combinations of n dissimilar things taken r at a time.

Let ${}^nC_r$ denote the required number of combinations.

Then each of these combinations consists of a group of r dissimilar things which can be arranged among themselves in $[r]$ ways. [Art. 346. Cor.]

Hence ${}^nC_r \times [r]$ is equal to the number of arrangements of n things taken r at a time; that is,

$${}^nC_r \times [r] = {}^nP_r = n(n-1)(n-2) \dots (n-r+1);$$

$$\therefore {}^nC_r = \frac{n(n-1)(n-2) \dots (n-r+1)}{[r]}.$$

Cor. This formula for ${}^nC_r$ may also be written in a different form; for if we multiply the numerator and the denominator by $\frac{n-r}{r}$ we obtain

$$\frac{n(n-1)(n-2)\dots(n-r+1) \times \frac{n-r}{r}}{\frac{n-r}{r}}, \text{ or } \frac{n}{r} \frac{n-r}{n-r};$$

since $n(n-1)(n-2)\dots(n-r+1) \times \frac{n-r}{r} = n$.

Example. From 12 books in how many ways can a selection of 5 be made, (1) when one specified book is always included, (2) when one specified book is always excluded?

(1) Since the specified book is to be included in every selection, we have only to choose 4 out of the remaining 11.

Hence the number of ways = ${}^{11}C_4 = \frac{11 \times 10 \times 9 \times 8}{1 \times 2 \times 3 \times 4} = 330$.

(2) Since the specified book is always to be excluded, we have to select the 5 books out of the remaining 11.

Hence the number of ways = ${}^{11}C_5 = \frac{11 \times 10 \times 9 \times 8 \times 7}{1 \times 2 \times 3 \times 4 \times 5} = 462$.

349. The number of combinations of n things r at a time is equal to the number of combinations of n things $n-r$ at a time.

In making all the possible combinations of n things, to each group of r things we select, there is left a corresponding group of $n-r$ things; that is, the number of combinations of n things r at a time is the same as the number of combinations of n things $n-r$ at a time;

$$\therefore {}^nC_r = {}^nC_{n-r}.$$

This result is frequently useful in enabling us to abridge arithmetical work.

Example. Out of 14 men in how many ways can an eleven be chosen?

The required number = ${}^{14}C_{11}$

$$= {}^{14}C_3 = \frac{14 \times 13 \times 12}{1 \times 2 \times 3} = 364.$$

If we had made use of the formula ${}^{14}C_{11}$, we should have had to reduce an expression whose numerator and denominator each contained 11 factors.

350. In the examples which follow it is important to notice that the formula for *permutations* should not be used until the suitable *selections* required by the question have been made.

ten in a different
the denominator by

$$\frac{n}{r(n-r)};$$

an a selection of 5
included, (2) when

in every selection,

$$= 330.$$

excluded, we have

$$\frac{3 \times 7}{5} = 462.$$

ngs r at a time is
 $n - r$ at a time.

n things, to each
ponding group of
ions of n things
ations of n things

g us to abridge

can an eleven be

ould have had to
minator each con-

portant to notice
be used until the
e been made.

Example 1. From 7 Englishmen and 4 Americans a committee of 6 is to be formed: in how many ways can this be done, (1) when the committee contains exactly 2 Americans, (2) at least 2 Americans?

(1) The number of ways in which the Americans can be chosen is ${}^4C_2$; and the number of ways in which the Englishmen can be chosen is ${}^7C_4$. Each of the first groups can be associated with each of the second; hence

the required number of ways = ${}^4C_2 \times {}^7C_4$

$$= \frac{4}{2} \frac{3}{2} \times \frac{7}{4} \frac{6}{3} = \frac{7}{2} \frac{7}{2} \frac{3}{1} = 210.$$

(2) We shall exhaust all the suitable combinations by forming all the groups containing 2 Americans and 4 Englishmen; then 3 Americans and 3 Englishmen; and lastly 4 Americans and 2 Englishmen.

The sum of the three results will give the answer. Hence the required number of ways = ${}^4C_2 \times {}^7C_4 + {}^4C_3 \times {}^7C_3 + {}^4C_4 \times {}^7C_2$

$$= \frac{4}{2} \frac{3}{2} \times \frac{7}{4} \frac{6}{3} + \frac{4}{3} \times \frac{7}{3} \frac{6}{2} + 1 \times \frac{7}{2} \frac{6}{1} \\ = 210 + 140 + 21 = 371.$$

In this example we have only to make use of the suitable formulæ for combinations, for we are not concerned with the possible arrangements of the members of the committee among themselves.

Example 2. Out of 7 consonants and 4 vowels, how many words can be made each containing 3 consonants and 2 vowels?

The number of ways of choosing the three consonants is ${}^7C_3$, and the number of ways of choosing the 2 vowels is ${}^4C_2$; and since each of the first groups can be associated with each of the second, the number of combined groups, each containing 3 consonants and 2 vowels, is ${}^7C_3 \times {}^4C_2$.

Further, each of these groups contains 5 letters, which may be arranged among themselves in 5 ways. Hence

$$\text{the required number of words} = \frac{7}{3} \frac{6}{2} \times \frac{4}{2} \frac{3}{1} \times 5 \\ = 5 \times 7 = 25200.$$

EXAMPLES XXXVII. a.

- Find the value of ${}^5P_4$, ${}^7P_6$, ${}^8C_5$, ${}^{25}C_{25}$.
- How many different arrangements can be made by taking (1) five, (2) all of the letters of the word *soldier*?
- If ${}^nC_3 : {}^{n-1}C_4 = 8 : 5$, find n .

4. How many different selections of four coins can be made from a bag containing a sovereign, a half-sovereign, a half-crown, a florin, a shilling, a franc, a sixpence, a penny, and a farthing?
5. How many numbers between 3000 and 4000 can be made with the digits 9, 3, 4, 6?
6. In how many ways can the letters of the word *volume* be arranged if the vowels can only occupy the even places?
7. If the number of permutations of n things four at a time is fourteen times the number of permutations of $n-2$ things three at a time, find n .
8. From 5 masters and 10 boys how many committees can be selected containing 3 masters and 6 boys?
9. If ${}^{20}C_7 = {}^{20}C_{10}$, find ${}^nC_{12}$, ${}^{18}C_7$.
10. Out of the twenty-six letters of the alphabet in how many ways can a word be made consisting of five different letters, two of which must be a and e ?
11. How many words can be formed by taking 3 consonants and 2 vowels from an alphabet containing 21 consonants and 5 vowels?
12. A railway carriage will accommodate 5 passengers on each side: in how many ways can 10 persons take their seats when two of them decline to face the engine, and a third cannot travel backwards?

351. Hitherto, in the formulæ we have proved, the things have been regarded as *unlike*. Before considering cases in which some one or more sets of things may be *like*, it is necessary to point out exactly in what sense the words *like* and *unlike* are used. When we speak of things being *dissimilar*, *different*, *unlike*, we imply that the things are *visibly unlike*, so as to be easily distinguishable from each other. On the other hand we shall always use the term *like* things to denote such as are alike to the eye and cannot be distinguished from each other. For instance, in Ex. 2, Art. 350, the consonants and the vowels may be said each to consist of a group of things united by a common characteristic, and thus in a certain sense to be of the same kind; but they cannot be regarded as like things, because there is an individuality existing among the things of each group which makes them easily distinguishable from each other. Hence, in the final stage of the example we considered each group to consist of five *dissimilar* things and therefore capable of [5] arrangements among themselves. [Art. 346. Cor.]

352. To find the number of ways in which n things may be arranged among themselves, taking them all at a time, when p of the things are exactly alike of one kind, q of them exactly alike of another kind, r of them exactly alike of a third kind, and the rest all different.

Let there be n letters; suppose p of them to be a , q of them to be b , r of them to be c , and the rest to be unlike.

Let x be the required number of permutations; then if the p letters a were replaced by p unlike letters different from any of the rest, from any one of the x permutations, without altering the position of any of the remaining letters, we could form $|p$ new permutations. Hence if this change were made in each of the x permutations, we should obtain $x \times |p$ permutations.

Similarly, if the q letters b were replaced by q unlike letters, the number of permutations would be $x \times |p \times |q$.

In like manner, by replacing the r letters c by r unlike letters, we should finally obtain $x \times |p \times |q \times |r$ permutations.

But the things are now all different, and therefore admit of $|n$ permutations among themselves. Hence

$$x \times |p \times |q \times |r = |n;$$

that is,

$$x = \frac{|n}{|p |q |r};$$

which is the required number of permutations.

Any case in which the things are not all different may be treated similarly.

Example 1. How many different permutations can be made out of the letters of the word *assassination* taken all together?

We have here 13 letters of which 4 are s , 3 are a , 2 are i , and 2 are n . Hence the number of permutations

$$\begin{aligned} &= \frac{|13}{|4 |3 |2 |2} \\ &= 13 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 3 \cdot 5 \\ &= 1001 \times 10800 = 10810800. \end{aligned}$$

Example 2. How many numbers can be formed with the digits 1, 2, 3, 4, 3, 2, 1, so that the odd digits always occupy the odd places?

The odd digits 1, 3, 3, 1 can be arranged in their four places in

$$\frac{\frac{4}{2}}{2} \text{ ways} \dots\dots\dots (1).$$

The even digits 2, 4, 2 can be arranged in their three places in

$$\frac{\frac{3}{2}}{2} \text{ ways} \dots\dots\dots (2).$$

Each of the ways in (1) can be associated with each of the ways in (2).

$$\text{Hence the required number} = \frac{\frac{4}{2}}{2} \times \frac{\frac{3}{2}}{2} = 6 \times 3 = 18.$$

353. To find the number of permutations of n things r at a time, when each thing may be repeated once, twice, ... up to r times in any arrangement.

Here we have to consider the number of ways in which r places can be filled up when we have n different things at our disposal, each of the n things being used as often as we please in any arrangement.

The first place may be filled up in n ways, and, when it has been filled up in any one way, the second place may also be filled up in n ways, since we are not precluded from using the same thing again. Therefore the number of ways in which the first two places can be filled up is $n \times n$ or n^2 .

The third place can also be filled up in n ways, and therefore the first three places in n^3 ways.

Proceeding in this manner, and noticing that at any stage the index of n is always the same as the number of places filled up, we shall have the number of ways in which the r places can be filled up equal to n^r .

Example. In how many ways can 5 prizes be given away to 4 boys, when each boy is eligible for all the prizes?

Any one of the prizes can be given in 4 ways; and then any one of the remaining prizes can also be given in 4 ways, since it may be obtained by the boy who has already received a prize. Thus two prizes can be given away in 4^2 ways, three prizes in 4^3 ways, and so on. Hence the 5 prizes can be given away in 4^5 , or 1024 ways.

354. To find for what value of r the number of combinations of n things r at a time is greatest.

$$\text{Since } {}^nC_r = \frac{n(n-1)(n-2) \dots (n-r+2)(n-r+1)}{1 \cdot 2 \cdot 3 \dots (r-1)r}$$

$$\text{and } {}^nC_{r-1} = \frac{n(n-1)(n-2) \dots (n-r+2)}{1 \cdot 2 \cdot 3 \dots (r-1)};$$

$$\therefore {}^nC_r = {}^nC_{r-1} \times \frac{n-r+1}{r}.$$

The multiplying factor $\frac{n-r+1}{r}$ may be written $\frac{n+1}{r} - 1$, which shews that it decreases as r increases. Hence as r receives the values 1, 2, 3, ... in succession, ${}^nC_r$ is continually increased, until $\frac{n+1}{r} - 1$ becomes equal to 1 or less than 1.

$$\text{Now } \frac{n+1}{r} - 1 > 1, \text{ so long as } \frac{n+1}{r} > 2; \text{ that is, } \frac{n+1}{2} > r.$$

We have to choose the greatest value of r consistent with this inequality.

(1) Let n be even, and equal to $2m$; then

$$\frac{n+1}{2} = \frac{2m+1}{2} = m + \frac{1}{2};$$

and for all values of r up to m inclusive this is greater than r .

Hence by putting $r = m = \frac{n}{2}$, we find that the greatest number of combinations is ${}^nC_{\frac{n}{2}}$.

(2) Let n be odd, and equal to $2m+1$; then

$$\frac{n+1}{2} = \frac{2m+2}{2} = m+1;$$

and for all values of r up to m inclusive this is greater than r ; but when $r = m+1$ the multiplying factor becomes equal to 1, and

$${}^nC_{m+1} = {}^nC_m; \text{ that is, } {}^nC_{\frac{n+1}{2}} = {}^nC_{\frac{n-1}{2}};$$

and therefore the number of combinations is greatest when the things are taken $\frac{n+1}{2}$, or $\frac{n-1}{2}$ at a time; the result being the same in the two cases.

EXAMPLES XXXVII. b.

1. Find the number of permutations which can be made from all the letters of the words,

(1) *irresistible*,

(2) *phenomenon*,

(3) *tittle-tattle*.

2. How many different numbers can be formed by using the seven digits 2, 3, 4, 3, 3, 1, 2? How many with the digits 2, 3, 4, 3, 3, 0, 2?

3. How many words can be formed from the letters of the word *Simoom*, so that vowels and consonants occur alternately in each word?

4. A telegraph has 5 arms and each arm has 4 distinct positions, including the position of rest: find the total number of signals that can be made.

5. In how many ways can n things be given to m persons, when there is no restriction as to the number of things each may receive?

6. How many different arrangements can be made out of the letters of the expression $a^6b^3c^6$ when written at full length?

7. There are four copies each of 3 different volumes; find the number of ways in which they can be arranged on one shelf.

8. In how many ways can 6 persons form a ring? Find the number of ways in which 4 gentlemen and 4 ladies can sit at a round table so that no two gentlemen sit together.

9. In how many ways can a word of 4 letters be made out of the letters a, b, e, c, d, o , when there is no restriction as to the number of times a letter is repeated in each word?

10. How many arrangements can be made out of the letters of the word *Toulouse*, so that the consonants occupy the first, fourth, and seventh places?

11. A boat's crew consists of eight men, of whom one can only row on bow side and one only on stroke side: in how many ways can the crew be arranged?

12. Shew that ${}^{n+1}C_r = {}^nC_r + {}^nC_{r-1}$.

13. A cricket eleven has to be chosen from 13 men of whom only 4 can bowl: in how many ways can the team be made up so as to include at least 2 bowlers?

14. In how many ways can n men be arranged in a row if two specified men are neither of them to be at either extremity of the row?

CHAPTER XXXVIII.

BINOMIAL THEOREM.

355. It may be shewn by actual multiplication that

$$\begin{aligned} & (x+a)(x+b)(x+c)(x+d) \\ &= x^4 + (a+b+c+d)x^3 + (ab+ac+ad+bc+bd+cd)x^2 \\ &+ (abc+abd+acd+bcd)x + abcd \dots\dots\dots (1). \end{aligned}$$

We may, however, write down this result by inspection; for the complete product consists of the sum of a number of partial products each of which is formed by multiplying together four letters, *one* being taken from *each* of the four factors. If we examine the way in which the various partial products are formed, we see that

(1) the term x^4 is formed by taking the letter x out of *each* of the factors.

(2) the terms involving x^3 are formed by taking the letter x out of *any three* factors, in every way possible, and *one* of the letters a, b, c, d out of the remaining factor.

(3) the terms involving x^2 are formed by taking the letter x out of *any two* factors, in every way possible, and *two* of the letters a, b, c, d out of the remaining factors.

(4) the terms involving x are formed by taking the letter x out of *any one* factor, and *three* of the letters a, b, c, d out of the remaining factors.

(5) the term independent of x is the product of all the letters a, b, c, d .

Example. Find the value of $(x-2)(x+3)(x-5)(x+9)$.

The product

$$\begin{aligned} &= x^4 + (-2+3-5+9)x^3 + (-6+10-18-15+27-45)x^2 \\ &+ (30-54+90-135)x + 270 \\ &= x^4 + 5x^3 - 47x^2 - 69x + 270. \end{aligned}$$

356. If in equation (1) of the preceding article we suppose $b=c=d=a$, we obtain

$$(x+a)^4 = x^4 + 4ax^3 + 6a^2x^2 + 4a^3x + a^4.$$

We shall now employ the same method to prove a formula known as the **Binomial Theorem**, by which any binomial of the form $x+a$ can be raised to any assigned positive integral power.

357. To find the expansion of $(x+a)^n$ when n is a positive integer.

Consider the expression

$$(x+a)(x+b)(x+c)\dots(x+k),$$

the number of factors being n .

The expansion of this expression is the continued product of the n factors, $x+a, x+b, x+c, \dots, x+k$, and every term in the expansion is of n dimensions, being a product formed by multiplying together n letters, one taken from each of these n factors.

The highest power of x is x^n , and is formed by taking the letter x from each of the n factors.

The terms involving x^{n-1} are formed by taking the letter x from any $n-1$ of the factors, and one of the letters $a, b, c, \dots, k$ from the remaining factor; thus the coefficient of x^{n-1} in the final product is the sum of the letters $a, b, c, \dots, k$; denote it by S_1 .

The terms involving x^{n-2} are formed by taking the letter x from any $n-2$ of the factors, and two of the letters $a, b, c, \dots, k$ from the two remaining factors; thus the coefficient of x^{n-2} in the final product is the sum of the products of the letters $a, b, c, \dots, k$ taken two at a time; denote it by S_2 .

And, generally, the terms involving x^{n-r} are formed by taking the letter x from any $n-r$ of the factors, and r of the letters $a, b, c, \dots, k$ from the r remaining factors; thus the coefficient of x^{n-r} in the final product is the sum of the products of the letters $a, b, c, \dots, k$ taken r at a time; denote it by S_r .

The last term in the product is $abc\dots k$; denote it by S_n .

Hence $(x+a)(x+b)(x+c)\dots(x+k)$

$$= x^n + S_1x^{n-1} + S_2x^{n-2} + \dots + S_rx^{n-r} + \dots + S_{n-1}x + S_n.$$

In S_1 the number of terms is n ; in S_2 the number of terms is the same as the number of combinations of n things 2 at a time; that is, ${}^nC_2$; in S_3 the number of terms is ${}^nC_3$; and so on.

Now suppose $b, c, \dots, k$, each equal to a ; then S_1 becomes ${}^nC_1a$; S_2 becomes ${}^nC_2a^2$; S_3 becomes ${}^nC_3a^3$; and so on; thus

$$(x+a)^n = x^n + {}^nC_1ax^{n-1} + {}^nC_2a^2x^{n-2} + {}^nC_3a^3x^{n-3} + \dots + {}^nC_na^n;$$

substituting for " $C_1, C_2, \dots$ " we obtain

$$(x+a)^n = x^n + nax^{n-1} + \frac{n(n-1)}{1 \cdot 2}a^2x^{n-2} + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}a^3x^{n-3} + \dots + a^n,$$

the series containing $n+1$ terms.

This is the *Binomial Theorem*, and the expression on the right is said to be **the expansion** of $(x+a)^n$.

358. The coefficients in the expansion of $(x+a)^n$ are very conveniently expressed by the symbols " $C_1, C_2, C_3, \dots, C_n$." We shall, however, sometimes further abbreviate them by omitting n , and writing $C_1, C_2, C_3, \dots, C_n$. With this notation we have

$$(x+a)^n = x^n + C_1ax^{n-1} + C_2a^2x^{n-2} + C_3a^3x^{n-3} + \dots + C_na^n.$$

If we write $-a$ in the place of a , we obtain

$$(x-a)^n = x^n + C_1(-a)x^{n-1} + C_2(-a)^2x^{n-2} + C_3(-a)^3x^{n-3} + \dots + C_n(-a)^n \\ = x^n - C_1ax^{n-1} + C_2a^2x^{n-2} - C_3a^3x^{n-3} + \dots + (-1)^nC_na^n.$$

Thus the terms in the expansion of $(x+a)^n$ and $(x-a)^n$ are *numerically* the same, but in $(x-a)^n$ they are alternately positive and negative, and the last term is positive or negative according as n is even or odd.

Example 1. Find the expansion of $(x+y)^6$.

By the formula, the expansion

$$= x^6 + {}^6C_1x^5y + {}^6C_2x^4y^2 + {}^6C_3x^3y^3 + {}^6C_4x^2y^4 + {}^6C_5xy^5 + {}^6C_6y^6 \\ = x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + y^6,$$

on calculating the values of ${}^6C_1, {}^6C_2, {}^6C_3, \dots$.

Example 2. Find the expansion of $(a-2x)^7$.

$(a-2x)^7 = a^7 - {}^7C_1a^6(2x) + {}^7C_2a^5(2x)^2 - {}^7C_3a^4(2x)^3 + \dots$ to 8 terms.

Now remembering that " $C_r = C_{n-r}$," after calculating the coefficients up to ${}^7C_3$, the rest may be written down at once; for ${}^7C_4 = {}^7C_3$; ${}^7C_5 = {}^7C_2$; and so on. Hence

$$(a-2x)^7 = a^7 - 7a^6(2x) + \frac{7 \cdot 6}{1 \cdot 2}a^5(2x)^2 - \frac{7 \cdot 6 \cdot 5}{1 \cdot 2 \cdot 3}a^4(2x)^3 + \dots \\ = a^7 - 7a^6(2x) + 21a^5(2x)^2 - 35a^4(2x)^3 + 35a^3(2x)^4 \\ - 21a^2(2x)^5 + 7a(2x)^6 - (2x)^7 \\ = a^7 - 14a^6x + 84a^5x^2 - 280a^4x^3 + 560a^3x^4 \\ - 672a^2x^5 + 448ax^6 - 128x^7.$$

359. In the expansion of $(x+a)^n$, the coefficient of the second term is ${}^nC_1$; of the third term is ${}^nC_2$; of the fourth term is ${}^nC_3$; and so on; the suffix in each term being one less than the number of the term to which it applies; hence ${}^nC_r$ is the coefficient of the $(r+1)^{\text{th}}$ term. This is called the **general term**, because by giving to r different numerical values any of the coefficients may be found from ${}^nC_r$; and by giving to x and a their appropriate indices any assigned term may be obtained. Thus the $(r+1)^{\text{th}}$ term may be written

$${}^nC_r x^{n-r} a^r, \text{ or } \frac{n(n-1)(n-2) \dots (n-r+1)}{r!} x^{n-r} a^r.$$

In applying this formula to any particular case, it should be observed that the index of a is the same as the suffix of C , and that the sum of the indices of x and a is n .

Example 1. Find the fifth term of $(a+2x^3)^{17}$.

$$\begin{aligned} \text{The required term} &= {}^{17}C_4 a^{13} (2x^3)^4 \\ &= \frac{17 \cdot 16 \cdot 15 \cdot 14}{1 \cdot 2 \cdot 3 \cdot 4} \times 16 a^{13} x^{12} \\ &= 38080 a^{13} x^{12}. \end{aligned}$$

Example 2. Find the fourteenth term of $(3-a)^{15}$.

$$\begin{aligned} \text{The required term} &= {}^{15}C_{13} (3)^2 (-a)^{13} \\ &= {}^{15}C_2 \times (-9a^{13}) \quad [\text{Art. 349.}] \\ &= -945a^{13}. \end{aligned}$$

360. The simplest form of the binomial theorem is the expansion of $(1+x)^n$. This is obtained from the general formula of Art. 357, by writing 1 in the place of x , and x in the place of a . Thus

$$\begin{aligned} (1+x)^n &= 1 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_r x^r + \dots + {}^nC_n x^n \\ &= 1 + nx + \frac{n(n-1)}{1 \cdot 2} x^2 + \dots + x^n; \end{aligned}$$

the general term being

$$\frac{n(n-1)(n-2) \dots (n-r+1)}{r!} x^r.$$

361. The expansion of a binomial may always be made to depend upon the case in which the first term is unity; thus

$$\begin{aligned} (x+y)^n &= \left\{ x \left(1 + \frac{y}{x} \right) \right\}^n \\ &= x^n (1+z)^n, \text{ where } z = \frac{y}{x}. \end{aligned}$$

of the second
term is ${}^nC_3$;
less than the
 ${}^nC_r$ is the co-
efficient of the
general term,
any of the
terms to x and a
be obtained.

$x^n - r a^r$.

, it should be
of C , and that

Example. Find the coefficient of x^{16} in the expansion of $(x^2 - 2x)^{10}$.

We have $(x^2 - 2x)^{10} = x^{20} \left(1 - \frac{2}{x}\right)^{10}$;

and, since x^{20} multiplies every term in the expansion of $\left(1 - \frac{2}{x}\right)^{10}$,
we have in this expansion to seek the coefficient of the term which
contains $\frac{1}{x^4}$.

Hence the required coefficient $= {}^{10}C_4 (-2)^4$

$$= \frac{10 \cdot 9 \cdot 8 \cdot 7}{1 \cdot 2 \cdot 3 \cdot 4} \times 16 \\ = 3360.$$

EXAMPLES XXXVIII. a.

Expand the following binomials :

- | | | |
|---------------------------------------|---------------------------------------|--|
| 1. $(x+2)^4$. | 2. $(x+3)^5$. | 3. $(a+x)^7$. |
| 4. $(a-x)^5$. | 5. $(1-2y)^5$. | 6. $\left(2x + \frac{y}{2}\right)^4$. |
| 7. $\left(2 - \frac{x}{2}\right)^6$. | 8. $\left(a - \frac{3}{b}\right)^7$. | 9. $\left(ax + \frac{y}{a}\right)^9$. |

Write down and simplify :

- The 4th term of $(1+x)^{12}$.
- The 6th term of $(2-y)^8$.
- The 5th term of $(a-5b)^7$.
- The 15th term of $(2x-1)^{17}$.
- The 7th term of $\left(1 - \frac{1}{x}\right)^{10}$.
- The 6th term of $\left(3x + \frac{a}{2}\right)^9$.
- The middle term of $\left(\frac{2}{3}a - \frac{3}{2a}\right)^6$.
- The 23rd term of $\left(x^2 + \frac{b}{x}\right)^{25}$.
- The 10th term of $(x^2 - x)^{17}$.

[Art. 349.]

em is the ex-
neral formula
r in the place

${}^nC_r x^r$

be made to
ity; thus

19. Find the value of $(x - \sqrt{3})^4 + (x + \sqrt{3})^4$.
 20. Expand $(\sqrt{1-x^2}+1)^5 - (\sqrt{1-x^2}-1)^5$.
 21. Find the coefficient of x^{12} in $(x^3+2x)^{10}$.
 22. Find the coefficient of x in $\left(x^2 - \frac{a}{2x}\right)^{14}$.
 23. Find the term independent of x in $\left(2x^2 - \frac{1}{x}\right)^{12}$.
 24. Find the coefficient of x^{-20} in $\left(\frac{x^2}{3} - \frac{2}{x^3}\right)^{15}$.

362. In the expansion of $(1+x)^n$ the coefficients of terms equidistant from the beginning and end are equal.

The coefficient of the $(r+1)^{\text{th}}$ term from the beginning is ${}^nC_r$.

The $(r+1)^{\text{th}}$ term from the end has $n+1-(r+1)$, or $n-r$ terms before it; therefore counting from the beginning it is the $(n-r+1)^{\text{th}}$ term, and its coefficient is ${}^nC_{n-r}$, which has been shewn to be equal to ${}^nC_r$. [Art. 349.] Hence the proposition follows.

363. To find the greatest coefficient in the expansion of $(1+x)^n$.

The coefficient of the general term of $(1+x)^n$ is ${}^nC_r$; and we have only to find for what value of r this is greatest.

By Art. 354, when n is even, the greatest coefficient is ${}^nC_{\frac{n}{2}}$; and when n is odd, it is ${}^nC_{\frac{n-1}{2}}$ or ${}^nC_{\frac{n+1}{2}}$; these two coefficients being equal.

364. To find the greatest term in the expansion of $(x+a)^n$.

We have $(x+a)^n = x^n \left(1 + \frac{a}{x}\right)^n$;

therefore, since x^n multiplies every term in $\left(1 + \frac{a}{x}\right)^n$, it will be sufficient to find the greatest term in this latter expansion.

Let the r^{th} and $(r+1)^{\text{th}}$ be any two consecutive terms. The $(r+1)^{\text{th}}$ term is obtained by multiplying the r^{th} term by $\frac{n-r+1}{r} \cdot \frac{a}{x}$; that is, by $\left(\frac{n+1}{r} - 1\right) \frac{a}{x}$. [Art. 359.]

The factor $\frac{n+1}{r} - 1$ decreases as r increases; hence the $(r+1)^{\text{th}}$ term is not always greater than the r^{th} term, but only until $\left(\frac{n+1}{r} - 1\right) \frac{a}{x}$ becomes equal to 1, or less than 1.

Now $\left(\frac{n+1}{r} - 1\right) \frac{a}{x} > 1$, so long as $\frac{n+1}{r} - 1 > \frac{x}{a}$;

that is $\frac{n+1}{r} > \frac{x}{a} + 1$, or $\frac{(n+1)a}{x+a} > r \dots \dots \dots (1)$.

If $\frac{(n+1)a}{x+a}$ be an integer, denote it by p ; then if $r=p$ the multiplying factor becomes 1, and the $(p+1)^{\text{th}}$ term is equal to the p^{th} ; and these are greater than any other term.

If $\frac{(n+1)a}{x+a}$ be not an integer, denote its integral part by q ; then the greatest value of r consistent with (1) is q ; hence the $(q+1)^{\text{th}}$ term is the greatest.

Since we are only concerned with the *numerically greatest term*, the investigation will be the same for $(x-a)^n$; therefore in any numerical example it is unnecessary to consider the sign of the second term of the binomial. Also it will be found best to work each example independently of the general formula.

Example. Find the greatest term in the expansion of $(1+4x)^8$, when x has the value $\frac{1}{3}$.

Denote the r^{th} and $(r+1)^{\text{th}}$ terms by T_r and T_{r+1} respectively; then

$$T_{r+1} = \frac{8-r+1}{r} \cdot 4x \times T_r = \frac{9-r}{r} \times \frac{4}{3} \times T_r;$$

hence $T_{r+1} > T_r$ so long as $\frac{9-r}{r} \times \frac{4}{3} > 1$;

that is, $36 - 4r > 3r$, or $36 > 7r$.

The greatest value of r consistent with this is 5; hence the greatest term is the sixth, and its value

$$= {}^8C_5 \times \left(\frac{4}{3}\right)^5 = {}^8C_3 \times \left(\frac{4}{3}\right)^5 = \frac{57344}{243}.$$

of terms equi-

inning is ${}^nC_r$
+1), or $n-r$
eginning it is
hich has been
he proposition

expansion of

${}^nC_r$; and we
st.

cient is ${}^nC_{\frac{n}{2}}$;
vo coefficients

f $(x+a)^n$.

)ⁿ, it will be
pansion.

utive terms.
 r^{th} term by

[Art. 359.]

365. To find the sum of the coefficients in the expansion of $(1+x)^n$.

In the identity $(1+x)^n = 1 + C_1x + C_2x^2 + C_3x^3 + \dots + C_nx^n$,
put $x=1$; thus

$$\begin{aligned} 2^n &= 1 + C_1 + C_2 + C_3 + \dots + C_n \\ &= \text{sum of the coefficients.} \end{aligned}$$

COR. $C_1 + C_2 + C_3 + \dots + C_n = 2^n - 1$;

that is, the total number of combinations of n things taking some or all of them at a time is $2^n - 1$.

366. To prove that in the expansion of $(1+x)^n$, the sum of the coefficients of the odd terms is equal to the sum of the coefficients of the even terms.

In the identity $(1+x)^n = 1 + C_1x + C_2x^2 + C_3x^3 + \dots + C_nx^n$,
put $x = -1$; thus

$$\begin{aligned} 0 &= 1 - C_1 + C_2 - C_3 + C_4 - C_5 + \dots; \\ \therefore 1 + C_2 + C_4 + \dots &= C_1 + C_3 + C_5 + \dots \end{aligned}$$

367. The Binomial Theorem may also be applied to expand expressions which contain more than two terms.

Example. Find the expansion of $(x^2 + 2x - 1)^3$.

Regarding $2x - 1$ as a single term, the expansion

$$\begin{aligned} &= (x^2)^3 + 3(x^2)^2(2x-1) + 3x^2(2x-1)^2 + (2x-1)^3 \\ &= x^6 + 6x^5 + 9x^4 - 4x^3 - 9x^2 + 6x - 1, \text{ on reduction.} \end{aligned}$$

368. For a full discussion of the Binomial Theorem when the index is not restricted to positive integral values the student is referred to the *Higher Algebra*, Chap. XIV. It is there shewn that when x is less than unity, the formula

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1 \cdot 2}x^2 + \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}x^3 + \dots$$

is true for any value of n .

When n is negative or fractional the number of terms in the expansion is unlimited, but in any particular case we may write down as many terms as we please, or we may find the coefficient of any assigned term.

the expansion

$$\dots + C_n x^n,$$

things taking

the sum of the
coefficients of

$$\dots + C_n x^n,$$

lied to expand

1)³
otion.

Theorem when
es the student
is there shewn

$$x^3 + \dots$$

terms in the
we may write
the coefficient

Example 1. Expand $(1+x)^{-3}$ to four terms.

$$\begin{aligned} (1+x)^{-3} &= 1 + (-3)x + \frac{(-3)(-3-1)}{1 \cdot 2} x^2 + \frac{(-3)(-3-1)(-3-2)}{1 \cdot 2 \cdot 3} x^3 + \dots \\ &= 1 - 3x + \frac{3 \cdot 4}{1 \cdot 2} x^2 - \frac{3 \cdot 4 \cdot 5}{1 \cdot 2 \cdot 3} x^3 + \dots \\ &= 1 - 3x + 6x^2 - 10x^3 + \dots \end{aligned}$$

Example 2. Expand $(4+3x)^{\frac{2}{3}}$ to four terms.

$$\begin{aligned} (4+3x)^{\frac{2}{3}} &= 4^{\frac{2}{3}} \left(1 + \frac{3x}{4}\right)^{\frac{2}{3}} = 8 \left(1 + \frac{3x}{4}\right)^{\frac{2}{3}} \\ &= 8 \left[1 + \frac{3}{2} \cdot \frac{3x}{4} + \frac{\frac{3}{2}(\frac{3}{2}-1)}{1 \cdot 2} \left(\frac{3x}{4}\right)^2 + \frac{\frac{3}{2}(\frac{3}{2}-1)(\frac{3}{2}-2)}{1 \cdot 2 \cdot 3} \left(\frac{3x}{4}\right)^3 + \dots \right] \\ &= 8 \left[1 + \frac{3}{2} \cdot \frac{3x}{4} + \frac{3}{8} \cdot \frac{9x^2}{16} - \frac{1}{16} \cdot \frac{27x^3}{64} + \dots \right] \\ &= 8 + 9x + \frac{27}{16} x^2 - \frac{27}{128} x^3 + \dots \end{aligned}$$

369. In finding the general term we must now use the formula

$$\frac{n(n-1)(n-2) \dots (n-r+1)}{r!} x^r$$

written in full; for the symbol ${}^nC_r$ cannot be employed when n is fractional or negative.

Example 1. Find the general term in the expansion of $(1+x)^{\frac{1}{2}}$.

$$\begin{aligned} \text{The } (r+1)^{\text{th}} \text{ term} &= \frac{\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2) \dots (\frac{1}{2}-r+1)}{r!} x^r \\ &= \frac{1(-1)(-3)(-5) \dots (-2r+3)}{2^r r!} x^r. \end{aligned}$$

The number of factors in the numerator is r , and $r-1$ of these are negative; therefore, by taking -1 out of each of these negative factors, we may write the above expression

$$(-1)^{r-1} \frac{1 \cdot 3 \cdot 5 \dots (2r-3)}{2^r r!} x^r,$$

Example 2. Find the general term in the expansion of $(1-x)^{-3}$.

$$\begin{aligned}\text{The } (r+1)^{\text{th}} \text{ term} &= \frac{(-3)(-4)(-5) \dots (-3-r+1)}{r!} (-x)^r \\ &= (-1)^r \frac{3 \cdot 4 \cdot 5 \dots (r+2)}{r!} (-1)^r x^r \\ &= (-1)^{2r} \frac{3 \cdot 4 \cdot 5 \dots (r+2)}{1 \cdot 2 \cdot 3 \dots r} x^r \\ &= \frac{(r+1)(r+2)}{1 \cdot 2} x^r,\end{aligned}$$

by removing like factors from the numerator and denominator.

370. The following example illustrates a useful application of the Binomial Theorem.

Example. Find the cube root of 126 to five places of decimals.

$$\begin{aligned}(126)^{\frac{1}{3}} &= (5^3 + 1)^{\frac{1}{3}} = 5 \left(1 + \frac{1}{5^3} \right)^{\frac{1}{3}} \\ &= 5 \left(1 + \frac{1}{3} \cdot \frac{1}{5^3} - \frac{1}{9} \cdot \frac{1}{5^6} + \frac{5}{81} \cdot \frac{1}{5^9} - \dots \right) \\ &= 5 + \frac{1}{3} \cdot \frac{1}{5^2} - \frac{1}{9} \cdot \frac{1}{5^5} + \frac{1}{81} \cdot \frac{1}{5^8} - \dots \\ &= 5 + \frac{1}{3} \cdot \frac{2^2}{10^3} - \frac{1}{9} \cdot \frac{2^5}{10^6} + \frac{1}{81} \cdot \frac{2^7}{10^9} - \dots \\ &= 5 + \frac{.04}{3} - \frac{.00032}{9} + \frac{.0000128}{81} - \dots \\ &= 5 + .013333 \dots - .000035 \dots + \dots \\ &= 5.01329, \text{ to five places of decimals.}\end{aligned}$$

EXAMPLES XXXVIII. b.

In the following expansions find which is the greatest term :

1. $(x+y)^{17}$ when $x=4$, $y=3$.
2. $(x-y)^{26}$ when $x=9$, $y=4$.
3. $(1+x)^4$ when $x=\frac{2}{3}$.

tion of $(1-x)^{-3}$.

$(-x)^r$

x^r

ominator.

ful application

s of decimals.

)

ls.

reatest term :

4. $(a-4b)^{15}$ when $a=12$, $b=2$.
 5. $(7x+2y)^{30}$ when $x=8$, $y=14$.
 6. $(2x+3)^n$ when $x=\frac{5}{2}$, $n=15$.
 7. In the expansion of $(1+x)^{25}$ the coefficients of the $(2r+1)^{\text{th}}$ and $(r+5)^{\text{th}}$ terms are equal; find r .
 8. Find n when the coefficients of the 16^{th} and 26^{th} terms of $(1+x)^n$ are equal.
 9. Find the relation between r and n in order that the coefficients of the $(r+3)^{\text{th}}$ and $(2r-3)^{\text{th}}$ terms of $(1+x)^{3n}$ may be equal.
 10. Find the coefficient of x^m in the expansion of $\left(x^2 + \frac{1}{x}\right)^{2m}$.
 11. Find the middle term of $(1+x)^{2m}$ in its simplest form.
 12. Find the sum of the coefficients of $(x+y)^{16}$.
 13. Find the sum of the coefficients of $(3x+y)^9$.
 14. Find the r^{th} term from the beginning and the r^{th} term from the end of $(a+2x)^n$.
 15. Expand $(a^2+2a+1)^3$ and $(x^3-4x+2)^3$.
- Expand to 4 terms the following expressions :
- | | | |
|-----------------------------|-------------------------------|-------------------------------|
| 16. $(1+x)^{\frac{1}{3}}$. | 17. $(1+x)^{\frac{2}{3}}$. | 18. $(1+x)^{\frac{5}{6}}$. |
| 19. $(1+3x)^{-2}$. | 20. $(1-x^2)^{-3}$. | 21. $(1+3x)^{-4}$. |
| 22. $(2+x)^{-3}$. | 23. $(1+2x)^{-\frac{1}{2}}$. | 24. $(a-2x)^{-\frac{3}{2}}$. |

Write down and simplify :

25. The 5^{th} term and the 10^{th} term of $(1+x)^{-\frac{5}{2}}$.
26. The 3^{rd} term and the 11^{th} term of $(1+2x)^{\frac{1}{2}}$.
27. The 4^{th} term and the $(r+1)^{\text{th}}$ term of $(1+x)^{-2}$.
28. The 7^{th} term and the $(r+1)^{\text{th}}$ term of $(1-x)^{\frac{1}{2}}$.
29. The $(r+1)^{\text{th}}$ term of $(a-bx)^{-1}$, and of $(1-nx)^{\frac{1}{2}}$.

Find to four places of decimals the value of

- | | | | |
|-----------------------|-----------------------|----------------------|---------------------|
| 30. $\sqrt[5]{122}$. | 31. $\sqrt[4]{620}$. | 32. $\sqrt[3]{31}$. | 33. $1+\sqrt{99}$. |
|-----------------------|-----------------------|----------------------|---------------------|

CHAPTER XXXIX.

LOGARITHMS.

371. DEFINITION. The **logarithm** of any number to a given **base** is the index of the power to which the base must be raised in order to equal the given number. Thus if $a^x = N$, x is called the logarithm of N to the base a .

Examples. (1) Since $3^4 = 81$, the logarithm of 81 to base 3 is 4.

(2) Since $10^1 = 10$, $10^2 = 100$, $10^3 = 1000$, ...

the natural numbers 1, 2, 3, ... are respectively the logarithms of 10, 100, 1000, ... to base 10.

372. The logarithm of N to base a is usually written $\log_a N$, so that the same meaning is expressed by the two equations

$$a^x = N; \quad x = \log_a N.$$

Example. Find the logarithm of $32\sqrt[4]{4}$ to base $2\sqrt{2}$.

Let x be the required logarithm; then

by definition,

$$(2\sqrt{2})^x = 32\sqrt[4]{4};$$

$$\therefore (2 \cdot 2^{\frac{1}{2}})^x = 2^5 \cdot 2^{\frac{1}{2}};$$

$$\therefore 2^{\frac{3}{2}x} = 2^{5\frac{1}{2}};$$

hence, by equating the indices, $\frac{3}{2}x = \frac{27}{5};$

$$\therefore x = \frac{18}{5} = 3.6.$$

373. When it is understood that a particular system of logarithms is in use, the suffix denoting the base is omitted. Thus in arithmetical calculations in which 10 is the base, we usually write $\log 2$, $\log 3$, ... instead of $\log_{10} 2$, $\log_{10} 3$, ...

Logarithms to the base 10 are known as **Common Logarithms**; this system was first introduced in 1615 by Briggs, a contemporary of Napier the inventor of Logarithms. Before discussing the properties of common logarithms we shall prove some general propositions which are true for all logarithms independently of any particular base.

374. *The logarithm of 1 is 0.*

For $a^0=1$ for all values of a ; therefore $\log 1=0$, whatever the base may be.

375. *The logarithm of the base itself is 1.*

For $a^1=a$; therefore $\log_a a=1$.

376. *To find the logarithm of a product.*

Let MN be the product; let a be the base of the system, and suppose

$$x=\log_a M, \quad y=\log_a N;$$

so that

$$a^x=M, \quad a^y=N.$$

Thus the product $MN=a^x \times a^y=a^{x+y}$;

whence, by definition, $\log_a MN=x+y$

$$=\log_a M + \log_a N.$$

Similarly,

$$\log_a MNP=\log_a M + \log_a N + \log_a P;$$

and so on for any number of factors.

Example. $\log 42=\log (2 \times 3 \times 7)=\log 2 + \log 3 + \log 7.$

377. *To find the logarithm of a fraction.*

Let $\frac{M}{N}$ be the fraction, and suppose

$$x=\log_a M, \quad y=\log_a N;$$

so that

$$a^x=M, \quad a^y=N.$$

Thus the fraction

$$\frac{M}{N}=\frac{a^x}{a^y}=a^{x-y};$$

whence, by definition, $\log_a \frac{M}{N}=x-y$

$$=\log_a M - \log_a N.$$

Example. $\log(2\frac{1}{7}) = \log \frac{15}{7} = \log 15 - \log 7$
 $= \log(3 \times 5) - \log 7 = \log 3 + \log 5 - \log 7.$

378. To find the logarithm of a number raised to any power, integral or fractional.

Let $\log_a(M^p)$ be required, and suppose

$$x = \log_a M, \text{ so that } a^x = M;$$

then

$$M^p = (a^x)^p = a^{px};$$

whence, by definition, $\log_a(M^p) = px;$

that is,

$$\log_a(M^p) = p \log_a M.$$

Similarly,

$$\log_a(M^{\frac{1}{r}}) = \frac{1}{r} \log_a M.$$

Example. Express the logarithm of $\frac{\sqrt{a^3}}{c^5 b^2}$ in terms of $\log a$, $\log b$, and $\log c$.

$$\begin{aligned} \log \frac{\sqrt{a^3}}{c^5 b^2} &= \log \frac{a^{\frac{3}{2}}}{c^5 b^2} = \log a^{\frac{3}{2}} - \log(c^5 b^2) \\ &= \frac{3}{2} \log a - (\log c^5 + \log b^2) = \frac{3}{2} \log a - 5 \log c - 2 \log b. \end{aligned}$$

379. From the equation $10^x = N$, it is evident that common logarithms will not in general be integral, and that they will not always be positive.

For instance, $3154 > 10^3$ and $< 10^4$;

$$\therefore \log 3154 = 3 + \text{a fraction.}$$

Again,

$$06 > 10^{-2} \text{ and } < 10^{-1};$$

$$\therefore \log 06 = -2 + \text{a fraction.}$$

380. DEFINITION. The integral part of a logarithm is called the **characteristic**, and the decimal part is called the **mantissa**.

The characteristic of the logarithm of any number to the base 10 can be written down by inspection, as we shall now shew.

381. *To determine the characteristic of the logarithm of any number greater than unity.*

It is clear that a number with two digits in its integral part lies between 10^1 and 10^2 ; a number with three digits in its integral part lies between 10^2 and 10^3 ; and so on. Hence a number with n digits in its integral part lies between 10^{n-1} and 10^n .

Let N be a number whose integral part contains n digits; then

$$N = 10^{(n-1)} + \text{a fraction};$$

$$\therefore \log N = (n-1) + \text{a fraction}.$$

Hence the characteristic is $n-1$; that is, *the characteristic of the logarithm of a number greater than unity is less by one than the number of digits in its integral part, and is positive.*

382. *To determine the characteristic of the logarithm of a decimal fraction.*

A decimal with one cipher immediately after the decimal point, such as .0324, being greater than .01 and less than .1, lies between 10^{-2} and 10^{-1} ; a number with two ciphers after the decimal point lies between 10^{-3} and 10^{-2} ; and so on. Hence a decimal fraction with n ciphers immediately after the decimal point lies between $10^{-(n+1)}$ and 10^{-n} .

Let D be a decimal beginning with n ciphers; then

$$D = 10^{-(n+1)} + \text{a fraction};$$

$$\therefore \log D = -(n+1) + \text{a fraction}.$$

Hence the characteristic is $-(n+1)$; that is, *the characteristic of the logarithm of a decimal fraction is greater by unity than the number of ciphers immediately after the decimal point and is negative.*

383. The logarithms to base 10 of all integers from 1 to 200000 have been found and tabulated; in most Tables they are given to seven places of decimals. This is the system in practical use, and it has two great advantages:

(1) From the results already proved it is evident that the characteristics can be written down by inspection, so that only the mantissæ have to be registered in the Tables.

(2) The mantissæ are the same for the logarithms of all

numbers which have the same significant digits; so that it is sufficient to tabulate the mantissæ of the logarithms of *integers*.

This proposition we proceed to prove.

384. Let N be any number, then since multiplying or dividing by a power of 10 merely alters the position of the decimal point without changing the sequence of figures, it follows that $N \times 10^p$, and $N \div 10^q$, where p and q are any integers, are numbers whose significant digits are the same as those of N .

$$\begin{aligned} \text{Now } \log(N \times 10^p) &= \log N + p \log 10 \\ &= \log N + p \dots\dots\dots(1). \end{aligned}$$

$$\begin{aligned} \text{Again, } \log(N \div 10^q) &= \log N - q \log 10 \\ &= \log N - q \dots\dots\dots(2). \end{aligned}$$

In (1) an integer is added to $\log N$, and in (2) an integer is subtracted from $\log N$; that is, the mantissa or decimal portion of the logarithm remains unaltered.

In this and the three preceding articles the mantissæ have been supposed positive. In order to secure the advantages of Briggs' system, we arrange our work so as *always* to keep the mantissa positive, so that when the mantissa of any logarithm has been taken from the Tables the characteristic is prefixed with its appropriate sign, according to the rules already given.

385. In the case of a negative logarithm the minus sign is written *over the characteristic*, and not before it, to indicate that the characteristic alone is negative, and not the whole expression. Thus $\bar{4}30103$, the logarithm of '0002, is equivalent to $-4 + '30103$, and must be distinguished from -430103 , an expression in which both the integer and the decimal are negative. In working with negative logarithms an arithmetical artifice will sometimes be necessary in order to make the mantissa positive. For instance, a result such as -3.69897 , in which the whole expression is negative, may be transformed by subtracting 1 from the characteristic and adding 1 to the mantissa. Thus

$$-3.69897 = -4 + (1 - .69897) = \bar{4}30103.$$

Example 1. Required the logarithm of '0002432.

In the Tables we find that 3859636 is the mantissa of $\log 2432$ (the decimal point as well as the characteristic being omitted); and, by Art. 382, the characteristic of the logarithm of the given number is -4 ;

$$\therefore \log '0002432 = \bar{4}3859636.$$

its; so that it is
rithms of *integers*.

e multiplying or
e position of the
figures, it follows
any integers, are
as those of N .

.....(1).

.....(2).

(2) an integer is
e decimal portion

e mantissæ have
the advantages of
ways to keep the
f any logarithm
ristic is prefixed
already given.

he minus sign is
to indicate that
whole expression.
t to $-4 + \cdot 30103$,
n expression in
ative. In work-
tifice will some-
a positive. For
e whole expres-
ting 1 from the
us

03.

tissa of $\log 2432$
(g omitted); and,
he given number

Example 2. Find the value of $\sqrt[5]{\cdot 00000165}$, given

$$\log 165 = 2 \cdot 2174839, \quad \log 697424 = 5 \cdot 8434968.$$

Let x denote the value required; then

$$\begin{aligned} \log x &= \log (\cdot 00000165)^{\frac{1}{5}} = \frac{1}{5} \log (\cdot 00000165) \\ &= \frac{1}{5} (\bar{6} \cdot 2174839); \end{aligned}$$

the *mantissa* of $\log \cdot 00000165$ being the same as that of $\log 165$, and the *characteristic* being prefixed by the rule.

$$\text{Now} \quad \frac{1}{5} (\bar{6} \cdot 2174839) = \frac{1}{5} (\bar{10} + 4 \cdot 2174839) = \bar{2} \cdot 8434968$$

and $\cdot 8434968$ is the mantissa of $\log 697424$; hence x is a number consisting of these same digits but with one cipher after the decimal point. [Art. 382.]

Thus

$$x = \cdot 0697424.$$

386. Suppose that the logarithms of all numbers to base a are known and tabulated, it is required to find the logarithms to base b .

Let N be any number whose logarithm to base b is required.

Let $y = \log_a N$, so that $b^y = N$;

$$\therefore \log_a (b^y) = \log_a N;$$

that is,

$$y \log_a b = \log_a N;$$

$$\therefore y = \frac{1}{\log_a b} \times \log_a N,$$

or

$$\log_b N = \frac{1}{\log_a b} \times \log_a N \dots\dots\dots(1).$$

Now since N and b are given, $\log_a N$ and $\log_a b$ are known from the Tables, and thus $\log_b N$ may be found.

Hence it appears that to transform logarithms from base a to base b we have only to multiply them all by $\frac{1}{\log_a b}$; this is a constant quantity and is given by the Tables; it is known as the *modulus*.

Cor. If in equation (1) we put a for N , we obtain

$$\log_b a = \frac{1}{\log_a b} \times \log_a a = \frac{1}{\log_a b};$$

$$\therefore \log_b a \times \log_a b = 1.$$

387. The following examples illustrate the utility of logarithms in facilitating arithmetical calculation.

Example 1. Given $\log 3 = .4771213$, find $\log \{(2.7)^3 \times (.81)^{\frac{1}{2}} \div (90)^{\frac{1}{2}}\}$.

$$\begin{aligned} \text{The required value} &= 3 \log \frac{27}{10} + \frac{4}{5} \log \frac{81}{100} - \frac{5}{4} \log 90 \\ &= 3(\log 3^3 - 1) + \frac{4}{5}(\log 3^4 - 2) - \frac{5}{4}(\log 3^2 + 1) \\ &= \left(9 + \frac{16}{5} - \frac{5}{2}\right) \log 3 - \left(3 + \frac{8}{5} + \frac{5}{4}\right) \\ &= \frac{97}{10} \log 3 - 5\frac{17}{20} \\ &= 4.6280766 - 5.85 \\ &= \bar{2}.7780766. \end{aligned}$$

The student should notice that the logarithm of 5 and its powers can always be obtained from $\log 2$; thus

$$\log 5 = \log \frac{10}{2} = \log 10 - \log 2 = 1 - \log 2.$$

Example 2. Find the number of digits in 875^{16} , given

$$\log 2 = .3010300, \quad \log 7 = .8450980.$$

$$\begin{aligned} \log(875^{16}) &= 16 \log(7 \times 125) \\ &= 16(\log 7 + 3 \log 5) \\ &= 16(\log 7 + 3 - 3 \log 2) \\ &= 16 \times 2.9420080 \\ &= 47.072128; \end{aligned}$$

hence the number of digits is 48. [Art. 381.]

Example 3. Given $\log 2$ and $\log 3$, find to two places of decimals the value of x from the equation

$$6^{3-4x} \cdot 4^{x+5} = 8.$$

Taking logarithms of both sides, we have

$$(3-4x) \log 6 + (x+5) \log 4 = \log 8;$$

$$\therefore (3-4x)(\log 2 + \log 3) + (x+5)2 \log 2 = 3 \log 2;$$

$$\therefore x(-4 \log 2 - 4 \log 3 + 2 \log 2) = 3 \log 2 - 3 \log 2 - 3 \log 3 - 10 \log 2;$$

$$\therefore x = \frac{10 \log 2 + 3 \log 3}{2 \log 2 + 4 \log 3}$$

$$= \frac{4.4416639}{2.5105452}$$

$$= 1.77 \dots$$

EXAMPLES XXXIX.

1. Find the logarithms of $\sqrt{32}$ and $\cdot 03125$ to base $\frac{3}{2}$, and 100 and $\cdot 00001$ to base $\cdot 01$.

2. Find the value of

$$\log_4 512, \log_5 \cdot 0016, \log_{\frac{1}{27}} \frac{1}{27}, \log_4 343.$$

3. Write down the numbers whose logarithms to bases

$$25, 3, \cdot 02, 1, -4, 1.7, 1000$$

are

$$\frac{1}{2}, -2, -3, 5, -1, 2, -\frac{2}{3} \text{ respectively.}$$

Simplify the expressions:

$$4. \log \frac{(ab^2c^4)^{\frac{1}{3}}}{\sqrt[3]{a^{-3}b^3c^6}}.$$

$$5. \log \left\{ \left(\frac{x^4y^{-3}}{x^{-1}y^3} \right)^{-3} \div \left(\frac{x^{-2}y^3}{xy^{-1}} \right)^5 \right\}.$$

6. Find by inspection the characteristics of the logarithms of 3174, 625.7, 3.502, .4, .374, .000135, 23.22065.

7. The mantissa of $\log 37203$ is $\cdot 5705780$: write down the logarithms of 37.203, $\cdot 000037203$, 372030000.

8. The logarithm of 7623 is 3.8821259 : write down the numbers whose logarithms are $\cdot 8821259$, 6.8821259 , 7.8821259 .

Given $\log 2 = \cdot 3010300$, $\log 3 = \cdot 4771213$, $\log 7 = \cdot 8450980$, find the value of

$$9. \log 729.$$

$$10. \log 8400.$$

$$1. \log \cdot 256.$$

$$12. \log 5.832.$$

$$13. \log \sqrt[3]{392}.$$

$$14. \log \cdot 3048.$$

15. Shew that $\log \frac{11}{15} + \log \frac{490}{297} - 2 \log \frac{7}{9} = \log 2$.
16. Find to six decimal places the value of $\log \frac{225}{224} - 2 \log \frac{20}{189} + \log \frac{512}{81}$.
17. Simplify $\log \{(10 \cdot 8)^{\frac{1}{2}} \times (24)^{\frac{1}{3}} \div (90)^{-2}\}$, and find its numerical value.
18. Find the value of $\log (\sqrt[2]{128} \cdot \sqrt{108} \div \sqrt[2]{1008} \cdot \sqrt[2]{162})$.
19. Find the value of $\log \sqrt{\frac{588 \times 768}{686 \times 972}}$.
20. Find the number of digits in 42^{42} .
21. Shew that $\left(\frac{81}{80}\right)^{1000}$ is greater than 100000.
22. How many ciphers are there between the decimal point and the first significant digit in $\left(\frac{2}{3}\right)^{1000}$?
23. Find the value of $\sqrt[2]{0.1008}$, having given $\log 398742 = 5.6006921$.
24. Find the seventh root of .00792, having given $\log 11 = 1.0413927$ and $\log 500.677 = 2.6998179$.
25. Find the value of $2 \log \frac{75}{49} + \log \frac{135}{32} - 3 \log \frac{45}{28}$.
- Find the numerical value of x in the following equations:
26. $3^{x+2} = 405$. 27. $10^{5-2x} = 27^{-2x}$.
28. $5^{x-3} = 8$. 29. $12^{2x-4} \cdot 187^{-2x} = 1458$.

CHAPTER XL.

SCALES OF NOTATION.

388. THE ordinary numbers with which we are acquainted in Arithmetic are expressed by means of multiples of powers of 10; for instance

$$25 = 2 \times 10 + 5;$$

$$4705 = 4 \times 10^3 + 7 \times 10^2 + 0 \times 10 + 5.$$

This method of representing numbers is called the **common** or **denary scale of notation**, and ten is said to be the **radix** of the scale. The symbols employed in this system of notation are the nine digits and zero.

In like manner any number other than ten may be taken as the radix of a scale of notation; thus if 7 is the radix, a number expressed by 2453 represents $2 \times 7^3 + 4 \times 7^2 + 5 \times 7 + 3$; and in this scale no digit higher than 6 can occur.

389. The names Binary, Ternary, Quaternary, Quinary, Senary, Septenary, Octenary, Nonary, Denary, Undenary, and Duodenary are used to denote the scales corresponding to the values *two, three, ... twelve* of the radix.

In the undenary, duodenary, ... scales we shall require symbols to represent the digits which are greater than nine. It is unusual to consider any scale higher than that with radix twelve; when necessary we shall employ the symbols *t, e, T* as digits to denote 'ten,' 'eleven' and 'twelve.'

It is especially worthy of notice that in every scale 10 is the symbol not for 'ten', but for the radix itself.

390. The ordinary operations of Arithmetic may be performed in any scale; but, bearing in mind that the successive powers of the radix are no longer powers of ten in determining the *carrying figures* we must not divide by ten, but by the radix of the scale in question.

Example 1. In the scale of eight subtract 371532 from 530225, and multiply the difference by 7.

$$\begin{array}{r} 530225 \\ 371532 \\ \hline 136473 \end{array}$$

$$\begin{array}{r} 136473 \\ 7 \\ \hline 1226235 \end{array}$$

Explanation. After the first figure of the subtraction, since we cannot take 3 from 2 we add 8; thus we have to take 3 from ten, which leaves 7; then 6 from ten, which leaves 4; then 2 from eight which leaves 6; and so on.

Again, in multiplying by 7, we have

$$3 \times 7 = \text{twenty-one} = 2 \times 8 + 5;$$

we therefore put down 5 and carry 2.

Next

$$7 \times 7 + 2 = \text{fifty-one} = 6 \times 8 + 3;$$

put down 3 and carry 6; and so on.

Example 2. Divide 15et20 by 9 in the scale of twelve.

$$\begin{array}{r} 9 \overline{) 15et20} \\ 1ee96 \dots 6. \end{array}$$

Explanation. Since $15 = 1 \times T + 5 = \text{seventeen} = 1 \times 9 + 8$, we put down 1 and carry 8.

Also $8 \times T + e = \text{one hundred and seven} = e \times 9 + 8$; we therefore put down e and carry 8; and so on.

391. To express a given integral number in any proposed scale.

Let N be the given number, and r the radix of the proposed scale.

Let $a_0, a_1, a_2, \dots, a_n$ be the required digits by which N is to be expressed, beginning with that in the unit's place; then

$$N = a_n r^n + a_{n-1} r^{n-1} + \dots + a_2 r^2 + a_1 r + a_0.$$

We have now to find the values of $a_0, a_1, a_2, \dots, a_n$.

Divide N by r , then the remainder is a_0 , and the quotient is

$$a_n r^{n-1} + a_{n-1} r^{n-2} + \dots + a_2 r + a_1.$$

If this quotient is divided by r , the remainder is a_1 ; if the next quotient..... a_2 ;

and so on, until there is no further quotient.

Thus all the required digits $a_0, a_1, a_2, \dots, a_n$ are determined by successive divisions by the radix of the proposed scale.

1532 from 530225,

36473
7
26235

subtraction, since we
take 3 from ten,
then 2 from eight

Example 1. Express the denary number 5213 in the scale of seven.

$$\begin{array}{r} 7 \overline{) 5213} \\ 7 \overline{) 744} \dots 5 \\ 7 \overline{) 106} \dots 2 \\ 7 \overline{) 15} \dots 1 \\ 2 \dots 1 \end{array}$$

Thus $5213 = 2 \times 7^4 + 1 \times 7^3 + 1 \times 7^2 + 2 \times 7 + 5$;
and the number required is 21125.

Example 2. Transform 21125 from scale seven to scale eleven.

$$\begin{array}{r} e \overline{) 21125} \\ e \overline{) 1244} \dots t \\ e \overline{) 61} \dots 0 \\ 3 \dots t \end{array}$$

$\therefore$ the required number is 3t0t.

Explanation. In the first line of work

$$21 = 2 \times 7 + 1 = \text{fifteen} = 1 \times e + 4;$$

therefore on dividing by e we put down 1 and carry 4.

$$\text{Next } 4 \times 7 + 1 = \text{twenty-nine} = 2 \times e + 7;$$

therefore we put down 2 and carry 7; and so on.

392. Hitherto we have only discussed whole numbers; but fractions may also be expressed in any scale of notation; thus

$$\cdot 25 \text{ in scale ten denotes } \frac{2}{10} + \frac{5}{10^2};$$

$$\cdot 25 \text{ in scale six denotes } \frac{2}{6} + \frac{5}{6^2};$$

$$\cdot 25 \text{ in scale } r \text{ denotes } \frac{2}{r} + \frac{5}{r^2};$$

Fractions thus expressed in a form analogous to that of ordinary decimal fractions are called **radix-fractions**, and the point is called the **radix-point**. The general type of such fractions in scale r is

$$\frac{b_1}{r} + \frac{b_2}{r^2} + \frac{b_3}{r^3} + \dots;$$

where $b_1, b_2, b_3, \dots$ are integers, all less than r , of which any one or more may be zero.

393. To express a given radix-fraction in any proposed scale.

Let F be the given fraction, and r the radix of the scale.

Let $b_1, b_2, b_3, \dots$ be the required digits beginning from the left; then

$$F = \frac{b_1}{r} + \frac{b_2}{r^2} + \frac{b_3}{r^3} + \dots$$

We have now to find the values of $b_1, b_2, b_3, \dots$

Multiply both sides of the equation by r ; then

$$rF = b_1 + \frac{b_2}{r} + \frac{b_3}{r^2} + \dots$$

Hence b_1 is equal to the integral part of rF ; and, if we denote the fractional part by F_1 , we have

$$F_1 = \frac{b_2}{r} + \frac{b_3}{r^2} + \dots$$

Multiply again by r ; then b_2 is the integral part of rF_1 . Similarly by successive multiplications by r , each of the digits may be found, and the fraction expressed in the proposed scale.

Example 1. Express $\frac{7}{8}$ as a radix-fraction in scale six.

$$\frac{7}{8} \times 6 = \frac{7 \times 3}{4} = 5 + \frac{1}{4};$$

$$\frac{1}{4} \times 6 = \frac{1 \times 3}{2} = 1 + \frac{1}{2};$$

$$\frac{1}{2} \times 6 = 3.$$

$$\therefore \text{the required fraction} = \frac{5}{6} + \frac{1}{6^2} + \frac{3}{6^3} = .513.$$

Example 2. Transform 1606·7 from scale eight to scale five.

Treating the integral and the fractional parts separately, we have

5) 1606	·7
5) 264 ... 2	5
5) 44 ... 0	4·3
5) 7 ... 1	5
1 ... 2	1·7

After this the digits in the fractional part recur; hence the required number is 12102·41.

y proposed scale.
of the scale.

inning from the

.....
en

rF ; and, if we

gral part of rF_1 .
 r , each of the
t in the proposed

le six.

to scale five.

parately, we have

$\begin{array}{r} 7 \\ 5 \\ \hline 3 \\ 5 \\ \hline 7 \end{array}$

recur; hence the

Example 3. In what scale is the septenary number 2403 represented by 735?

Let r be the radix of the scale required; then

$$7r^2 + 3r + 5 = 2 \times 7^3 + 4 \times 7^2 + 3 = 885;$$

that is, $7r^2 + 3r - 880 = 0$; whence $r = 11$ or $-\frac{80}{7}$.

Thus the scale is the undenary.

394. In any scale of notation of which the radix is r , the sum of the digits of any whole number divided by $r-1$ will leave the same remainder as the whole number divided by $r-1$.

Let N denote the number, $a_0, a_1, a_2, \dots, a_n$ the digits beginning with that in the unit's place, and S the sum of the digits; then

$$N = a_0 + a_1r + a_2r^2 + \dots + a_{n-1}r^{n-1} + a_nr^n;$$

$$S = a_0 + a_1 + a_2 + \dots + a_{n-1} + a_n.$$

$$\therefore N - S = a_1(r-1) + a_2(r^2-1) + \dots + a_{n-1}(r^{n-1}-1) + a_n(r^n-1).$$

Now every term on the right-hand side is divisible by $r-1$;

$$\therefore \frac{N-S}{r-1} = \text{an integer} = I \text{ suppose;}$$

that is, $\frac{N}{r-1} = I + \frac{S}{r-1}$; which proves the proposition.

Hence a number in scale r will be divisible by $r-1$ when the sum of its digits is divisible by $r-1$. For example, in the ordinary scale a number is divisible by 9 when the sum of its digits is divisible by 9.

EXAMPLES XL.

1. Add together 352, 21435, 3505, 35 in the scale of six.
2. From 35260013 take 7471235 in the scale of eight.
3. Multiply 31044 by 4302 in the quinary scale,
4. Find the product of the undenary numbers 9483 and 347.

E.A.

Z

5. Divide 31664435 by 6541 in the scale of seven.
6. Find the square of 3024 in the quinary scale.
7. Express 75013 in the nonary, and 5210 in the quaternary scale.
8. Transform 987504 to the scale of twelve.
9. Express the octenary number 76543 in the denary scale.
10. Transform 54321 from scale six to scale seven.
11. Express the duodenary number *te* in the binary scale.
12. Express a thousand and one in powers of two, and one hundred thousand in powers of eleven.
13. Express the sum of the septenary numbers 532, 2106, 3261, 53 in the undenary scale; also express the difference of the ternary numbers 2021121 and 1221212 in the same scale, and find the product of the two results.
14. Find the difference between 53774 in the scale of 8 and 32875 in the scale of 9, expressing the result in the denary scale.
15. Express 131·890625 in scale eight.
16. Transform 1001·12211 from the ternary to the nonary scale.
17. Express the octenary fraction ·2037 in the scale of 4.
18. Express $\frac{27}{32}$ and $\frac{500}{729}$ as radix fractions in scale 6.
19. Reduce the undenary fraction $\frac{587}{749}$ to its lowest terms.
20. In what scale is a hundred denoted by 400?
21. In what scale is 647 the square of 25?
22. In what scale are the numbers denoted by 432, 565, 708 in arithmetical progression?
23. In what scale are the numbers denoted by ·22, 2·6, 34 in geometrical progression?
24. Find the square root of 443001 in the scale of 5; 2434524 in the scale of 7; and 4985679 in the scale of eleven.

even.

scale.

in the quaternary

e denary scale.

even.

binary scale.

s of two, and one

ers 532, 2106, 3261,

ence of the ternary

and find the product

scale of 8 and 32875

ary scale.

ary to the nonary

e scale of 4.

scale 6.

lowest terms.

00?

by 432, 565, 708 in

l by .22, 2.6, 34 in

le of 5; 2434524 in

.

CHAPTER XLI.

EXPONENTIAL AND LOGARITHMIC SERIES.

395. THE advantages of common logarithms have been explained in Art. 383, and in practice no other system is used. But in the first place these logarithms are calculated to another base and then transformed to the base 10.

In the present chapter we shall prove certain formulæ known as the **Exponential and Logarithmic Series**, and give a brief explanation of the way in which they are used in constructing a table of logarithms.

396. *To expand a^x in ascending powers of x .*

By the Binomial Theorem, if $n > 1$,

$$\begin{aligned} \left(1 + \frac{1}{n}\right)^{nx} &= 1 + nx \cdot \frac{1}{n} + \frac{nx(nx-1)}{2} \cdot \frac{1}{n^2} + \frac{nx(nx-1)(nx-2)}{3} \cdot \frac{1}{n^3} + \dots \\ &= 1 + x + \frac{x\left(x - \frac{1}{n}\right)}{2} + \frac{x\left(x - \frac{1}{n}\right)\left(x - \frac{2}{n}\right)}{3} + \dots \dots \dots (1). \end{aligned}$$

By putting $x=1$, we obtain

$$\left(1 + \frac{1}{n}\right)^n = 1 + 1 + \frac{1 - \frac{1}{n}}{2} + \frac{\left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)}{3} + \dots \dots (2).$$

But

$$\left(1 + \frac{1}{n}\right)^{nx} = \left\{ \left(1 + \frac{1}{n}\right)^n \right\}^x;$$

hence the series (1) is the x^{th} power of the series (2); that is,

$$\begin{aligned} &+ x + \frac{x\left(x - \frac{1}{n}\right)}{2} + \frac{x\left(x - \frac{1}{n}\right)\left(x - \frac{2}{n}\right)}{3} + \dots \\ &= \left\{ 1 + 1 + \frac{1 - \frac{1}{n}}{2} + \frac{\left(1 - \frac{1}{n}\right)\left(1 - \frac{2}{n}\right)}{3} + \dots \right\}^x; \end{aligned}$$

and this is true however great n may be. If therefore n be indefinitely increased we have

$$1+x+\frac{x^2}{2}+\frac{x^3}{3}+\dots=\left(1+1+\frac{1}{2}+\frac{1}{3}+\dots\right)^x.$$

The series $1+1+\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+\dots$

is usually denoted by e ; hence

$$e^x=1+x+\frac{x^2}{2}+\frac{x^3}{3}+\frac{x^4}{4}+\dots$$

Write cx for x , then

$$e^{cx}=1+cx+\frac{c^2x^2}{2}+\frac{c^3x^3}{3}+\dots$$

Now let $e^c=a$, so that $c=\log_e a$; by substituting for c we obtain

$$a^x=1+x\log_e a+\frac{x^2(\log_e a)^2}{2}+\frac{x^3(\log_e a)^3}{3}+\dots$$

This is the *Exponential Theorem*.

397. The series

$$1+1+\frac{1}{2}+\frac{1}{3}+\frac{1}{4}+\dots,$$

which we have denoted by e , is very important as it is the base to which logarithms are first calculated. Logarithms to this base are known as the Napierian system, so named after Napier their inventor. They are also called *natural* logarithms from the fact that they are the first logarithms which naturally come into consideration in algebraical investigations.

When logarithms are used in theoretical work it is to be remembered that the base e is always understood, just as in arithmetical work the base 10 is invariably employed.

From the series the approximate value of e can be determined to any required degree of accuracy; to 10 places of decimals it is found to be 2.7182818284.

If therefore n be

$$\left(\frac{1}{3} + \dots\right)^n.$$

Example 1. Find the sum of the infinite series

$$1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots$$

We have

$$e = 1 + 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots;$$

and by putting $x = -1$ in the series for e^x , we obtain

$$e^{-1} = 1 - 1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} - \dots$$

$$\therefore e + e^{-1} = 2 \left(1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots \right);$$

hence the sum of the series is $\frac{1}{2}(e + e^{-1})$.

Example 2. Find the coefficient of x^r in the expansion of $\frac{a-bx}{e^x}$.

$$\frac{a-bx}{e^x} = (a-bx)e^{-x}$$

$$= (a-bx) \left\{ 1 - x + \frac{x^2}{2} - \frac{x^3}{3} + \dots + \frac{(-1)^r x^r}{r} + \dots \right\}.$$

$$\text{The coefficient required} = \frac{(-1)^r}{r} \cdot a - \frac{(-1)^{r-1}}{r-1} \cdot b$$

$$= \frac{(-1)^r}{r} (a + rb).$$

398. To expand $\log_e(1+x)$ in ascending powers of x .

From Art. 396,

$$a^y = 1 + y \log_e a + \frac{y^2 (\log_e a)^2}{2} + \frac{y^3 (\log_e a)^3}{3} + \dots$$

In this series write $1+x$ for a ; thus $(1+x)^y$

$$= 1 + y \log_e(1+x) + \frac{y^2}{2} \{\log_e(1+x)\}^2 + \frac{y^3}{3} \{\log_e(1+x)\}^3 + \dots (1).$$

Also by the Binomial Theorem, when $y < 1$ we have

$$(1+x)^y = 1 + yx + \frac{y(y-1)}{2} x^2 + \frac{y(y-1)(y-2)}{3} x^3 + \dots (2).$$

Now in (2) the coefficient of y is

$$x + \frac{(-1)}{1.2}x^2 + \frac{(-1)(-2)}{1.2.3}x^3 + \frac{(-1)(-2)(-3)}{1.2.3.4}x^4 + \dots;$$

that is,
$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

Equate this to the coefficient of y in (1); thus we have

$$\log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

This is known as the *Logarithmic Series*.

399. Except when x is very small the series for $\log_e(1+x)$ is of little use for numerical calculations. We can, however, deduce from it other series by the aid of which Tables of Logarithms may be constructed.

400. In Art. 398 we have proved that

$$\log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots;$$

replacing x by $-x$, we have

$$\log_e(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \dots$$

By subtraction,

$$\log_e \frac{1+x}{1-x} = 2 \left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \right).$$

Put $\frac{1+x}{1-x} = \frac{n+1}{n}$, so that $x = \frac{1}{2n+1}$; we thus obtain

$$\log_e(n+1) - \log_e n = 2 \left\{ \frac{1}{2n+1} + \frac{1}{3(2n+1)^3} + \frac{1}{5(2n+1)^5} + \dots \right\}.$$

From this formula by putting $n=1$ we can obtain $\log_e 2$. Again, by putting $n=2$ we obtain $\log_e 3 - \log_e 2$; whence $\log_e 3$ is found, and therefore also $\log_e 9$ is known.

Now by putting $n=9$ we obtain $\log_e 10 - \log_e 9$; thus the value of $\log_e 10$ is found to be 2.30258509...

To convert Napierian logarithms into logarithms to base 10 we multiply by $\frac{1}{\log_e 10}$, which is the *modulus* [Art. 386] of the

common system, and its value is $\frac{1}{2.30258509...}$, or .43429448...; we shall denote this modulus by μ .

By multiplying the last series throughout by μ we obtain a formula adapted to the calculation of common logarithms. Thus

$$\mu \log_e(n+1) - \mu \log_e n = 2\mu \left\{ \frac{1}{2n+1} + \frac{1}{3(2n+1)^3} + \frac{1}{5(2n+1)^5} + \dots \right\};$$

that is,

$$\log_{10}(n+1) - \log_{10} n = 2 \left\{ \frac{\mu}{2n+1} + \frac{\mu}{3(2n+1)^3} + \frac{\mu}{5(2n+1)^5} + \dots \right\}.$$

From this result we see that if the logarithm of one of two consecutive numbers be known, the logarithm of the other may be found, and thus a table of logarithms can be constructed.

EXAMPLES XII.

1. Shew that

$$(1) \quad e^{-2} = 1 - \frac{2^2}{2!} + \frac{2^4}{4!} - \frac{2^6}{6!} + \dots$$

$$(2) \quad \frac{e^2 - 1}{2e} = 1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \dots$$

2. Expand $\log \sqrt{1+x}$ in ascending powers of x .

3. Prove that $\log_e 2 = \frac{1}{2} + \frac{1}{12} + \frac{1}{30} + \frac{1}{56} + \dots$

4. Shew that

$$\log_{10} \left(\frac{1}{1-x} \right) = \frac{1}{\log_e 10} \left(x + \frac{x^2}{2} + \frac{x^3}{3} + \dots \right).$$

5. Prove that

$$\log \frac{1+x}{1-3x} = 4x + 4x^2 + \frac{28}{3}x^3 + 20x^4 + \dots$$

6. Shew that if $x > 1$,

$$\log \sqrt{x^2 - 1} = \log x - \frac{1}{2x^2} - \frac{1}{4x^4} - \frac{1}{6x^6} - \dots$$

7. Shew that

$$\log \sqrt{\frac{1+x}{1-x}} - \log \sqrt{\frac{1-x}{1+x}} = 2 \left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \right).$$

8. If
$$a = b - \frac{b^2}{2} + \frac{b^3}{3} - \frac{b^4}{4} + \dots,$$

express b in ascending powers of a .

9. Calculate the value of $\sqrt{e}$ to 4 places of decimals.

10. Prove that

$$\log_e(1+x-2x^2) = x - \frac{5x^2}{2} + \frac{7x^3}{3} - \frac{17x^4}{4} + \dots,$$

and find the general term of the series.

11. Prove that
$$e^{-1} = 2 \left(\frac{1}{3} + \frac{2}{5} + \frac{3}{7} + \dots \right).$$

12. Prove that
$$\log_e 3 = 1 + \frac{1}{3 \cdot 2^2} + \frac{1}{5 \cdot 2^4} + \frac{1}{7 \cdot 2^6} + \dots$$

13. Shew that

$$\log_e(1-3x+2x^2)^{-1} = 3x + \frac{5x^2}{2} + 3x^3 + \frac{17x^4}{4} + \dots,$$

and find the general term of the series.

14. Prove that the expansion of $\log_e(1-x+x^2)$ is

$$-x + \frac{x^2}{2} + \frac{2x^3}{3} + \frac{x^4}{4} - \frac{x^5}{5} - \frac{x^6}{3} - \dots$$

15. If $x > 1$, prove that

$$\frac{1}{x} + \frac{1}{2x^2} + \frac{1}{3x^3} + \dots = \frac{1}{x-1} - \frac{1}{2(x-1)^2} + \frac{1}{3(x-1)^3} - \dots$$

CHAPTER XLII.

MISCELLANEOUS EQUATIONS.

401. MANY kinds of miscellaneous equations may be solved by the ordinary rules for quadratic equations as explained in Art. 202; but others require some special artifice for their solution. These will be illustrated in the present chapter.

Example 1. Solve $\frac{x^2-6}{x} + \frac{5x}{x^2-6} = 6$.

Write y for $\frac{x^2-6}{x}$; thus

$$y + \frac{5}{y} = 6, \text{ or } y^2 - 6y + 5 = 0;$$

whence

$$y = 5, \text{ or } 1.$$

$$\therefore \frac{x^2-6}{x} = 5, \text{ or } \frac{x^2-6}{x} = 1;$$

that is,

$$x^2 - 5x - 6 = 0, \text{ or } x^2 - x - 6 = 0.$$

Thus

$$x = 6, -1; \text{ or } x = 3, -2.$$

Example 2. Solve $3^{2x+3} - 55 = 28(3^x - 2)$.

This equation may be written $3^3 \cdot 3^{2x} - 28 \cdot 3^x + 1 = 0$.

By writing y for 3^x , we obtain

$$27y^2 - 28y + 1 = 0; \text{ that is, } (27y - 1)(y - 1) = 0$$

whence

$$y = \frac{1}{27}, \text{ or } 1.$$

Thus

$$3^x = \frac{1}{27} = 3^{-3}, \text{ or } 3^x = 1 = 3^0.$$

and therefore

$$x = -3, \text{ or } 0.$$

Example 3. Solve $2x^2 - 3\sqrt{2x^2 - 7x + 7} = 7x - 3$.

On transposition, $(2x^2 - 7x) - 3\sqrt{2x^2 - 7x + 7} = -3$.

By putting $\sqrt{2x^2 - 7x + 7} = y$, so that $2x^2 - 7x + 7 = y^2$, we obtain

$$(y^2 - 7) - 3y = -3, \text{ or } y^2 - 3y - 4 = 0;$$

whence

$$y = 4, \text{ or } -1.$$

Thus

$$\sqrt{2x^2 - 7x + 7} = 4, \text{ or } \sqrt{2x^2 - 7x + 7} = -1;$$

that is,

$$2x^2 - 7x - 9 = 0, \text{ or } 2x^2 - 7x + 6 = 0.$$

From the first of these quadratics we obtain $x = \frac{9}{2}$, or -1 , and from the second $x = 2$, or $\frac{3}{2}$.

It should be noticed that in this solution we have tacitly assumed y to be the *positive* value of the expression $\sqrt{2x^2 - 7x + 7}$, so that the roots obtained from the solution of $\sqrt{2x^2 - 7x + 7} = -1$ will only satisfy the original equation in the modified form obtained by changing the sign of the radical.

Thus $x = \frac{9}{2}$, or -1 satisfies $2x^2 - 3\sqrt{2x^2 - 7x + 7} = 7x - 3$.

and $x = 2$, or $\frac{3}{2}$ satisfies $2x^2 + 3\sqrt{2x^2 - 7x + 7} = 7x - 3$.

EXAMPLES XLII. a.

Solve the following equations :

- $x^2 + x + 1 = \frac{42}{x^2 + x}$.
- $\frac{x}{x^2 - 1} + \frac{x^2 - 1}{x} = 2\frac{1}{2}$.
- $\left(x + \frac{1}{x}\right)^2 - 4\left(x + \frac{1}{x}\right) = 5$.
- $8x^6 + 65x^3 + 8 = 0$.
- $\frac{x+8}{x+12} + \frac{5}{x+4} = \frac{3x+14}{3x+8}$.
- $4^x + 8 = 9 \cdot 2^x$.
- $\frac{3x-6}{5-x} + \frac{11-2x}{10-4x} = 3\frac{1}{2}$.
- $3\sqrt{x-3} \cdot x^{-\frac{1}{2}} = 8$.
- $\left(x - \frac{6}{x}\right)^2 + 4x - \frac{24}{x} = 5$.
- $27x^{\frac{2}{3}} - 1 = 26x^{\frac{1}{3}}$.
- $7\sqrt{x-8} - \sqrt{21x+12} = 2\sqrt{3}$.
- $4^{2x+1} + 16 = 65 \cdot 4^x$.
- $x+2 = \sqrt{4+x}\sqrt{8-x}$.
- $3^{\frac{x}{2}} + 3^{-\frac{x}{2}} = 2$.

$$15. \quad 2x^3 - 2x + 2\sqrt{2x^2 - 7x + 6} = 5x - 6.$$

$$16. \quad x^2 + 6\sqrt{x^2 - 2x + 5} = 11 + 2x.$$

$$17. \quad 2\sqrt{x^2 - 6x + 2} + 4x + 1 = x^2 - 2x.$$

$$18. \quad \sqrt{4x^2 + 2x + 7} = 12x^2 + 6x - 11.$$

$$19. \quad 3x(3-x) = 11 - 4\sqrt{x^2 - 3x + 5}.$$

$$20. \quad x^2 - x + 3\sqrt{2x^2 - 3x + 2} = \frac{x}{2} + 7.$$

$$21. \quad \sqrt{\frac{2-x}{3x}} - \sqrt{\frac{3x}{2-x}} = \frac{3}{2}.$$

$$22. \quad \sqrt{\frac{a}{x}} - \sqrt{\frac{x}{a}} = \frac{a^2 - 1}{a}.$$

$$23. \quad (a-b)x^2 + (b-c)x + c - a = 0.$$

$$24. \quad a(b-c)x^2 + b(c-a)x + c(a-b) = 0.$$

$$25. \quad \sqrt{a-x} + \sqrt{b-x} = \sqrt{a+b-2x}.$$

$$26. \quad \frac{1}{a-x} + \frac{1}{b-x} = \frac{1}{a-c} + \frac{1}{b}.$$

$$27. \quad \sqrt{x-p} + \sqrt{x-q} = \frac{p}{\sqrt{x-p}} + \frac{q}{\sqrt{x-p}}.$$

$$28. \quad \sqrt{(x-2)(x-3)} + 5\sqrt{\frac{x-2}{x-3}} = \sqrt{x^2 + 6x + 8}.$$

$$29. \quad \sqrt{x^2 + 4x - 4} + \sqrt{x^2 + 4x - 10} = 6.$$

$$30. \quad \sqrt[3]{x-a} - \sqrt[3]{x-b} = \sqrt[3]{b-a}.$$

402. No general methods can be given for the solution of simultaneous equations containing two or more unknowns. The simpler cases have been considered in Chapter XXVI.; the following examples illustrate useful artifices to be employed in special cases.

Example 1. Solve $x + y = 4$ (1),

$$(x^2 + y^2)(x^3 + y^3) = 280 \text{(2).}$$

We have $x^2 + y^2 = (x + y)^2 - 2xy = 16 - 2xy$;

and $x^3 + y^3 = (x + y)^3 - 3xy(x + y) = 64 - 12xy$.


By substituting in (2), we obtain

$$(16 - 2xy)(64 - 12xy) = 280; \text{ that is, } 3x^2y^2 - 40xy + 93 = 0;$$

whence


$$xy = 3, \text{ or } \frac{31}{3}.$$





MICROCOFY RESOLUTION TEST CHART

(ANSI and ISO TEST CHART No. 2)



APPLIED IMAGE Inc

1653 East Main Street
Rochester, New York 14609 USA
(716) 482-0300 - Phone
(716) 288-5989 - Fax

$$\begin{aligned} \text{Thus } \left. \begin{array}{l} x+y=4, \\ xy=3; \end{array} \right\} & \text{whence we obtain } \left. \begin{array}{l} x=3, \text{ or } 1; \\ y=1, \text{ or } 3; \end{array} \right\} \\ \text{or } \left. \begin{array}{l} x+y=4, \\ xy=\frac{31}{3}; \end{array} \right\} & \text{whence } \left. \begin{array}{l} x=2 \pm \sqrt{-\frac{19}{3}}; \\ y=2 \mp \sqrt{-\frac{19}{3}}. \end{array} \right\} \end{aligned}$$

Example 2. Solve $x^2y^2z=225$, $xy^2z^2=75$, $x^2yz^2=45$.

By multiplying the three equations together, we have

$$x^5y^5z^5=225 \times 75 \times 45=5^5 \times 3^5;$$

whence

$$xyz=5 \times 3=15.$$

By squaring this equation and dividing by each of the given equations in succession, we obtain

$$z=1, \quad x=3, \quad y=5.$$

Example 3. Solve the equations

$$x^2+xy+xz=48, \quad xy+y^2+yz=12, \quad xz+yz+z^2=84.$$

These equations may be written

$$x(x+y+z)=48, \quad y(x+y+z)=12, \quad z(x+y+z)=84.$$

By addition,

$$(x+y+z)(x+y+z)=144;$$

whence

$$x+y+z=\pm 12.$$

On dividing each of the given equations in turn by this last equation, we obtain

$$x=\pm 4, \quad y=\pm 1, \quad z=\pm 7.$$

It is clear that the roots must be taken either all positively or all negatively.

Example 4. Solve $x+y-z=14$ (1),

$$y^2+z^2-x^2=46$$
(2),

$$yz=9$$
(3).

From (2) and (3), $(y-z)^2-x^2=28$.

Put u for $y-z$; then this equation becomes

$$u^2-x^2=28.$$

Also from (1),

$$u+x=14;$$

by division,

$$u-x=2;$$

whence

$$x=6, \text{ and } u=8.$$

Thus $y-z=8$, and $yz=9$; whence $y=9$, or -1 ; $z=1$, or -9 ; and the solution is $x=6$, $y=9$, $z=1$; or $x=6$, $y=-1$, $z=-9$.

EXAMPLES XLII. b.

$$\begin{array}{lll} 1. & 3x - 2y = 11, & 2. & x^3 + y^3 = 91, & 3. & x^3 - y^3 = 335, \\ & 9x^2 - 4y^2 = 209. & & x^2y + xy^2 = 84. & & x^2y - xy^2 = 70. \end{array}$$

$$\begin{array}{ll} 4. & x^2 + xy + y^2 = 84, \\ & x + \sqrt{xy} + y = 14. \end{array} \quad \begin{array}{l} 5. & x^2 + xy + y^2 = 189, \\ & x - \sqrt{xy} + y = 9. \end{array}$$

$$\begin{array}{ll} 6. & \frac{3}{x^2} - \frac{1}{xy} - \frac{2}{y^2} = \frac{2}{9}, \\ & \frac{3}{x} + \frac{2}{y} = \frac{4}{3}. \end{array} \quad \begin{array}{l} 7. & \frac{2}{x^2} - \frac{3}{xy} - \frac{2}{y^2} = 17, \\ & \frac{1}{x} - \frac{2}{y} = 1. \end{array}$$

$$\begin{array}{ll} 8. & x^2y + y^2x = 20, \\ & \frac{1}{x} + \frac{1}{y} = \frac{5}{4}. \end{array} \quad \begin{array}{l} 9. & x^2 - 7xy + 4y^2 = 34, \\ & \frac{2x+y}{x-3y} - \frac{x-3y}{2x+y} = 2\frac{2}{3}. \end{array}$$

$$\begin{array}{ll} 10. & x^2 - xy + x = 35, \\ & xy - y^2 + y = 15. \end{array} \quad \begin{array}{l} 11. & (x+y)^2 + 3(x-y) = 30, \\ & xy + 3(x-y) = 11. \end{array}$$

$$\begin{array}{ll} 12. & (x-y)^2 = 3 - 2x - 2y, \\ & y(x-y+1) = x(y-x+1). \end{array} \quad \begin{array}{l} 13. & x^2 + 1 = 81(y^2 + y) \\ & x^2 + x = 9(y^3 + 1). \end{array}$$

14. Find the rational roots of

$$\begin{array}{ll} (1) & \left. \begin{array}{l} x+y=5 \\ (x^2+y^2)(x^3+y^3)=455 \end{array} \right\} \\ (2) & \left. \begin{array}{l} x-y=2 \\ (x^2+y^2)(x^3-y^3)=260 \end{array} \right\}. \end{array}$$

$$\begin{array}{ll} 15. & \sqrt{\frac{x}{y}} + \sqrt{\frac{y}{x}} = 4\frac{1}{4}, \\ & \frac{x}{\sqrt{y}} + \frac{y}{\sqrt{x}} = 16\frac{1}{4}. \end{array} \quad \begin{array}{l} 16. & \sqrt{\frac{x+y}{x-y}} + \sqrt{\frac{x-y}{x+y}} = \frac{34}{15}, \\ & \frac{x+y}{\sqrt{x-y}} + \frac{x-y}{\sqrt{x+y}} = \frac{152}{15}. \end{array}$$

$$17. \quad x^2yz = 72, \quad xy^2z = 48, \quad xyz^2 = 96.$$

$$18. \quad xyz = 30, \quad xyu = 120, \quad xzu = 20, \quad yzu = 24.$$

$$19. \quad yz + zx = 13, \quad zx + xy = 25, \quad xy + yz = 20.$$

$$20. \quad y(x+z)=112, \quad z(x+y)=132, \quad x(y+z)=90.$$

$$21. \quad (x+a)(y-b)=2, \quad (y-b)(z+c)=3, \quad (z+c)(x+a)=6.$$

$$22. \quad (x+y)(x+z)=63, \quad (y+z)(y+x)=42, \quad (z+x)(z+y)=54.$$

$$23. \quad x^2-xy-xz=14, \quad xy-y^2-yz=6, \quad xz-yz-z^2=4.$$

$$24. \quad x(3z-2y)=42, \quad y(x-2z)=4, \quad z(x+5y)+34=0.$$

$$25. \quad xy+x+y=11, \quad yz+y+z=3, \quad zx+z+x=2.$$

$$26. \quad (x+y)^2-z^2=65, \quad x^2-(y+z)^2=13, \quad x+y-z=5.$$

$$27. \quad z+x=9xyz, \quad x+y=5xyz, \quad y+z=8xyz.$$

$$28. \quad x^2+y^2+z^2=84, \quad x+y+z=14, \quad xy=z^2.$$

$$29. \quad x+y-z=1, \quad x^2-y^2+z^2=15, \quad xz=12.$$

$$30. \quad y+z-x=9, \quad x^2-y^2-z^2=15, \quad yz=3.$$

$$31. \quad x^2+y^2+z^2=133, \quad y+z-x=7, \quad yz=x^2.$$

$$32. \quad 3^x=9^{y-1}, \quad 16^{3-x}=8^{y-2}.$$

$$33. \quad 2^{y-1}=16^{x-2}, \quad 3^{\frac{1}{x}}=9^{\frac{1}{z}}, \quad \sqrt[3]{2^{y-3}}=\sqrt[2]{8^{x-2}}.$$

$$34. \quad x^2-(y-z)^2=a^2, \quad y^2-(z-x)^2=b^2, \quad z^2-(x-y)^2=c^2.$$

CHAPTER XLIII.

INTEREST AND ANNUITIES.

403. QUESTIONS involving Simple Interest are easily solved by the ordinary rules of Arithmetic; but in Compound Interest the calculations are often extremely laborious. We shall now shew how these arithmetical calculations may be simplified by the aid of logarithms. Instead of taking as the rate of interest the interest on £100 for one year, it will be found more convenient to take the interest on £1 for one year. If this be denoted by $\mathcal{L}r$, and the amount of £1 for 1 year by $\mathcal{L}R$, we have $R = 1 + r$.

404. *To find the interest and amount of a given sum in a given time at compound interest.*

Let P denote the principal, R the amount of £1 in one year, n the number of years, I the interest, and M the amount.

The amount of P at the end of the first year is PR ; and, since this is the principal for the second year, the amount at the end of the second year is $PR \times R$ or PR^2 . Similarly the amount at the end of the third year is PR^3 , and so on; hence the amount in n years is PR^n ; that is,

$$M = PR^n;$$

and therefore

$$I = P(R^n - 1).$$

Example. Find the amount of £100 in a hundred years, allowing compound interest at the rate of 5 per cent., payable quarterly; having given

$$\log 2 = \cdot 3010300, \quad \log 3 = \cdot 4771213, \quad \log 14 \cdot 3906 = 1 \cdot 15808.$$

The amount of £1 in a quarter of a year is $\mathcal{L} \left(1 + \frac{1}{4} \cdot \frac{5}{100} \right)$ or $\mathcal{L} \frac{81}{80}$.

The number of payments is 400. If M be the amount, we have

$$M = 100 \left(\frac{81}{80} \right)^{400};$$

$$\begin{aligned}\therefore \log M &= \log 100 + 400(\log 81 - \log 80) \\ &= 2 + 400(4 \log 3 - 1 - 3 \log 2) \\ &= 2 + 400(4 \cdot 0053952) = 4 \cdot 15808;\end{aligned}$$

whence

$$M = 14390 \cdot 6.$$

Thus the amount is £14390. 12s.

Note. At simple interest the amount is £600.

405. *To find the present value and discount of a given sum due in a given time, allowing compound interest.*

Let P be the given sum, V the present value, D the discount, R the amount of £1 for one year, n the number of years.

Since V is the sum which, put out to interest at the present time, will in n years amount to P , we have

$$P = VR^n;$$

$$\therefore V = PR^{-n},$$

and

$$D = P - V = P(1 - R^{-n}).$$

Annuities.

406. An **annuity** is a fixed sum paid periodically under certain stated conditions; the payment may be made either once a year or at more frequent intervals. Unless it is otherwise stated we shall suppose the payments annual.

407. *To find the amount of an annuity left unpaid for a given number of years allowing compound interest.*

Let A be the annuity, R the amount of £1 for one year, n the number of years, M the amount.

At the end of the first year A is due, and the amount of this sum in the remaining $n-1$ years is AR^{n-1} ; at the end of the second year another A is due, and the amount of this sum in the remaining $n-2$ years is AR^{n-2} ; and so on.

$$\begin{aligned}\therefore M &= AR^{n-1} + AR^{n-2} + \dots + AR^2 + AR + A \\ &= A(1 + R + R^2 + \dots \text{ to } n \text{ terms}) \\ &= A \frac{R^n - 1}{R - 1}.\end{aligned}$$

408. *To find the present value of an annuity to continue for a given number of years allowing compound interest.*

Let A be the annuity, R the amount of £1 in one year, n the number of years, V the required present value.

The present value of A due in 1 year is AR^{-1} ;
 the present value of A due in 2 years is AR^{-2} ;
 the present value of A due in 3 years is AR^{-3} ; and so on.

[Art. 405.]

Now V is the sum of the present values of the different payments;

$$V = AR^{-1} + AR^{-2} + AR^{-3} + \dots \text{ to } n \text{ terms}$$

$$= AR^{-1} \frac{1 - R^{-n}}{1 - R^{-1}}$$

$$= A \frac{1 - R^{-n}}{R - 1}.$$

Note. This result may also be obtained by dividing the value of M , given in Art. 407, by R^n . [Art. 404.]

Cor. If we make n infinite we obtain for the present value of a perpetual annuity

$$V = \frac{A}{R - 1} = \frac{A}{r}.$$

409. If mA is the present value of an annuity A , the annuity is said to be worth m **years' purchase**.

In the case of a perpetual annuity $mA = \frac{A}{r}$; hence

$$m = \frac{1}{r} = \frac{100}{\text{rate per cent.}};$$

that is, the number of years' purchase of a perpetual annuity is obtained by dividing 100 by the rate per cent.

A good test of the credit of a Government is furnished by the number of years' purchase of its Stocks; thus the $2\frac{3}{4}$ p.c. Consols at $96\frac{1}{4}$ are worth 35 years' purchase; Russian 4 p.c. Stock at 96 is worth 24 years' purchase; while Austrian 5 p.c. Stock at 80 is only worth 16 years' purchase.

410. A **freehold estate** is an estate which yields a perpetual annuity called the *rent*; and thus the value of the estate is equal to the present value of a perpetual annuity equal to the rent.

It follows from Art. 409 that if we know the number of years' purchase that a tenant pays in order to buy his farm, we obtain the rate per cent. at which interest is reckoned by dividing 100 by the number of years' purchase.

E.A.

2A

EXAMPLES XLIII.

1. If in the year 1600 a sum of £1000 had been left to accumulate for 300 years, find its amount in the year 1900, reckoning compound interest at 4 per cent. per annum. Given

$$\log 104 = 2.0170333 \text{ and } \log 12885.5 = 4.10999.$$

2. Find in how many years a sum of money will amount to one hundred times its value at $5\frac{1}{2}$ per cent. per annum compound interest. Given $\log 1055 = 3.023$.

3. Find the present value of £6000 due in 20 years, allowing compound interest at 8 per cent. per annum. Given

$$\log 2 = .30103, \log 3 = .47712, \text{ and } \log 12875 = 4.10975.$$

4. Find at what rate per cent. per annum £1200 will amount to £20000 in 15 years at compound interest. Given

$$\log 2 = .30103, \log 3 = .47712, \text{ and } \log 12063 = 4.08145.$$

5. Find the amount of an annuity of £100 in 15 years, allowing compound interest at 4 per cent. per annum. Given

$$\log 1.04 = .01703, \text{ and } \log 180075 = 5.25545.$$

6. A freehold estate worth £280 a year is sold for £7000; find the rate of interest.

7. If a perpetual annuity is worth 40 years' purchase, find what an annuity of £300 will amount to in 10 years at the same rate of interest. Given $\log 10.25 = 1.01072$, and $\log 1280 = 3.1072$.

8. Find the present value of an annuity of £900 to continue for 20 years at $4\frac{1}{2}$ per cent. compound interest. Given

$$\log 1.045 = .01912, \text{ and } \log 41458 = 4.6176.$$

9. A man borrows £20000 at 5 per cent. compound interest. If the principal and interest are to be paid by 20 equal annual instalments, find the amount of each of these; having given

$$\log 105 = 2.0212, \text{ and } \log 3767 = 3.576.$$

10. A man has a capital of £100000, for which he receives interest at $3\frac{1}{2}$ per cent.; if he spends £7000 a year, find in what time he will be ruined. Given

$$\log 2 = .301, \log 3 = .477, \text{ and } \log 2' = 1.362.$$

MISCELLANEOUS EXAMPLES VI.

1. Simplify $b - \{b - (a + b) - [b - (b - \overline{a - b})] + 2a\}$.
2. Find the sum of $a + b - 2(c + d)$, $b + c - 3(d + a)$, and $c + d - 4(a + b)$.
3. Multiply $\frac{1}{2}x + \frac{2}{3}y$ by $x - \frac{1}{3}y$.
4. If $x=6$, $y=4$, $z=3$, find the value of $\sqrt[3]{2x+3y+z}$.
5. Find the square of $2 - 3x + x^2$.
6. Solve $\frac{x+3}{x-1} + \frac{x-4}{x-6} = 2$.
7. Find the H.C.F. of $a^3 - 2a - 4$ and $a^3 - a^2 - 4$.
8. Simplify $\frac{2a}{a+b} + \frac{2b}{a-b} - \frac{a^2+b^2}{a^2-b^2}$.
9. Solve
$$\left. \begin{aligned} \frac{3}{5}x + \frac{y}{4} &= 13 \\ \frac{1}{3}x - \frac{y}{8} &= 3 \end{aligned} \right\}$$
.
10. Two digits, which form a number, change places when 18 is added to the number, and the sum of the two numbers thus formed is 44: find the digits.
11. If $a=1$, $b=-2$, $c=3$, $d=-4$, find the value of
$$\frac{a^2b^2 + b^2c + d(a-b)}{10a - (c+b)^2}$$
.
12. Subtract $-x^2 + y^2 - z^2$ from the sum of $\frac{1}{3}x^2 + \frac{1}{4}y^2$, $\frac{1}{5}y^2 + \frac{1}{3}z^2$, and $\frac{1}{3}z^2 - \frac{1}{4}x^2$.
13. Write down the cube of $x + 8y$.
14. Simplify $\frac{x^2+xy}{x^2+y^2} \times \frac{x^4-y^4}{xy} \times \frac{y}{x}$.
15. Solve $\frac{3}{5}(2x-7) - \frac{2}{3}(x-8) = \frac{4x+1}{15} + 4$.
16. Find the H.C.F. and L.C.M. of $x^4 + x^3 + 2x - 4$ and $x^3 + 3x^2 - 4$.

17. Find the square root of $4a^4 + 9(1 - 2a) + 3a^2(7 - 4a)$.

18. Solve
$$\left. \begin{aligned} y &= \frac{x+a}{2} + \frac{b}{3} \\ x &= \frac{y+b}{2} + \frac{a}{3} \end{aligned} \right\}$$
.

19. Simplify $\left(\frac{a}{x+a} - \frac{x}{x-a} \right) \div \frac{x^2+a^2}{x^2+ax}$.

20. When 1 is added to the numerator and denominator of a certain fraction the result is equal to $\frac{3}{2}$; and when 1 is subtracted from its numerator and denominator the result is equal to 2: find the fraction.

21. Shew that the sum of $12a+6b-c$, $-7a-b+c$, and $a+b+6c$, is six times the sum of $25a+13b-8c$, $-13a-13b-c$, and $-11a+b+10c$.

22. Divide $x^2 - xy + \frac{3}{16}y^2$ by $x - \frac{1}{4}y$.

23. Add together $18 \left\{ \frac{2x}{9} - \frac{1}{6} \left(\frac{2y}{3} + z \right) \right\}$,
 $24 \left(\frac{3x}{8} - \frac{2y-3z}{12} \right)$, and $30 \left\{ \frac{7z}{15} - \frac{4}{5}(2x-y) \right\}$.

24. Find the factors of

$$(1) 10x^2 + 79x - 8. \quad (2) 729x^6 - y^6.$$

25. Solve $\frac{2x-1}{5} + \frac{5x+3}{17} = 3 - \frac{4x-118}{11}$.

26. Find the value of

$$(5a-3b)(a-b) - b\{3a-c(4a-b) - b^2(a+c)\},$$

$$\text{when } a=0, b=-1, c=\frac{1}{2}.$$

27. Find the H.C.F. of

$$7x^3 - 10x^2 - 7x + 10 \text{ and } 2x^3 - x^2 - 2x + 1.$$

28. Simplify $\frac{x^3 - 7xy + 12y^2}{x^2 + 5xy + 6y^2} \div \frac{x^3 - 5xy + 4y^2}{x^2 + xy - 2y^2}$.

29. Solve
$$\left. \begin{aligned} 3abx + y &= 9b \\ 4abx + 3y &= 17b \end{aligned} \right\}$$
.

30. Find the two times between 7 and 8 o'clock when the hands of a watch are separated by 15 minutes.

31. If $a=1$, $b=-2$, $c=3$, $d=-4$, find the value of

$$\sqrt{a^3-4b+a^2}-\sqrt{c^3+b^3+a+d}.$$
32. Multiply the product of $\frac{1}{4}x^2-\frac{1}{2}xy+y^2$ and $\frac{1}{2}x+y$ by x^3-8y^3 .
33. Simplify by removing brackets

$$a^4-\{4a^3-(6a^2-4a+1)\}$$

$$-[-2-\{a^4-(-4a^3-6a^2-4a)\}-(8a-1)].$$
34. Find the remainder when $5x^4-7x^3+3x^2-x+8$ is divided by $x-4$.
35. Simplify $\frac{x^2+y^2}{x^2-xy} \times \frac{xy-y^2}{x^4-y^4} \times \frac{x}{y}$.
36. Solve
$$\left. \begin{aligned} \frac{x-11}{3}+y &= 18 \\ 2x+\frac{y-13}{4} &= 29 \end{aligned} \right\}.$$
37. Find the square root of $4x^6-12x^4+28x^3+9x^2-42x+49$.
38. Solve $\cdot 006x - \cdot 491 + \cdot 723x = -\cdot 005$.
39. Find the L.C.M. of x^3+y^3 , $3x^2+2xy-y^2$, and $x^3-x^2y+xy^2$.
40. A bill of 25 guineas is paid with crowns and half-guineas, and twice the number of half-guineas exceeds three times that of the crowns by 17: how many of each are used?
41. Simplify

$$(a+b+c)^2-(a-b+c)^2+(a+b-c)^2-(-a+b+c)^2.$$
42. Find the remainder when $a^4-3a^3b+2a^2b^2-b^4$ is divided by $c^2-ab+2b^2$.
43. If $a=0$, $b=1$, $c=-2$, $d=3$, find the value of

$$(3abc-2bcd)^{\frac{2}{3}}/a^3bc-c^3bd+3.$$
44. Find an expression which will divide both $4x^2+3x-10$ and $4x^3+7x^2-3x-15$ without remainder.
45. Simplify
$$\frac{a+\frac{ab}{a-b}}{a^2-\frac{2a^2b^2}{a^2+b^2}} \times \frac{\frac{1}{a^2}-\frac{1}{b^2}}{\frac{1}{a}-\frac{1}{b}}.$$
46. Find the cube root of $8x^3-2x^2y+\frac{xy^2}{6}-\frac{y^3}{216}$.
47. Solve
$$\left. \begin{aligned} 9x+8y &= 43xy \\ 8x+9y &= 42xy \end{aligned} \right\}.$$

48. Simplify $\frac{3}{x-4} - \frac{2}{x-5} - \frac{x-7}{(x-2)(x-3)}$.
49. Find the L.C.M. of $8x^3 + 38x^2 + 59x + 30$
and $6x^3 - 13x^2 - 13x + 30$.
50. A boy spent half of his money in one shop, one-third of the remainder in a second, and one-fifth of what he had left in a third. He had one shilling at last: how much had he at first?
51. Find the remainder when $x^7 - 10x^6 + 8x^5 - 7x^3 + 3x - 11$ is divided by $x^2 - 5x + 4$.
52. Simplify $4 \left\{ a - \frac{3}{2} \left(b - \frac{4c}{3} \right) \right\} \left\{ \frac{1}{2} (2a - b) + 2(b - c) \right\}$.
53. If $a = \frac{25}{16}$, $b = 1$, $c = \frac{3}{4}$, prove that

$$(a - \sqrt{b})(\sqrt{a} + b)\sqrt{a - b} = \frac{3c^4}{\sqrt{a - c^2}}$$
54. Find the L.C.M. of $x^2 - 7x + 12$, $3x^2 - 6x - 9$, and $2x^2 - 6x - 9$.
55. Find the sum of the squares of $ax + by$, $bx - ay$, $ay + bx$, $by - ax$; and express the result in factors.
56. Solve $\frac{x}{6} + \frac{y}{4} = -\frac{3x - 5z}{4} = \frac{z}{8} + \frac{7y}{16} = 1$.
57. Simplify $\frac{a^3 + b^3}{a^4 - b^4} - \frac{a + b}{a^2 - b^2} - \frac{1}{2} \left\{ \frac{a - b}{a^2 + b^2} - \frac{1}{a - b} \right\}$.
58. Solve $x - \left(3x - \frac{2x + 5}{10} \right) = \frac{1}{6} (2x + 67) + \frac{5}{3} \left(1 + \frac{x}{5} \right)$.
59. Add together the following fractions:

$$\frac{2}{x^2 + xy + y^2}, \quad \frac{-4x}{x^3 - y^3}, \quad \frac{x^2}{y^2(x - y)^2}, \quad \frac{-x^2}{x^3y - y^4}.$$
60. A man agreed to work for 30 days, on condition that for every day's work he should receive 3s. 4d., and that for every day's absence from work he should forfeit 1s. 6d.; at the end of the time he received £3. 11s.: how many days did he work?
61. Divide $\frac{3x^5}{4} + 27 - \frac{43x^2}{4} - 4x^4 + \frac{77x^3}{8} - \frac{33x}{4}$ by $\frac{x^2}{2} + 3 - x$.
62. Find the value of

$$\frac{4y}{5}(y - x) - 35 \left[\frac{3x - 4y}{5} - \frac{1}{10} \left\{ 3x - \frac{5}{7}(7x - 4y) \right\} \right]$$

 when $x = -\frac{1}{2}$ and $y = 2$.

63. Simplify $\frac{10x-11}{3(x^2-1)} - \frac{10x-1}{3(x^2+x+1)} + \frac{x^2-2x+5}{(x^3-1)(x+1)}$.

64. Find the cube root of $\frac{a^3c^3}{b^3}x^6 - \frac{3a^2c}{b}x^5 + \frac{3ab}{c}x^4 - \frac{b^3}{c^3}x^3$.

65. Solve $\frac{4x-17}{x-4} + \frac{10x-13}{2x-3} = \frac{8x-30}{2x-7} + \frac{5x-4}{x-1}$.

66. Find the factors of

(1) x^3+5x^2+x+5 . (2) $x^2-2xy-323y^2$.

67. Solve
$$\left. \begin{aligned} \frac{1}{3}(x+y)+2z &= 21 \\ 3x-\frac{1}{2}(y+z) &= 65 \\ x+\frac{1}{2}(x+y-z) &= 38 \end{aligned} \right\}$$

68. Simplify $\frac{x+2y}{\frac{2}{7}x-y} - \frac{3x^2+63xy+70y^2}{2x^2+3xy-35y^2}$.

 69. Find the square root of $-(3b-2c-2a)^3\{2(a+c)-3b\}$.

70. The united ages of a man and his wife are six times the united ages of their children. Two years ago their united ages were ten times the united ages of their children, and six years hence their united ages will be three times the united ages of the children. How many children have they?

71. Find the sum of

$$x^2-3xy-\frac{2}{3}y^2, \quad 2y^2-\frac{2}{3}y^3+z^2, \quad xy-\frac{1}{3}y^2+y^3, \quad \text{and} \quad 2xy-\frac{1}{3}y^3.$$

 72. From $\{(a+b)(a-x)-(a-b)(b-x)\}$ subtract $(a+b)^2-2bx$.

 73. If $a=5$, $b=4$, $c=3$, find the value of

$$\sqrt[2]{6abc+(b+c)^3+(c+a)^3+(a+b)^3-(a+b+c)^3}.$$

74. Find the factors of

(1) $3x^3+6x^2-189x$. (2) $a^2+2ab+b^2+a+b$.

75. Solve

$$\left. \begin{aligned} px &= qy \\ (p+q)x - (q-p)y &= r \end{aligned} \right\}$$

76. Simplify $\frac{x+\frac{y}{2}}{2x^2+xy+\frac{y^2}{2}} - \frac{x^2-\frac{y^2}{2}}{4\left(x^3-\frac{y^3}{8}\right)}$.

77. Solve $\frac{x-7}{x+7} + \frac{1}{2(x+7)} = \frac{2x-15}{2x-6}$.

78. Reduce $\frac{x^4 - x^2 - 2x + 2}{2x^3 - x - 1}$ to its lowest terms.

79. Add together the fractions :

$$\frac{1}{2x^2 - 4x + 2}, \quad \frac{1}{2x^2 + 4x + 2}, \quad \text{and} \quad \frac{1}{1 - x^2}.$$

80. A number consists of three digits, the right-hand one being zero. If the left-hand and middle digits be interchanged the number is diminished by 180; if the left-hand digit be halved, and the middle and right-hand digit be interchanged, the number is diminished by 336: find the number.

81. Divide $1 - 5x + \frac{152}{15}x^3 - \frac{106}{225}x^4 - \frac{28}{9}x^5$ by $1 - x - \frac{14}{15}x^2$.

82. If $p=1$, $q=\frac{1}{2}$, find the value of

$$\frac{(p^2 + q^2) - (p - q)\sqrt{p^2 + 2pq + q^2}}{2p + q - \{p - (q - p)\}}.$$

83. Multiply $\frac{3x^3}{2} - 5x^2 + \frac{x}{4} + 9$ by $\frac{x^2}{2} - x + 3$.

84. Find the L.C.M. of

$$(a^2b - 2ab^2)^2, \quad 2a^2 - 3ab - 2b^2, \quad \text{and} \quad 2(2a^2 + ab)^2.$$

85. Solve $\frac{2x+3}{x+1} = \frac{4x+5}{4x+4} + \frac{3x+3}{3x+1}$.

86. Reduce $\frac{5x^3 - 14x^2 + 16}{3x^3 - 2x^2 + 16x - 48}$ to its lowest terms.

87. Find the square root of

$$4a^4 + 9\left(a^2 + \frac{1}{a^2}\right) + 12a(a^2 + 1) + 18.$$

88. Solve
$$\left. \begin{aligned} \frac{x}{2a} + \frac{y}{3b} &= a + b \\ \frac{3x}{a} - \frac{2y}{b} &= 6(b - a) \end{aligned} \right\}$$

89. Multiply

$$3x + 4y + \frac{11xy}{3} \quad \text{by} \quad 10x - 3y - \frac{11xy}{\frac{x}{4} + y}$$

90. A bag contained five pounds in shillings and half-crowns; after 17 shillings and 6 half-crowns were taken out, thrice as many half-crowns as shillings were left: find the number of each coin.

91. Find the value of

$$5(a-b) \cdot 2\{3a-(a+b)\} + 7\{(a-2b)-(5a-2b)\},$$

$$\text{when } a = -\frac{1}{9}b.$$

92. Divide $3x^4 - 5x^3 + 7x^2 - 11x - 13$ by $3x - 2$.

93. Find the L.C.M. of

$$15(p^3 + q^3), \quad 5(p^2 - pq + q^2), \quad 4(p^2 + pq + q^2), \quad \text{and} \quad 6(p^2 - q^2).$$

94. Resolve into factors:

$$(1) \quad a^3 - 8b^3.$$

$$(2) \quad -x^2 + 2x - 1 + x^4.$$

95. Solve $\frac{x+a}{x+b} = \frac{x+3a}{x+a+b}$.

96. Simplify

$$(1) \quad \frac{35a^2b^2c^2 - 49b^3c^3}{65a^3bc - 91a^3b^2c^2}.$$

$$(2) \quad \frac{y^4 - 7y^3 + 8y^2 - 12y}{2y^2 - 2y - 60}.$$

97. Solve $7x - 9y + 4z = 16$

$$\left. \begin{aligned} \frac{x+y}{3} &= \frac{x+y+z}{2} \end{aligned} \right\}$$

$$2x - 3y + 4z - 5 = 0$$

98. Simplify $\frac{y^2 - \frac{2y}{y-1}}{y^2 - \frac{2y}{y+1}} \div \left(\frac{y^2 - 5y - 6}{y^2 - 6y + 5} \times \frac{y-2}{y+2} \right)$.

99. Find the square root of

$$\frac{4a^2 - 12ab - 8bc + 4ac + 9b^2 + c^2}{4a^2 + 9c^2 - 12ac}.$$

100. The express leaves Bristol at 3 p.m. and reaches London at 6; the ordinary train leaves London at 1.30 p.m. and arrives at Bristol at 6. If both trains travel uniformly, find the time when they will meet.

101. Solve (1) $\cdot 6x + \cdot 75x - \cdot 16 = x - \cdot 583x + 5$.

$$(2) \quad \frac{37}{x^2 - 5x + 6} + \frac{4}{x - 2} = \frac{7}{3 - x}.$$

102. Simplify (1) $\frac{a+x}{a^2+ax+x^2} + \frac{a-x}{a^2-ax+x^2} + \frac{2x^3}{a^4+a^2x^2+x^4}$

(2) $(1+x)^2 \div \left\{ 1 + \frac{x}{1-x + \frac{x}{1+x+x^2}} \right\}$.

103. Find the square root of

$$a^6 + \frac{1}{a^6} - 6\left(a^4 + \frac{1}{a^4}\right) + 15\left(a^2 + \frac{1}{a^2}\right) - 20;$$

also the cube root of the result.

104. Divide $1-2x$ by $1+3x$ to 4 terms.

105. I bought a horse and carriage for £75; I sold the horse at a gain of 5 per cent., and the carriage at a gain of 20 per cent., making on the whole a gain of 16 per cent. Find the original cost of the horse.

106. Find the divisor when $(4a^2+7ab+5b^2)^2$ is the dividend, $8(a+2b)^2$ the quotient, and $b^2(9a+11b)^2$ the remainder.

107. Solve (1) $5x(x-3)=2(x-7)$.

(2) $\frac{1}{(x-1)(x-2)} + 6 = \frac{3}{x-2} + \frac{2}{x-1}$.

108. If $x = a + b + \frac{(a-b)^2}{4(a+b)}$, and $y = \frac{a+b}{4} + \frac{ab}{a+b}$,
prove that $(x-a)^2 - (y-b)^2 = b^2$.

109. Find the square root of

$$49x^4 + \frac{1051x^2}{25} - \frac{14x^3}{5} - \frac{6x}{5} + 9.$$

110. Solve $\frac{a+x}{a^2+ax+x^2} + \frac{a-x}{a^2-ax+x^2} = \frac{3a}{x(a^4+a^2x^2+x^4)}$.

111. Subtract $\frac{x+3}{x^2+x-12}$ from $\frac{x+4}{x^2-x-12}$,
and divide the difference by $1 + \frac{2(x^2-12)}{x^2+7x+12}$.

112. Find the H.C.F. and L.C.M. of

$$2x^2 + (6a-10b)x - 30ab \text{ and } 3x^2 - (9a+15b)x + 45ab.$$

113. Solve (1) $2cx^2 - abx + 2abd = 4cdx$.

(2) $\frac{x}{2(x+3)} - 2\frac{5}{4} = \frac{x^2}{x^2-9} - \frac{8x-1}{4(x-3)}$.

114. If $a=1$, $b=2$, $c=3$, $d=4$, find the value of

$$\frac{a^b + b^c + c^d}{b^a + c^b + d^c + (a+b)(b+c)} + 3(a^a + b^b + c^c) \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right).$$
115. I rode one-third of a journey at 10 miles an hour, one-third more at 9, and the rest at 8 miles an hour; if I had ridden half the journey at 10, and the other half at 8 miles per hour, I should have been half a minute longer on the way: what distance did I ride?
116. The product of two factors is $(3x+2y)^3 - (2x+3y)^3$, and one of the factors is $x-y$; find the other factor.
117. If $a+b=1$, prove that $(a^2-b^2)^2 = a^3 + b^3 - ab$.
118. Resolve into factors:
 (1) $x^3 + y^3 + 3xy(x+y)$. (2) $m^3 - n^3 - m(m^2 - n^2) + n(m-n)^2$.
119. Solve (1) $\left. \begin{aligned} x^3 - y^3 &= 28 \\ x^2 + xy + y^2 &= 7 \end{aligned} \right\}$. (2) $\left. \begin{aligned} x^2 - 6xy + 11y^2 &= 9 \\ x - 3y &= 1 \end{aligned} \right\}$.
120. Find the square root of
 $(a-b)^4 - 2(a^2+b^2)(a-b)^2 + 2(a^4+b^4).$
121. Simplify the fractions:
 (1) $\frac{1}{a^2 - \frac{a^3-1}{a + \frac{1}{a+1}}}$. (2) $\frac{\left(1 + \frac{1}{x}\right) \times \left(1 - \frac{1}{x}\right)^2}{x - \frac{1}{x}}.$
122. Find the H.C.F. of
 $a^2b + b^2c - abc - ab^2$ and $ax^2 + ab - a^2 - bx^2$.
123. A constituency had two-thirds of its number Conservatives; in an election 25 refused to vote, and 60 went over to the Liberals; the voters were now equal. How many voters were there altogether?
124. Solve (1) $\frac{x^2}{a+b} + (a-b) = \frac{2ax}{a+b}$.
 (2) $\frac{3}{x} + \frac{2}{y} = 6 \left(\frac{1}{y} - \frac{1}{2x} \right) = 2.$
125. Simplify (1) $\left(1 + \frac{y^2 + z^2 - x^2}{2yz}\right) \div \left(1 - \frac{x^2 + y^2 - z^2}{2xy}\right).$
 (2) $\frac{(x+1)^3 - (x-1)^3}{(x+1)^4 - (x-1)^4}.$
126. Divide

$$\frac{x^4 + (a-1)x^3 - (2a+1)x^2 + (a^2+4a-5)x + 3a+6}{x^2 - 3x + a + 2}.$$

127. Resolve into factors :

$$(1) x^3 + 5xy - 24y^2 + x - 3y. \quad (2) x^3 - \frac{4}{x}$$

128. Find the square root of $p^2 - 3q$ to three terms.

$$129. \text{ Solve } (1) \frac{x-5}{x-6} - \frac{x-6}{x-7} = \frac{x-1}{x-2} - \frac{x-2}{x-3}$$

$$(2) ax+1=by+1=ay+bx.$$

130. Find the H.C.F. of $3x^2 + (4a-2b)x - 2ab + a^2$ and $x^3 + (2a-b)x^2 - (2ab-a^2)x - a^2b$.

131. Simplify

$$(1) \frac{(x^a)^3}{x^{b+c}} \times \frac{(x^b)^3}{x^{c+a}} \times \frac{(x^c)^3}{x^{a+b}}. \quad (2) x^{\frac{1}{2}}y^{\frac{1}{3}} \left(\frac{y^{\frac{1}{3}}}{x^{\frac{1}{6}}} \right)^2 \div \frac{y^{-\frac{1}{4}}}{x^{\frac{1}{4}}}$$

132. At a cricket match the contractor provided dinner for 24 persons, and fixed the price so as to gain $12\frac{1}{2}$ per cent. upon his outlay. Three of the cricketers being absent, the remaining 21 paid the fixed price for their dinner, and the contractor lost 1s. : what was the charge for the dinner ?

133. Prove that $x(y+2) + \frac{x}{y} + \frac{y}{x}$ is equal to a , if

$$x = \frac{y}{y+1} \text{ and } y = \frac{a-2}{2}.$$

134. Find the cube root of

$$x^3 - 12x^2 + 54x - 112 + \frac{108}{x} - \frac{48}{x^2} + \frac{8}{x^3}.$$

135. Find the H.C.F. and L.C.M. of

$$x^3 + 2ax^2 + a^2x + 2a^3 \text{ and } x^3 - 2ax^2 + a^2x - 2a^3.$$

136. Simplify

$$(1) 42 \left\{ \frac{4x-3y}{6} - \frac{3x-4y}{7} \right\} - 56 \left\{ \frac{3x-2y}{7} - \frac{2x-3y}{8} \right\},$$

$$(2) \frac{4b+a}{3b+a} + \frac{a-4b}{a-3b} + \frac{a^2-3b^2}{a^2-9b^2}.$$

137. Resolve $4a^2(x^3+18ab^3) - (32a^5+9b^5x^3)$ into four factors.

138. Solve (1) $5\sqrt{3x-1} = \sqrt{75x-29}$.

$$(2) \frac{xy}{x+y} = 70, \quad \frac{xz}{x+z} = 84, \quad \frac{yz}{y+z} = 140.$$

139. Shew that the difference between

$$\frac{x}{x-a} + \frac{x}{x-b} + \frac{x}{x-c} \quad \text{and} \quad \frac{a}{x-a} + \frac{b}{x-b} + \frac{c}{x-c}$$

is the same whatever value x may have.

140. Multiply $x^{\frac{3}{2}} + 2y^{\frac{3}{2}} + 3z^{\frac{3}{2}}$ by $x^{\frac{3}{2}} - 2y^{\frac{3}{2}} - 3z^{\frac{3}{2}}$.

141. Walking $4\frac{1}{4}$ miles an hour, I start $1\frac{1}{2}$ hours after a friend whose pace is 3 miles an hour: how long shall I be in overtaking him?

142. Express in the simplest form

$$(1) (8^{\frac{3}{2}} + 4^{\frac{3}{2}}) \times 16^{-\frac{3}{2}}. \quad (2) \frac{\left\{ 9^n \cdot 3^2 \times \frac{1}{3^{-n}} \right\} - 27^n}{3^{2n} \times 9}.$$

143. Find the square root of

$$\frac{x}{y} + \frac{y}{x} + 3 - 2\sqrt{\frac{x}{y}} - 2\sqrt{\frac{y}{x}}.$$

144. Simplify

$$(1) \left(\frac{x}{x-1} - \frac{1}{x+1} \right) \cdot \frac{x^3-1}{x^6+1} \cdot \frac{(x-1)^2(x+1)^2+x^2}{x^4+x^2+1}.$$

$$(2) \left\{ \frac{a^4-y^4}{a^3-2ay+y^3} \div \frac{a^2+ay}{a-y} \right\} \times \left\{ \frac{a^5-a^3y^2}{a^3+y^3} \div \frac{a^4-2a^2y+y^2}{a^2-ay+y^2} \right\}.$$

145. Find the value of

$$(1) \sqrt{8} + \sqrt{50} - \sqrt{18} + \sqrt{48}. \quad (2) \sqrt{35} + 14\sqrt{6}.$$

146. Solve $(1) \frac{x-b}{x-a} - \frac{x-a}{x-b} = \frac{2(a-b)}{x-(a+b)}.$

$$(2) \left. \begin{aligned} 2x+3y &= 1\frac{1}{2} \\ 4x^2+9xy+9y^2 &= 11 \end{aligned} \right\}.$$

147. Shew that

$$\frac{(a+b)^3-c^3}{(a+b)-c} + \frac{(b+c)^3-a^3}{b+c-a} + \frac{(c+a)^3-b^3}{c+a-b}$$

is equal to $2(a+b+c)^2 + a^2 + b^2 + c^2.$

148. Divide $a-x+4a^{\frac{1}{2}}x^{\frac{3}{2}}-4a^{\frac{1}{2}}x^{\frac{5}{2}}$

$$\text{by} \quad a^{\frac{1}{2}}+2a^{\frac{1}{2}}x^{\frac{1}{2}}-x^{\frac{1}{2}}.$$

149. Find the square root of

$$(a-1)^4 + 2(a^4+1) - 2(a^2+1)(a-1)^2.$$

150. How much are pears a gross when 120 more for a sovereign lowers the price 2*d.* a score?
151. Shew that if a number of two digits is six times the sum of its digits, the number formed by interchanging the digits is five times their sum.
152. Find the value of

$$\frac{1}{(a-b)(b-c)} - \frac{1}{(b-c)(a-c)} - \frac{1}{(c-a)(b-a)}$$

153. Multiply

$$3+5x - \frac{12+41x+36x^2}{4+7x} \text{ by } 5-2x + \frac{26x-8x^2-14}{3-4x}.$$

154. If $x - \frac{1}{x} = 1$, prove that $x^2 + \frac{1}{x^2} = 3$, and $x^3 - \frac{1}{x^3} = 4$.

155. Solve
- $$\left. \begin{array}{l} (1) \frac{3x}{11} + \frac{23}{x+4} = \frac{1}{3}(x+5). \\ (2) \begin{array}{l} 2x^2 - 3y^2 = 23 \\ 2xy - 3y^2 = 3 \end{array} \right\}.$$

156. Simplify

$$(1) 1\frac{3}{5}\sqrt{20} - 3\sqrt{5} - \sqrt{\frac{1}{5}} \quad (2) \frac{\sqrt{x}\left(\frac{\sqrt[4]{y}}{x^{\frac{1}{2}}}\right)}{y^{-\frac{1}{2}}\left(x^{\frac{1}{2}}\right)} \div \frac{y^{-\frac{1}{2}}}{x^{\frac{1}{2}}}.$$

157. Find the H.C.F. of $(p^2-1)x^3 + (3p-1)x - p(p-1)$ and $p(p+1)x^2 - (p^2-2p-1)x - (p-1)$.

158. Reduce to its simplest form

$$\frac{ax + \frac{a}{y}}{x - \frac{1}{y}} \times \frac{x^2 + \frac{1}{y^2}}{bx^2 - \frac{b}{y^2}} \times \frac{\frac{1}{5}(xy-1)^2}{\frac{1}{3}(x^4y^4-1)}.$$

159. Find the square root of

$$(1) 1 - 2^{2n+1} + 4^{2n}. \quad (2) 9^n - 2 \cdot 6^n + 4^n.$$

160. A clock gains 4 minutes a day. What time should it indicate at 6 o'clock in the morning, in order that it may be right at 7.15 p.m. on the same day?

161. If $x = 2 + \sqrt{2}$, find the value of $x^2 + \frac{4}{x^2}$.

162. Solve

$$(1) \frac{\sqrt{x+a}}{\sqrt{x-b}} = \frac{\sqrt{x-a}}{\sqrt{x}}. \quad (2) \frac{\sqrt{1+x} + \sqrt{1-x}}{\sqrt{1+x} - \sqrt{1-x}} = 3.$$

163. Simplify

$$\frac{a^2}{(b-a)(c-a)} + \frac{b^2}{(c-b)(a-b)} + \frac{c^2}{(a-c)(b-c)}.$$

164. Find the product of $\frac{1}{5}\sqrt{5}$, $\frac{1}{2}\sqrt[3]{2}$, $\sqrt[4]{80}$, $\sqrt[5]{5}$, and divide

$$\frac{8-4\sqrt{5}}{\sqrt{5}+1} \text{ by } \frac{3\sqrt{5}-7}{5+\sqrt{7}}.$$

165. Resolve $9x^6y^2 - 576y^2 - 4x^3 + 256x^2$ into six factors.

166. Simplify

$$(1) \frac{1 - \frac{a^2}{(x+a)^2}}{(x+a)(x-a)} \div \frac{x(x+2a)}{(x^2-a^2)(x+a)^2}.$$

$$(2) \frac{6x^2y^2}{m+n} \div \left[\frac{3(m-n)x}{7(r+s)} \div \left\{ \frac{4(r-s)}{21xy^2} \div \frac{r^3-s^2}{4(m^2-n^2)} \right\} \right].$$

167. Simplify

$$(1) \left(a^{1+\frac{q}{p}} \right)^{\frac{p}{p+q}} \div \sqrt[p]{\frac{a^{2p}}{(a^{-1})^{-p}}}.$$

$$(2) \sqrt{14} - \sqrt{132}.$$

168. Find the H.C.F. and L.C.M. of

$$20x^4 + x^2 - 1, \quad 25x^4 + 5x^3 - x - 1, \quad 25x^4 - 10x^3 + 1.$$

169. Solve

$$(1) a + x + \sqrt{2ax + x^2} = b.$$

$$(2) x + 9\frac{5}{8} + \frac{1}{x + \frac{11}{8}} = 8.$$

170. The price of photographs is raised 3s. per dozen, and customers consequently receive seven less than before for a guinea: what were the prices charged?

171. If $\left(a + \frac{1}{a}\right)^2 = 3$, prove that $a^3 + \frac{1}{a^3} = 0$.

172. Find the value of

$$\frac{x+2a}{2b-x} + \frac{x-2a}{2b+x} + \frac{4ab}{x^2-4b^2}, \text{ when } x = \frac{ab}{a+b}.$$

173. Reduce to fractions in their lowest terms.

$$(1) \left(\frac{1}{x} + \frac{1}{y} + \frac{1}{z} \right) \div \left(\frac{x+y+z}{x^2+y^2+z^2-xy-yz-zx} - \frac{1}{x+y+z} \right) + 1.$$

$$(2) \left(1 - \frac{56}{x+4} + \frac{42}{x+3} \right) \left(1 + \frac{56}{x-4} - \frac{42}{x-3} \right).$$

174. Express as a whole number

$$(27)^{\frac{2}{3}} + (16)^{\frac{1}{2}} - \frac{2}{(8)^{-\frac{2}{3}}} + \frac{5/2}{(4)^{-\frac{1}{2}}}.$$

175. Simplify

$$(1) \frac{n}{1-x^n} + \frac{n}{1-x^{-n}} \quad (2) \sqrt[n]{97-56}\sqrt{3}.$$

176. Solve

$$(1) \frac{x-4a}{x-3a} + \frac{x-5a}{x-4a} = \frac{x+6a}{x-4a} + \frac{x+5a}{x-3a}.$$

$$(2) \left. \begin{aligned} 3x^2 + xy + 3y^2 &= 8\frac{1}{4} \\ 8x^2 - 3xy + 8y^2 &= 17\frac{3}{4} \end{aligned} \right\}.$$

177. Find the square root of $\frac{a^2x^2+2a^{1/2}x^3+b^4x^4}{a^{2n}+2a^nx^n+x^{2n}}.$

178. Simplify

$$(1) \frac{b}{\sqrt{a}} \times \sqrt[3]{ac} \times \frac{\sqrt[3]{c^3}}{\sqrt{b}} \times \frac{\sqrt{b^{-1}}}{a^{-\frac{1}{2}}}. \quad (2) \left\{ \frac{(9^{n+1}) \times \sqrt{3 \cdot 3^n}}{3\sqrt{3^{-n}}} \right\}^{\frac{1}{n}}.$$

179. A boat's crew can row 8 miles an hour in still water; what is the speed of a river's current if it take them 2 hours and 40 minutes to row 8 miles up and 8 miles down?

180. If $a = x^2 - yz$, $b = y^2 - zx$, $c = z^2 - xy$, prove that $a^2 - bc = x(ax + by + cz).$ 181. Find a quantity such that when it is subtracted from each of the quantities a , b , c , the remainders are in continued proportion.

182. Simplify

$$(1) \left(x + y - \frac{1}{x+y - \frac{xy}{x+y}} \right) \times \frac{x^3 - y^3}{x^2 - y^2}.$$

$$(2) \frac{2(7x-4)}{6x^2-7x+2} + \frac{x-10}{6x^2-x-2} - \frac{2(4x-1)}{4x^2-1}.$$

183. Find the sixth root of

$$729 - 2916x^2 + 4860x^4 - 4320x^6 + 2160x^8 - 576x^{10} + 64x^{12}.$$

184. Simplify

$$(1) \frac{1}{x + \sqrt{x^2 - 1}} + \frac{1}{x - \sqrt{x^2 - 1}}.$$

$$(2) \sqrt[4]{16} + \sqrt[3]{81} - \sqrt[3]{-512} + \sqrt[3]{192} - 7\sqrt[3]{9}.$$

185. Solve

$$(1) \frac{5}{6 - \frac{5}{6 - x}} = x.$$

$$(2) \begin{cases} x^2y^2 + 192 = 28xy \\ x + y = 8 \end{cases}.$$

186. Simplify

$$\frac{b-c}{a^2 - (b-c)^2} + \frac{c-a}{b^2 - (c-a)^2} + \frac{a-b}{c^2 - (a-b)^2}.$$

187. Solve

$$(1) x - 15\frac{3}{4} + \frac{5}{x - 15\frac{3}{4}} = 6.$$

$$(2) 2(x + y^{-1}) = 3(x^{-1} - y) = 4.$$

 188. If $xy = ab(a+b)$ and $x^2 - xy + y^2 = a^3 + b^3$ prove that

$$\left(\frac{x}{a} - \frac{y}{b}\right)\left(\frac{x}{b} - \frac{y}{a}\right) = 0.$$

189. Find the H.C.F. of

$$(2a^2 - 3a - 2)x^2 + (a^2 + 7a + 2)x - a^2 - 2a$$

and

$$(4a^2 + 4a + 1)x^2 - (4a^2 + 2a)x + a^2.$$

190. Multiply

$$\sqrt{2x} + \sqrt{2(2x-1)} - \frac{1}{\sqrt{2x}}$$

by

$$\frac{1}{\sqrt{2x}} + \sqrt{2(2x-1)} - \sqrt{2x}.$$

191. Divide

$$a^4b^2 + b^4c^2 + c^4a^2 - a^2b^4 - b^2c^4 - c^2a^4$$

by

$$a^2b + b^2c + c^2a - ab^2 - bc^2 - ca^2.$$

192. Simplify

$$(1) \frac{7}{2(x+1)} - \frac{1}{6(x-1)} - \frac{10x-1}{3(x^2+x+1)}.$$

$$(2) \left\{ \frac{\sqrt{x+a}}{\sqrt{x-a}} - \frac{\sqrt{x-a}}{\sqrt{x+a}} \right\} \times \frac{\sqrt{x^3-a^3}}{\sqrt{(x+a)^2-ax}}$$

E A.

B

193. If p be the difference between any quantity and its reciprocal, q the difference between the square of the same quantity and the square of its reciprocal, shew that

$$p^2(p^2+4)=q^2.$$

194. A man started for a walk when the hands of his watch were coincident between three and four o'clock. When he finished, the hands were again coincident between five and six o'clock. What was the time when he started, and how long did he walk?

195. If n be an integer, shew that $7^{2n+1}+1$ is always divisible by 8.

196. Simplify
$$\frac{\left(p+\frac{1}{q}\right)^p \left(p-\frac{1}{q}\right)^q}{\left(q+\frac{1}{p}\right)^p \left(q-\frac{1}{p}\right)^q}.$$

197. Find the value of

$$(1) \frac{7+3\sqrt{5}}{7-3\sqrt{5}} + \frac{7-3\sqrt{5}}{7+3\sqrt{5}}.$$

$$(2) \frac{\sqrt{1+x}+\sqrt{1-x}}{\sqrt{1+x}-\sqrt{1-x}} \text{ when } x=\frac{2b}{b^2+1}.$$

198. If $a+b+c+d=2s$, prove that

$$4(ab+cd)^2-(a^2+b^2-c^2-d^2)^2=16(s-a)(s-b)(s-c)(s-d).$$

199. A man buys a number of articles for £1, and sells for £1. 1s. all but two at 2d. apiece more than they cost: how many did he buy?

200. Find the square root of

$$2(81x^4+y^4)-2(9x^2+y^2)(3x-y)^2+(3x-y)^4.$$

201. If $x:a::y:b::z:c$, prove that

$$(bc+ca+ab)^2(x^2+y^2+z^2)=(bz+cx+ay)^2(a^2+b^2+c^2).$$

202. If a man save £10 more than he did the previous year, and if he saved £20 the first year, in how many years will his savings amount to £1700?

203. Given that 4 is a root of the quadratic $x^2-5x+q=0$, find the value of q and the other root.

204. A person having 7 miles to walk increases his speed one mile an hour after the first mile, and finds that he is half an hour less on the road than he would have been had he not altered his rate. How long did he take?

205. If $(a+b+c)x = (-a+b+c)y = (a-b+c)z = (a+b-c)w$,

shew that
$$\frac{1}{y} + \frac{1}{z} + \frac{1}{w} = \frac{1}{x}.$$

206. Find a Geometrical Progression of which the sum of the first two terms is $2\frac{2}{3}$, and the sum to infinity $4\frac{1}{6}$.

207. Simplify
$$\frac{\left(1 + \frac{x}{y}\right)^m \left(1 - \frac{y}{x}\right)^n}{\left(1 + \frac{y}{x}\right)^n \left(1 - \frac{x}{y}\right)^m}$$

208. A man has a stable containing 10 stalls; in how many ways could he stable 5 horses?

209. In boring a well 400 feet deep the cost is 2s. 3d. for the first foot and an additional penny for each subsequent foot: what is the cost of boring the last foot, and also of boring the entire well?

210. If α, β are the roots of $x^2 + px + q = 0$, shew that p, q are the roots of the equation

$$x^2 + (a + \beta - \alpha\beta)x - \alpha\beta(a + \beta) = 0.$$

211. Multiply together the duodenary numbers *tele* and *ete*.

212. If $\frac{x+z}{y} = \frac{z}{x} = \frac{x}{z-y}$; determine the ratios $x : y : z$.

213. If α, b, c are in H.P. shew that

$$\left(\frac{3}{a} + \frac{3}{b} - \frac{2}{c}\right)\left(\frac{3}{c} + \frac{3}{b} - \frac{2}{a}\right) + \frac{9}{b^2} = \frac{25}{ac}.$$

214. Find the number of permutations which can be made from all the letters of the words

(1) *Consequences*, (2) *Acarnania*.

215. Expand by the Binomial Theorem $(2a-3x)^n$; and find the numerically greatest term in the expansion of $(1+x)^n$, if $x = \frac{3}{5}$, and $n=7$.
216. When $x = \frac{\sqrt{3}}{4}$, find the value of
$$\frac{1+2x}{1+\sqrt{1+2x}} + \frac{1-2x}{1-\sqrt{1-2x}}.$$
217. Simplify $\frac{x^2-bc}{(a-b)(a-c)} + \frac{x^2-ca}{(b-c)(b-a)} + \frac{x^2-ab}{(c-a)(c-b)}.$
218. Solve the equations:
- (1) $(x^2-5x+2)^2 = x^2-5x+22.$
 - (2) $\left(x^2 + \frac{1}{x^2}\right)^2 + 4\left(x^2 + \frac{1}{x^2}\right) = 12.$
219. Prove that
$$(y-z)^3 + (x-y)^3 + 3(x-y)(x-z)(y-z) = (x-z)^3.$$
220. Out of 16 consonants and 5 vowels, how many words can be formed each containing 4 consonants and 2 vowels?
221. If $b-a$ is a harmonic mean between $c-a$ and $d-a$, shew that $d-c$ is a harmonic mean between $a-c$ and $b-c$.
222. In how many ways may 2 red balls, 3 black, 1 white, 2 blue be selected from 4 red, 6 black, 2 white and 5 blue; and in how many ways may they be arranged?
223. The sum of a certain number of terms of an arithmetical series is 36, and the first and last of these terms are 1 and 11 respectively: find the number of terms, and the common difference of the series.
224. Expand by the Binomial Theorem
- (1) $\left(2 - \frac{3a}{4}\right)^5$; (2) $\left(1 - \frac{2}{3}x\right)^{\frac{2}{3}}$ to 5 terms.
225. In what scale is the denary number 418 represented by 1534?

226. Simplify $\frac{2\sqrt{10}}{3\sqrt{27}} \times \frac{15\sqrt{21}}{4\sqrt{15}} \div \frac{5\sqrt{14}}{7\sqrt{48}}$; and find the value of $\frac{1}{3\sqrt{5}-6}$, given that $\sqrt{5}=2.236$.

227. By the Binomial Theorem find the cube root of 128 to six places of decimals.

228. There are 9 books of which 4 are Greek, 3 are Latin, and 2 are English; in how many ways could a selection be made so as to include at least one of each language?

229. Simplify

$$(1) \frac{\sqrt{45x^3} - \sqrt{80x^3} + \sqrt{5a^2x}}{a-x};$$

$$(2) \left\{ \frac{x^{\frac{1}{2}} + x^{-\frac{1}{2}}}{x^2 - x + 1} - \frac{x^{\frac{1}{2}} - x^{-\frac{1}{2}}}{x^2 + x + 1} \right\} \div \left\{ \frac{x^{\frac{1}{2}} + 2x^{-\frac{1}{2}}}{x^3 - 1} - \frac{2x^{-\frac{1}{2}}}{x^3 + 1} \right\}.$$

230. Form the quadratic equation whose roots are $5 \pm \sqrt{6}$.

If the roots of $x^2 - px + q = 0$ are two consecutive integers prove that $p^2 - 4q - 1 = 0$.

231. Subtract 4.72473 from 7.641 in the scale of eight. Find the square root of 108404 in the scale of twelve.

232. Find $\log_{16} 128$, $\log_4 \sqrt{128}$, $\log_2 \frac{1}{4}$; and having given

$$\log 2 = .3010300 \text{ and } \log 3 = .4771213,$$

find the logarithm of .00001728.

233. A and B start from the same point, B five days after A; A travels 1 mile the first day, 2 miles the second, 3 miles the third, and so on; B travels 12 miles a day. When will they be together? Explain the double answer.

234. Solve the equations:

$$(1) 2^x = 8^{y+1}, \quad 9^y = 3^{x-6};$$

$$(2) x^z = y^{2x}, \quad 2^z = 2 \times 4^x, \quad x + y + z = 16.$$

235. The sum of the first 10 terms of an arithmetical series is to the sum of the first 5 terms as 13 is to 4. Find the ratio of the first term to the common difference.

236. Find the greatest term in the expansion of $(1-x)^{-\frac{1}{3}}$ when $x = \frac{12}{13}$.
237. Five gentlemen and one lady wish to enter an omnibus in which there are only three vacant places; in how many ways can these places be occupied (1) when there is no restriction, (2) when one of the places is to be occupied by the lady?
238. Given $\log 2 = .301030$, $\log 3 = .477121$, and $\log 7 = .845098$, find the logarithms of .005, 6.3, and $\left(\frac{49}{216}\right)^{\frac{1}{3}}$.
Find x from the equation $18^3 - 4x = (54\sqrt{2})^{3x-2}$.
239. If P and Q vary respectively as $y^{\frac{1}{3}}$ and $y^{\frac{1}{3}}$ when z is constant, and as $z^{\frac{1}{3}}$ and $z^{\frac{1}{3}}$ when y is constant, and if $x = P + Q$, find the equation between x , y , z ; it being known that when $y = z = 64$, $x = 12$; and that when $y = 4z = 16$, $x = 2$.
240. Simplify

$$\log \frac{133}{65} + 2 \log \frac{13}{7} - \log \frac{143}{90} + \log \frac{77}{171}.$$
241. If the number of permutations of n things 4 at a time is to the number of combinations of $2n$ things 3 at a time as 22 to 3, find n .
242. If $\frac{1}{a} + \frac{1}{c} = \frac{1}{2b-a} + \frac{1}{2b-c}$, prove that $2b$ is either the arithmetic mean between $2a$ and $2c$, or the harmonic mean between a and c .
243. If ${}^nC_r$ denote the number of combinations of n things taken r together, prove that

$${}^{n+2}C_{r+1} = {}^nC_{r+1} + {}^nC_{r-1} + (2 \times {}^nC_r).$$
244. Find (1) the characteristic of $\log 54$ to base 3;
 (2) $\log_{10} (.0125)^{\frac{1}{3}}$; (3) the number of digits in 3^4 .
 Given $\log_{10} 2 = .30103$, $\log_{10} 3 = .47712$.

245. Write down the $(r+1)^{\text{th}}$ term of $(2ax^2 - x^3)^{\frac{1}{2}}$, and express it in its simplest form.

246. At a meeting of a Debating Society there were 9 speakers; 5 spoke for the Government, and 4 for the Opposition. In how many ways could the speeches have been made, if a member of the Government always speaks first, and the speeches are alternately for the Government and the Opposition?

247. Form the quadratic equation whose roots are

$$a + b + \sqrt{a^2 + b^2} \text{ and } \frac{2ab}{a + b + \sqrt{a^2 + b^2}}.$$

248. A point moves with a speed which is different in different miles, but invariable in the same mile, and its speed in any mile varies inversely as the number of miles travelled before it commences this mile. If the second mile be described in 2 hours, find the time taken to describe the n^{th} mile.

249. Solve the equations:

$$(1) \quad x^2(b-c) + ax(c-a) + a^2(a-b) = 0,$$

$$(2) \quad (x^3 - px + p^2)(qx + pq + p^2) = qx^3 + p^2q^2 + p^4.$$

250. Prove by the Binomial Theorem that

$$1 + \frac{3}{4} + \frac{3 \cdot 5}{4 \cdot 8} + \frac{3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12} + \dots \text{ad inf.} = \sqrt[3]{8}.$$

THE UNIVERSITY OF CHICAGO PRESS
PUBLISHED BY THE UNIVERSITY OF CHICAGO PRESS
CHICAGO, ILL. 60637

ANSWERS.

I. a. PAGE 4.

1. 70.	2. 125.	3. 105.	4. 343.	5. 30.
6. 32.	7. 12.	8. 70.	9. 6.	10. 108.
11. 7.	12. 144.	13. 48.	14. 189.	15. 200.
16. 27.	17. 1000.	18. 3.	19. 1.	20. 567.
21. 4.	22. 125.	23. 81.	24. 1.	25. 243.
26. 512.	27. 5.	28. 4096.	29. 64.	30. 90.
31. 24.	32. 2.	33. 81.	34. 64.	35. 8.
36. 2401.	37. 56.	38. 3.	39. 48.	40. 16.

I. b. PAGE 5.

1. 700.	2. 686.	3. 96.	4. 135.	5. 15.
6. 60.	7. 162.	8. 0.	9. 0.	10. 3000.
11. 98.	12. 225.	13. 3.	14. 1.	15. $5\frac{1}{3}$.
16. 2.	17. 36.	18. 160.	19. $1\frac{1}{8}$.	20. 40.
21. 0.	22. 72.	23. 2048.	24. 81.	25. $11\frac{1}{4}$.
26. $13\frac{1}{2}$.	27. $\frac{1}{64}$.	28. $3\frac{3}{8}$.		

I. c. PAGE 6.

1. 4.	2. 6.	3. 8.	4. 36.	5. 3.
6. 8.	7. 8.	8. 0.	9. 32.	10. 60.
11. 0.	12. 16.	13. $2\frac{2}{3}$.	14. $3\frac{1}{3}$.	15. $1\frac{1}{7}$.
16. 0.	17. $\frac{2}{3}$.	18. $2\frac{2}{3}$.	19. 0.	20. 3.
21. 8.	22. 12.	23. 105.	24. 16.	25. $\frac{1}{40}$.
26. $\frac{1}{24}$.	27. $\frac{1}{48}$.	28. $\frac{1}{20}$.	29. $\frac{1}{10}$.	30. $\frac{1}{4}$.
31. $\frac{5}{6}$.	32. $\frac{1}{10}$.	33. 9.	34. 64.	35. 1.
36. 3.				

E. A.

I. d. PAGE 8.

1. 19.	2. 0.	3. 7.	4. 11.	5. 21.
6. 6.	7. 18.	8. 36.	9. 6.	10. 14.
11. 85.	12. 96.	13. 36.	14. 0.	15. 0.
16. 12.	17. 24.	18. 43.	19. 4.	20. 8.
21. 12.	22. 0.	23. 1.	24. 6000.	25. $3\frac{1}{3}$.
26. 10.	27. $5\frac{1}{2}$.	28. $5\frac{1}{3}$.	29. 37.	30. 6.
31. $4\frac{1}{3}$.	32. $5\frac{1}{8}$.			

II. PAGE 12.

1. $47a$.	2. $24x$.	3. $39b$.	4. $151c$.	5. $-26x$.
6. $-40b$.	7. $-17y$.	8. $-66c$.	9. $-20b$.	10. $2x$.
11. 0.	12. $-16f$.	13. $-s$.	14. $7y$.	15. 0.
16. $2ab$.	17. x^2 .	18. $-14a^2x$.	19. $-21a^3$.	20. $-16x^3$.
21. 0.	22. $-19x^4$.	23. $-43abcd$.	24. $\frac{11}{6}x$.	25. $\frac{8}{5}a$.
26. $-3b$.	27. $-x^2$.	28. $-\frac{5}{6}ab$.	29. $\frac{5}{4}x$.	30. $-5x^2$.

III. a. PAGE 16.

1. 0.	2. $4a+4b+4c$.	3. 0.
4. $4x+4y+4z$.	5. $3a+5b-2c$.	6. $b-c$.
7. $39a-5b+4c$.	8. $5c$.	9. $3ax-3by+3cz$.
10. $22p-18q-20r$.	11. ab .	12. $-20ab+ca$.
13. $5ab+bc$.	14. $pq+qr+rp$.	15. $6x$.
16. $20a$.	17. $2xy+2zx$.	18. $14ab-11bc$.
19. $13z$.	20. $a+b+c$.	

III. b. PAGE 18.

1. abc .	2. x^2+xy+y^2 .	3. $a^2+3ab-2b^2$.
4. $yz+zx+xy$.	5. $3x^2+2xy-y^2$.	6. $-2x^3+x^2+4x+2$.
7. x^2-7x .	8. $15x^2-32x-18$.	9. $15x^3-4x^2+3x-1$.
10. $a^3+b^3+c^3$.	11. $a^3+b^3+c^3+d^3$.	12. x^3+x^2+x+3 .
13. $9a^3-3a^2$.	14. $3x^2-2y^2-2xy-4yz-3xz$.	
15. $-x^3+x^2+2y^2+y$.	16. $3x^2y+xy^2$.	17. $2a^2b$.
18. $x^5-x^4y-y^5$.	19. $a^3+b^3+c^3-3abc$.	

11. 5. 21.
6. 10. 14.
0. 15. 0.
4. 20. 8.
6000. 25. $3\frac{1}{3}$.
37. 30. 6.

20. $x^3 + x^2y + 7xy^2 + 3y^3$.

21. $\frac{1}{4}a - \frac{2}{3}b$.

22. $-3a - \frac{1}{2}b$.

23. $-\frac{7}{3}a + \frac{2}{3}b - \frac{1}{2}c$.

24. $\frac{3}{8}a - \frac{4}{5}b - \frac{15}{4}c$.

25. $\frac{1}{3}x^2 - \frac{4}{3}xy + \frac{1}{2}y^2$.

26. $\frac{5}{6}a^2 + \frac{3}{5}ab - \frac{1}{6}b^2$.

27. $\frac{3}{8}x^2 - \frac{2}{5}xy - \frac{1}{2}y^2$.

28. $-\frac{1}{4}x^3 + \frac{3}{8}ax^2 + \frac{5}{8}a^2x$.

29. $-\frac{1}{4}x^2 - xy + \frac{3}{5}y^2$.

30. $-a^3 - \frac{1}{2}a^2b + \frac{1}{4}ab^2 + b^3$.

IV. a. PAGE 21.

- 151c. 5. $-26x$.
 $-20b$. 10. $2x$.
 $7y$. 15. 0.
 $-21a^3$. 20. $-16x^3$.
 $\frac{11}{6}x$. 25. $\frac{8}{5}a$.
 $\frac{5}{4}x$. 30. $-5x^2$.

1. $-2a - 2c$.

2. $3a - 5b - 4c$.

3. $13x + 18y - 19z$.

4. $-5a + 30b - 4c$.

5. $11x + 13y - 16z$.

6. $12ab - 10bc - 10cd$.

7. $21a - 13b - 33c$.

8. $11x + 26y + 22z$.

9. $2ac + 2bd$.

10. $2ab - 2cd + 2ac - 2bd$.

11. $-cd - ac - bd$.

12. $2xy$.

13. $-3x^3 - x^2 - 2x + 1$.

14. $-12x^2y + 21xy^2 + 15xyz$.

15. $\frac{1}{6}a - \frac{3}{2}b + \frac{5}{6}c$.

16. $\frac{1}{4}x + \frac{3}{2}y - \frac{2}{3}z$.

17. $-\frac{5}{2}a - \frac{10}{3}b + \frac{1}{2}c$.

18. $x - y + \frac{1}{5}z$.

19. $-\frac{4}{3}x - \frac{4}{3}z$.

20. $-\frac{5}{6}x + \frac{13}{6}y$.

3. 0.
5. $b - c$.
9. $3ax - 3by + 3cz$.
2. $-20ab + ca$.
5. $6x$.
3. $14ab - 11bc$.

IV. b. PAGE 22.

3. $a^2 + 3ab - 2b^2$.
6. $-2x^3 + x^2 + 4x + 2$.
9. $15x^3 - 4x^2 + 3x - 1$.
2. $x^3 + x^2 + x + 3$.
 $z - 3xz$.
7. $2a^2b$.

1. $7xy - 7yz + 18xz$.

2. $-12x^2y^2 + 8x^3y + 21xy^3$.

3. $-12 + 9ab + 6a^2b^2$.

4. $-2a^2bc + 6b^2ca + 5c^2ab$.

5. $-12a^2b + 15ab^2 - 5cd$.

6. $-16x^2y + 10xy^2 - 2x^2y^2$.

7. $20a^2b^2 + 16a^2b$.

8. $9x^2 - 9x + 9$.

9. $x^3 + 3x^2 + 5x + 7$.

10. $-17a^2x^2 + 13x^2 + 20$.

11. $2x^2 - 2x$.

12. $6x^2y + 2y^3$.

13. $a^3 - c^3 - abc$.

14. $3x^3 + 10x^2y - 10xy^2$.

15. $4x^4 - 5x^3 - 2x^2 - x + 2$.

16. $-4a^3 + 4b^3 - 2c^3 + 10abc$.

17. $-x^5 + 2x^4 + x^3 - x^2 + 2x - 2$.

18. $4a^5 - 7a^4 - 5a^3 + 9a^2 - a - 7$.

19. $-5a^2b - 14ab^2 + a^3b^3 + b^4$.

20. $-a^3 + 22a^2b - 16ab^2 + 2b^3$.

21. $2x^2 - \frac{4}{3}xy - \frac{1}{2}y^2$.

22. $\frac{4}{3}a^2 - \frac{7}{2}a - \frac{1}{2}$.

23. $-\frac{1}{6}x^2 - \frac{5}{6}x + \frac{7}{6}$.

24. $\frac{5}{8}x^2 + \frac{1}{6}ax - \frac{1}{3}$.

25. $\frac{3}{4}x^3 - \frac{1}{2}x^2y - \frac{1}{6}y^2$.

26. $-\frac{1}{8}a^3 - \frac{2}{3}a^2x - \frac{1}{2}ax^2$.

Miscellaneous Examples I. PAGE 23.

- | | |
|-------------------------------------|-----------------------|
| 1. (1) $2x + x^2$; (2) $-3a + b$. | 2. $2a + 2c$. |
| 3. (1) 21; (2) 108. | 4. (1) 11; (2) 18. |
| 5. $7x^2 - 10x^2$. | |
| 6. $8a^3 - 2a$. | 9. $2x^3 - 2x^2$. |
| 11. $2a - (3b + 5c)$. | |
| 12. 47; 12. | 13. $-y^2 + y$. |
| 15. 36. | 16. 0. |
| 17. a^2b . | |
| 18. $x + 2x$. | 20. $7xy$. |
| 21. 8. | 23. $4a$. |
| 24. 118. | |
| 25. 30 B.C. | 26. $2x^2 - 2x$. |
| 28. $a + b - (c - d)$. | |
| 29. $a + b$ miles south of O. | 30. $2x^2 + 7x - 3$. |

V. a. PAGE 27.

- | | | | |
|--|-------------------------------------|--------------------------|---------------------|
| 1. $35x^7$. | 2. $20a^{11}$. | 3. $56a^4b^3$. | 4. $30x^4y^2$. |
| 5. $8a^3b^6$. | 6. $6a^2bc^4$. | 7. $4a^6b^6$. | 8. $10a^3b$. |
| 9. $28a^7b^3$. | 10. $5a^4b^3x^2y^2$. | 11. $6a^2x^7y^3$. | 12. $abcxyz$. |
| 13. $15a^7b^3x^4$. | 14. $28a^3b^3x^5$. | 15. $40a^2cx^2$. | 16. $30a^2x^3y^3$. |
| 17. $2x^7y^8$. | 18. $3a^5x^9y^{16}$. | 19. $a^4b^3 + a^3b^2c$. | |
| 20. $20a^3b^2x^3 - 28a^2b^2x^4$. | 21. $10x^3 + 6x^2y$. | | |
| 22. $a^5b + a^3b^3 - a^3bc^2$. | 23. $ab^2c^2 + a^2b^2c - a^3b^2c$. | | |
| 24. $20a^4bc^3 + 12a^2b^3c^3 - 8a^2bc^5$. | 25. $15x^5y + 3x^4y^2 - 21x^3y^2$. | | |
| 26. $48x^6y^3 - 40x^4y^4 + 56x^3y^5$. | 27. $6a^5b^3c - 7a^3b^4c^2$. | | |

V. b. PAGE 30.

- | | | | | |
|----------|-----------|----------|------------|------------|
| 1. 36. | 2. -48. | 3. 5. | 4. 24. | 5. -16. |
| 6. -12. | 7. -9. | 8. -24. | 9. -168. | 10. 480. |
| 11. -16. | 12. 375. | 13. 500. | 14. 140. | 15. -2000. |
| 16. 500. | 17. -180. | 18. -56. | 19. -1000. | 20. -224. |
| 21. 40. | 22. -63. | 23. 118. | 24. -130. | 25. -54. |
| 26. 3. | 27. 1. | 28. 0. | 29. 29. | 30. -13. |

V. c. PAGE 31.

- | | | | |
|---|---------------------------------------|-------------------------------|------------------|
| 1. $-3a^2x^2$. | 2. $14a^3y^2x^2$. | 3. $-a^5b^3$. | 4. $-60x^2y^3$. |
| 5. $3a^3b^4c^5d^6$. | 6. $-5x^3y^4z^2$. | 7. $-36x^2y^2z - 48xy^2z^2$. | |
| 8. $a^3b^2c^3 - a^2b^2c^4$. | 9. $3x^2 + 3xy + 3xz$. | 10. $a^3bc - ab^3c + abc^3$. | |
| 11. $a^2b^2c - ab^2c^2 + a^2bc^2$. | | 12. $14a^4b^3 + 28a^3b^4$. | |
| 13. $15x^2y^2 - 18x^2y^3 + 24x^3y^3$. | | 14. $56x^6y^4 + 40x^4y^6$. | |
| 15. $-5x^2y^3z^2 + 3x^2y^2z^3 - 8x^3y^2z^2$. | 16. $-48x^3y^3z^5 + 96x^4y^2z^4$. | | |
| 17. $91x^4y^6 + 105x^5y^4$. | 17. $-8x^2y^2z^2 + 10x^4y^2z^4$. | | |
| 19. $-a^2b^2c^2 + a^3b^2c^2 + a^2b^3c^2$. | 20. $a^3b^2c - a^2b^3c + a^2b^2c^2$. | | |

PAGE 23.

3. $2a + 2c$.
 5. $7x^2 - 10x^2$.
 1. $2a - (3b + 5c)$.
 3. 0. 17. a^2b .
 5. 4a. 24. 118.
 7. $a + b - (c - d)$.
 9. $2x^2 + 7x - 3$.

21. $-3a^2 + \frac{9}{2}ab - 6ac$. 22. $-\frac{5}{2}x^2 + \frac{5}{3}xy + \frac{10}{3}x$.
 23. $\frac{1}{4}a^2x - \frac{1}{16}abx - \frac{3}{8}acx$. 24. $-2a^5x^3 + \frac{7}{2}a^4x^4$.
 25. $\frac{5}{2}a^4x^2 - \frac{5}{3}a^3x^3 + a^2x^4$. 26. $\frac{21}{2}x^3y - x^2y^2$.
 27. $\frac{1}{2}x^5y^2 - 3x^3y^4$. 28. $-x^2y^3 + \frac{16}{49}x^5y^6$.

V. d. PAGE 32.

4. $30x^4y^2$. 2. $x^2 - 25$. 3. $x^2 - 17x + 70$.
 8. $10a^3b$. 5. $x^2 - 3x - 70$. 6. $x^2 + 17x + 70$.
 12. $abcxyz$. 7. $x^2 - 36$. 8. $x^2 + 4x - 32$. 9. $x^2 - 13x + 12$.
 16. $30a^3x^6y^3$. 10. $x^2 + 11x - 12$. 11. $x^2 - 225$. 12. $-x^2 + 18x - 45$.
 13. $x^2 + 5x + 6$. 14. $-x^2 + 14x - 49$. 15. $x^2 - 25$.
 16. $x^2 + x - 182$. 17. $x^2 + x - 306$. 18. $x^2 - x - 380$.
 19. $x^2 - 256$. 20. $-x^2 + 42x - 441$. 21. $2x^2 + 13x - 24$.
 22. $2x^2 - 13x - 24$. 23. $2x^2 - 11x + 5$. 24. $2x^2 - 7x + 5$.
 25. $6x^2 + 11x - 35$. 26. $6x^2 - 11x - 35$. 27. $10x^2 + 3x - 18$.
 28. $10x^2 - 3x - 18$. 29. $9x^2 - 25y^2$. 30. $9x^2 - 30xy + 25y^2$.
 31. $a^2 + ab - 6b^2$. 32. $a^2 + ab - 56b^2$. 33. $3a^2 - 30ab + 48b^2$.
 34. $a^2 - 4ab - 45b^2$. 35. $x^2 + ax - bx - ab$. 36. $x^2 - ax + bx - ab$.
 37. $x^2 - 2ax + 3bx - 6ab$. 38. $a^2x^2 - b^2y^2$.
 39. $x^2y^2 - a^2b^2$. 40. $4p^2q^3 - 9r^2$.

4. 5. -16.
 168. 10. 480.
 40. 15. -2000.
 1000. 20. -224.
 130. 25. -54.
 9. 30. -13.

V. e. PAGE 34.

1. $a^2 + 2ab + b^2 - c^2$. 2. $a^2 - 4b^2 + 4bc - c^2$.
 3. $a^4 + a^2b^2 + b^4$. 4. $x^2 + 4x^2y + 3xy^2 + 12y^3$.
 5. $x^4 - 4x^2 + 8x + 16$. 6. $x^6 + y^6$. 7. $x^3 - y^3$.
 8. $a^4 + 4a^2x^2 + 16x^4$. 9. $64a^3 - 27b^3$. 10. $x - a^4$.
 11. $x^4 + 2x^3 - 7x^2 - 8x + 12$. 12. $4x^5 - x^3 + 4x$.
 13. $a^6 + a^3b^3$. 14. $x^5 - 2x^4 - 4x^3 + 19x^2 - 31x + 15$.
 15. $a^5 + 4ab^4$. 16. $8x^5 - 27y^3$.
 17. $-x^4 + 4x^3y - x^2y^2 - 4xy^3 - y^4$. 18. $a^6 - a^4b^4 + 2a^3b^3 + b^6$.
 19. $x^4 - 2x^2y^2 + y^4$. 20. $a^3b^2 + c^2a^2 - a^2c^2 - b^2c^2$.
 21. $75a^5b^3 - 28a^3b^5 + 13a^2b^6 - 12ab^7$. 22. $81x^4 - 256a^4$.
 23. $a^4 - 25a^2b^2 - 10ab^3 - b^4$. 24. $x^3 + 3xy + y^3 - 1$.
 25. $a^3 + b^3 + c^3 - 3abc$. 26. $x^5 + y^5$. 27. $x^{15} + y^{10}$.

28. $a^6 - 2a^3 + 1$.
 31. $\frac{1}{4}a^3 + \frac{1}{72}a - \frac{1}{12}$.
 33. $\frac{2}{9}x^3 - \frac{3}{4}y^3$.
 35. $\frac{1}{4}x^4 - \frac{43}{36}x^2 + \frac{9}{16}$.
29. $a^2x^3 + 27a^2y^6$.
 32. $\frac{1}{4}x^3 - \frac{5}{6}x^2 + \frac{1}{12}x + \frac{1}{2}$.
 34. $\frac{9}{8}x^4 - \frac{3}{2}ax^3 + \frac{1}{2}a^2x^2 - \frac{2}{9}a^4$.
 36. $\frac{1}{4}a^4 + x^4$.

V. f. PAGE 36.

- | | | |
|---------------------------|----------------------------|----------------------------|
| 1. $x^2 + 3x - 40$. | 2. $x^3 + 5x - 6$. | 3. $x^3 + 7x - 30$. |
| 4. $x^2 + 4x - 5$. | 5. $x^2 - 2x - 63$. | 6. $x^2 - 18x + 80$. |
| 7. $x^2 + 7x - 44$. | 8. $x^2 + 2x - 8$. | 9. $x^2 - 4$. |
| 10. $a^3 - 1$. | 11. $a^2 + 4a - 45$. | 12. $a^2 + 9a - 36$. |
| 13. $a^3 - 4a - 32$. | 14. $a^2 - 64$. | 15. $a^2 + 7a - 78$. |
| 16. $a^2 + 6a + 9$. | 17. $a^2 - 121$. | 18. $a^2 - 16a + 64$. |
| 19. $x^2 - ax - 6a^2$. | 20. $x^3 + ax - 30a^2$. | 21. $x^3 - 9a^2$. |
| 22. $x^2 + 2xy - 8y^2$. | 23. $x^2 - 49y^2$. | 24. $x^2 - 6xy + 9y^2$. |
| 25. $a^3 + 6ab + 9b^2$. | 26. $a^2 + 5ab - 50b^2$. | 27. $a^2 - 17ab + 72b^2$. |
| 28. $2x^2 - x - 10$. | 29. $2x^2 - 9x + 10$. | 30. $2x^2 - 3x - 9$. |
| 31. $3x^2 + 2x - 1$. | 32. $4x^2 + 8x - 5$. | 33. $6x^2 + 5x - 21$. |
| 34. $8x^2 - 6x - 9$. | 35. $9x^2 - 64$. | 36. $4x^2 - 20x + 25$. |
| 37. $9x^2 - 3xy - 2y^2$. | 38. $9x^2 + 12xy + 4y^2$. | 39. $4x^2 + 4xy - 35y^2$. |
| 40. $25x^2 - 9a^2$. | 41. $2x^2 + 5ax - 25a^2$. | 42. $4x^2 + 4ax + a^2$. |

VI. a. PAGE 40.

- | | | | |
|---------------------------|--|---|--------------------------------|
| 1. $3x$. | 2. $-3x$. | 3. $-5x^3$. | 4. $-bx$. |
| 5. xy^2 . | 6. $-a^2$. | 7. $4ac$. | 8. $-4a^2b^4c^5$. |
| 9. a^4c^6 . | 10. $3x^3y^5z^2$. | 11. $4x^2$. | 12. $6a^6$. |
| 13. $5a^4$. | 14. $7a^2b^3$. | 15. -1 . | 16. $-7ab^2$. |
| 17. $-8b^2x$. | 18. $10y^2$. | 19. $x - 2y$. | 20. $x^2 - 3x + 1$. |
| 21. $x^4 - 7x^3 + 4x^2$. | 22. $10x^4 - 8x^3 + 3x$. | 23. $-3x^2 + 5x$. | 24. $3x - 4$. |
| 25. $3x^3 + 4x$. | 26. $2x^2y - 3xy^2$. | 27. $-a + b + c$. | 28. $a - b - b^2$. |
| 29. $-x^2 + 3xy + 4y^2$. | 30. $-2x^3y^3 + 4x^2y - 3y^2$. | 31. $2a - 3b + 4c$. | 32. $-\frac{1}{3}x^2 + 2y^2$. |
| 33. $3x - 2y - 4$. | 34. $-\frac{6}{7}a^2x^2 + \frac{3}{2}ax^3$. | 35. $\frac{2}{3}a - \frac{1}{6}b - c$. | |

$$10. x^6 + 2x^3y^3 + y^6.$$

$$^2 + \frac{1}{12}x + \frac{1}{2}.$$

$$x^3 + \frac{1}{2}a^2x^2 - \frac{2}{9}a^4.$$

$$3. x^2 + 7x - 30.$$

$$6. x^2 - 18x + 80.$$

$$9. x^2 - 4.$$

$$2. a^2 + 9a - 36.$$

$$5. a^2 + 7a - 78.$$

$$8. a^2 - 16a + 64.$$

$$1. x^2 - 9a^2.$$

$$4. x^2 - 6xy + 9y^2.$$

$$7. a^2 - 17ab + 72b^2.$$

$$10. 2x^2 - 3x - 9.$$

$$3. 6x^2 + 5x - 21.$$

$$6. 4x^2 - 20x + 25.$$

$$9. 4x^2 + 4xy - 35y^2.$$

$$2. 4x^2 + 4ax + a^2.$$

$$4. -bx.$$

$$8. -4a^2b^4c^5.$$

$$12. 6a^6.$$

$$16. -7ab^3.$$

$$20. x^2 - 3x + 1.$$

$$^3 + 3x.$$

$$26. 2x^2y - 3xy^2.$$

$$y + 4y^2.$$

$$4c.$$

$$4.$$

$$c.$$

VI. b. PAGE 42.

$$1. x + 2.$$

$$2. x - 4.$$

$$3. a - 6.$$

$$4. a - 24.$$

$$5. 3x + 1.$$

$$6. x + 5.$$

$$7. 5x + 1.$$

$$8. x + 7.$$

$$9. 5x + 1.$$

$$10. x + 11.$$

$$11. x + 5.$$

$$12. 3x + 1.$$

$$13. 3x + 7.$$

$$14. 3x - 7.$$

$$15. 3x - 5.$$

$$16. 4x - 7.$$

$$17. 4a + 3x.$$

$$18. 5a - x.$$

$$19. 3a + 4c.$$

$$20. 3a - 5c.$$

$$21. 6x + 5y.$$

$$22. 8x + 3y.$$

$$23. x^2 + 14x.$$

$$24. 4x^2 - x.$$

$$25. 9x^2 + 9x + 5.$$

$$26. 2a^2 - 5a + 3.$$

$$27. 3 + 3a + a^2.$$

$$28. 8 - 36x + 54x^2 - 27x^3.$$

VI. c. PAGE 44.

$$1. x - 4.$$

$$2. y + 1.$$

$$3. 2m - 3.$$

$$4. 2a^2 - 3a.$$

$$5. x^2 - x + 1.$$

$$6. a^2 - 3a + 1; \text{ rem. } a - 6.$$

$$7. a^2 + 3a + 2.$$

$$8. 2x^2 + x - 1; \text{ rem. } 3x + 4.$$

$$9. x^3 - 2x^2 + x + 1.$$

$$10. x^3 - 3x^2 + 2x - 1.$$

$$11. 10x^2 - 3x - 12; \text{ rem. } 7x - 45.$$

$$12. 7y^2 + 5y - 3; \text{ rem. } -39y + 27.$$

$$13. 2k^2 - 5k + 2.$$

$$14. 5 - 7m - m^3.$$

$$15. x^2 + 5x + 6.$$

$$16. x^2 - 2x + 3; \text{ rem. } 31x - 15.$$

$$17. 12 + 8x + x^2.$$

$$18. 7x^2 + 5xy + 2y^2.$$

$$19. x^2 - xy + y^2; \text{ rem. } x^2.$$

$$20. x^3 + x - y.$$

$$21. a^6 + a^3b^3 + b^6.$$

$$22. x^7 - x^6y + x^4y^3 - x^3y^4 + xy^6 - y^7.$$

$$23. x^6 + 2x^5y^2 - 3x^4y^4 - 6x^3y^6 + 2x^2y^8 + 4xy^{10} + y^{12}.$$

$$24. a^2 + 2ab + b^2 + a + b + 1.$$

$$25. x^5 - x^4y + xy^4 - y^5.$$

$$26. a^{10} + a^8b^2 + a^6b^4 + a^4b^6 + a^2b^8 + b^{10}.$$

$$27. a^8 - 2a^6b^2 + 3a^4b^4 - 2a^2b^6 + b^8.$$

$$28. 1 + c + a^2 + 2x - 2ax + 4x^2.$$

$$29. \frac{1}{4}a^2 - 3ax + 9x^2.$$

$$30. \frac{1}{9}a^2 - \frac{1}{6}a + \frac{1}{16}.$$

$$31. \frac{6}{25}a^3 - \frac{3}{5}a^2c + \frac{3}{2}ac^2.$$

$$32. \frac{3}{8}a^2 - \frac{1}{4}a - \frac{2}{3}.$$

$$33. 6x - \frac{1}{3}y - \frac{1}{2}.$$

$$34. \frac{4}{9}a^4 + \frac{1}{2}a^3x + \frac{9}{16}a^2x^2 + \frac{81}{128}ax^3.$$

VII. a. PAGE 47.

$$1. a + b - c.$$

$$2. a.$$

$$3. a + 3b - 4c.$$

$$4. 3a - b - c.$$

$$5. -2a - 4b - 2c.$$

$$6. -a + b - c.$$

$$7. b - a.$$

$$8. x - y.$$

$$9. 2a - 2b.$$

$$10. -2x - 5y.$$

$$11. x - a.$$

$$12. 2a - b - d.$$

$$13. -3c + 4y.$$

$$14. -x + 2y + 6z.$$

$$15. -5x.$$

$$16. -25x + 2y.$$

$$17. 11x - 36y.$$

$$18. 2x - 2z.$$

$$19. 2a.$$

$$20. a.$$

VII. b. PAGE 48.

- | | | |
|--------------------|---------------------------------------|--------------------------|
| 1. $3a$. | 2. a . | 3. $6a+2b-2c-2d$. |
| 4. $2x-3y+12z$. | | 5. b . |
| 7. $2b+4c$. | 8. $-a^2+8b^2-9c^2$. | 9. $-2a+6b+2c-2d$. |
| 10. $4a+b+c$. | 11. $-50c$. | 12. $-11a-2b$. |
| 13. $-a+b+5c$. | 14. $-2a+10b-11c$. | 15. $-227a+216b+84$. |
| 16. $2a-12c+84d$. | 17. $3a+4x$. | 18. $-10a$. |
| 19. $4a$. | 20. 0 . | 21. $\frac{11}{5}a-2b$. |
| 22. $12x-30y$. | 23. $a-\frac{13}{3}b+\frac{10}{3}c$. | 24. 0 . |

VII. c. PAGE 50.

1. $(a+2)x^4+(b-5)x^2+(2b-3)x+5$.
2. $(5a-b)x^3+(3b-4)x^2+(c-2)x+ab-7$.
3. $(9a-7)x^3+(5a-3)x^2+(7-2c)x+2$.
4. $(2c-a^2)x^3+(1-3b)x^2+(4d-3ab)x$.
5. $-(a^2+b)x^4-(2b-5)x^3-(3-a)x^2$.
6. $-(ab-7)x^3-(abc-7)x^2-(3c^2-5a)x$.
7. $-(c-a^2)x^3-(b+5-a)x^2$.
8. $-(a+c+7-3b^2)x^4-(b+5c^2)x$.
9. $(a-b)x^3-(b+2c)x^2-(b+c+d)x$.
10. $(5a+4c)x^3+(3a-6b+7c)x^2+(2a-7b)x$.
11. $(3a+2c)x^3+(a+8b)x^2-(8a+9b)x$.
12. $(6b+1)x^5-(a+2b)x^4-(2a+3c)x$.
13. $(a+b)x^3-(a+b)x^2+(a-b)x$.

VII. d. PAGE 51.

1. $(a-c)x^3+(b+c)x^2-(2c+1)x$.
2. $(1-b)x^3+(a+1)x^2+(b-1)x-1$.
3. $(a^2-5a+2)x^3+(2a-b)x^2-(a+5)x$.
4. $(a-p+1)x^3+(b+q+2)x-c-r+3$.
5. $(p+q-1)x^3+(p+q)x^2-(p+q)x+q$.
6. $acx^3+(2a+bc)x^2+(2b+c)x+2$.
7. $acx^3-(2a+bc)x^2+(3a+2b)x-3b$.
8. $apx^3+(aq-bp)x^2-(bq+cp)x-cq$.
9. $2bx^3-(3b-2c)x^2-(b+3c)x-c$.
10. $ax^3-(a+2b)x^2+(2b+3c)x-3c$.

$$-2c-2d.$$

$$6. 21a+b.$$

$$-2a+6b+2c-2d.$$

$$-11a-2b.$$

$$-227a+216b+84.$$

$$-10a.$$

$$\frac{11}{5}a-2b.$$

$$0.$$

$$11. apx^3 - (2a+3p)x^2 + (6-aq)x + 3q.$$

$$12. x^6 - (a^2+2b)x^4 + (2ac+b^2)x^2 - c^2.$$

$$13. a^2x^6 + (6a-1)x^4 + (9-2b)x^2 - b^2.$$

$$14. x^8 - (a^2+2b)x^6 + (2ac+b^2+2d)x^4 - (2bd+c^2)x^2 + d^2.$$

VIII. a. PAGE 55.

1. 5.	2. 4.	3. 7.	4. 4.	5. 3.	6. 1.
7. 5.	8. 3.	9. 15.	10. 13.	11. 13.	12. 5.
13. 1.	14. 16.	15. 10.	16. 30.	17. 5.	18. 1.
19. 2.	20. 1.	21. 1.	22. 2.	23. 3.	24. 1.
25. 4.	26. 3.	27. 3.	28. 3.	29. 1.	30. 4.
31. 7.	32. 3.	33. 4.	34. 4.	35. 1.	36. 1.
37. 2.	38. 2.	39. 1.	40. 2.		

VIII. b. PAGE 58.

1. 20.	2. 15.	3. 8.	4. 16.	5. 25.	6. 17.
7. 13.	8. 10.	9. 7.	10. 4.	11. $-\frac{1}{7}$.	12. $\frac{1}{7}$.
13. 5.	14. 7.	15. 6.	16. 10.	17. 6.	18. 8.
19. 7.	20. 25.	21. $3\frac{1}{7}$.	22. 8.	23. 12.	24. 5.
25. 5.	26. 12.	27. $\frac{4}{7}$.	28. $-5\frac{1}{8}$.	29. 8.	30. $66\frac{2}{3}$.
31. 7.	32. 7.	33. 2.	34. 12.	35. 27.	36. 5.

VIII. c. PAGE 60.

1. $2\frac{2}{3}$.	2. 6.	3. 10.	4. -6.	5. $9\frac{2}{3}$.	6. $1\frac{1}{3}$.
7. -12.	8. $\frac{3}{8}$.	9. $14\frac{4}{5}$.	10. $-\frac{3}{4}$.	11. $\frac{4}{5}$.	12. $-\frac{2}{21}$.
13. $1\frac{1}{2}$.	14. $-\frac{2}{3}$.	15. $1\frac{3}{4}$.	16. 12.	17. $3\frac{6}{7}$.	18. $2\frac{1}{4}$.
19. $\frac{5}{7}$.	20. $1\frac{3}{8}$.	21. $\frac{3}{7}$.			

IX. a. PAGE 62.

1. $y-x$.	2. $\frac{a}{3}$.	3. $5b$.	4. $3d-2c$.
5. $2k$.	6. $100-x$.	7. $\frac{b}{a}$.	8. $20-c$.

- | | | | |
|------------------------|----------------------------|--------------------|--------------------------|
| 9. $\frac{5}{6}a$. | 10. $\frac{600}{x}$. | 11. $x+11$. | 12. $c-20$. |
| 13. $90-x$. | 14. $x-30$. | 15. 20. | 16. $2x$. |
| 17. $36-x$. | 18. $x+a$. | 19. $5x$ days. | 20. 4. |
| 21. $\frac{x}{2}$. | 22. $\frac{x}{4}$. | 23. xy miles. | 24. $\frac{y}{x}$ miles. |
| 25. $\frac{60x}{a}$. | 26. $\frac{120}{x}$ hours. | 27. $5p$. | 28. $\frac{44}{x}$. |
| 29. $5a+2b$. | 30. $400-x$. | 31. $240a+12b-c$. | |
| 32. $x-6$. | 33. b . | 34. $40x$. | 35. $20a+2b-c$. |
| 36. $\frac{100}{xy}$. | 37. $y-\frac{13x}{5}$. | 38. $100-x-y-z$. | |
| 39. $240x+12y+z-30$. | 40. $2y+2z-x$. | | |

IX. b. PAGE 65.

- | | |
|-----------------------------------|-------------------------------|
| 1. $x, x+1, x+2, x+3$. | 2. $y-2, y-1, y$. |
| 3. $x-2, x-1, x, x+1, x+2$. | 4. $2n+2$. |
| 5. $2x-1$. | |
| 6. $6n+3$. | 7. $x-a-b$ miles. |
| 8. $n(a+b)$. | 9. $x+y+5$. |
| 10. $2x+5$. | 11. $mx+y$. |
| 12. $6x$. | 13. $\text{£}10bc$. |
| 14. $\text{£}\frac{ax}{20}$. | 15. $\text{£}\frac{a^3}{2}$. |
| 16. $\text{£}\frac{x^2y^2}{50}$. | 17. $3xy$. |
| 18. $\frac{x^3}{9}$. | 19. $\frac{px}{2}$. |
| 20. $\frac{abc}{60}$. | 21. $\frac{4yz}{x}$. |
| 22. $ab-\frac{c^2}{9}$. | 23. $\frac{3a}{4x}$. |
| 24. $\frac{bc}{20}$ hours. | 25. $\frac{22a}{15b}$. |
| 26. $\frac{15xy}{22}$. | 27. $\frac{y}{xz}$ days. |
| 28. yz . | 29. $\frac{y}{10r}$. |
| 30. $\frac{100p}{ar}$. | |

X. a. PAGE 71.

- | | | | |
|--|---|-------------|--------------------|
| 1. 17, 12. | 2. 13, 5. | 3. 75. | 4. 20 miles. |
| 5. 15, 43. | 6. 162. | 7. 1. | 8. 50, 55. |
| 9. 27, 28, 29. | 10. 3, 5. | 11. 15, 5. | 12. $\text{£}20$. |
| 13. 5. | 14. 60, 61. | 15. 6, 3. | |
| 16. $A \text{ £}100, B \text{ £}130, C \text{ £}150$. | 17. 53 florins, 71 shillings. | | |
| 18. Silk 6s., Linen 1s. | 19. 48, 12. | 20. 65, 40. | |
| 21. 60, 10. | 22. 20 half-crowns, 5 crowns, 10 shillings. | | |
| 23. 25, 5. | 24. 123 runs, 10 byes, 5 wides. | | |
| 25. 15 ft., 12 ft. | 26. 18 ft., 10 ft. | | |

12. $c - 20$.16. $2x$.

20. 4.

24. $\frac{y}{x}$ miles.28. $\frac{44}{x}$.

240a + 12b - c.

35. $20a + 2b - c$.38. $100 - x - y - z$.

+ 2z - x.

- 2, $y - 1$, y .5. $2x - 1$.9. $x + y + 5$.

13. £10bc.

17. $3xy$.21. $\frac{4yz}{x}$.25. $\frac{22a}{15b}$.30. $\frac{100p}{ar}$.

4. 20 miles.

8. 50, 55.

12. £20.

15. 6, 3.

s, 71 shillings.

20. 65, 40.

crown, 10 shillings.

6 wises.

0 ft.

X. b. PAGE 73.

- | | | | |
|-----------------------------|-----------------------------|--------------|------------|
| 1. 54. | 2. 24. | 3. 60. | 4. 35. |
| 5. 75. | 6. 24, 25. | 7. 224, 252. | 8. 49, 50. |
| 9. 50, 51, 52. | 10. £33. | 11. 27. | |
| 12. 90 Port, 150 Claret. | 13. A £450, B £180, C £140. | | |
| 14. A £525, B £600, C £160. | 15. £49. | | |
| 16. 12 ft. 18 ft. | 17. £12000. | 18. 44. | |

XI. a. PAGE 74.

- | | | | |
|--------------------|--------------------|-------------------|-----------------|
| 1. $2ab$. | 2. x^2y^2 . | 3. $2xy^2z$. | 4. abc . |
| 5. $5ab$. | 6. $3xy^2z$. | 7. $2a^2b^2c^2$. | 8. $7ab^2c^3$. |
| 9. $3x^2yz^2$. | 10. $2ax$. | 11. $7a$. | 12. $17abc$. |
| 13. xy . | 14. $8a^2b^2c^2$. | 15. $25xy$. | 16. bx . |
| 17. $5a^3b^3c^3$. | 18. abc . | | |

XI. b. PAGE 75.

In examples 19-20 the H.C.F. stands first; the L.C.M. second.

- | | | | |
|------------------------------|-------------------------------|--------------------------------------|----------------------|
| 1. $2a^2bc$. | 2. x^3y^2z . | 3. $12x^2y^2z$. | 4. $20a^2b^2c^3$. |
| 5. $15a^4b^3c^5$. | 6. $24abxyz$. | 7. abc . | 8. $a^2b^2c^2$. |
| 9. $12abc$. | 10. $12xyz$. | 11. $12x^2y^2z^2$. | 12. $42a^2b^3$. |
| 13. $a^2b^2c^2$. | 14. $30a^2b^2c^2$. | 15. $12x^3y^4$. | 16. $56x^4y^5$. |
| 17. $210a^3b^3c^3$. | 18. $264a^4b^4c^4$. | 19. ac , $12abc$. | 20. $2y$, $12xyz$. |
| 21. bc , $9ab^2c$. | 22. $13a^2bc$, $39a^3bc^2$. | 23. $17xy$, $51x^2yz^2$. | |
| 24. $5xy^2z$, $75x^3yz^2$. | 25. b , $30abc$. | 26. $17m^2p^2$, $51m^4n^4p^4$. | |
| 27. y^2 , $x^3y^5z^4$. | 28. $5p^2$, $60m^2p^3q^4$. | 29. $36k^2m^2n^4$, $216k^3m^3n^5$. | |

XII. a. PAGE 76.

- | | | | |
|----------------------------|------------------------------|----------------------------|--------------------------|
| 1. $\frac{1}{2b}$. | 2. $\frac{a}{4b}$. | 3. $\frac{2y}{5x}$. | 4. $\frac{1}{5ab}$. |
| 5. $\frac{z^2}{xy}$. | 6. $\frac{3a}{5c}$. | 7. $\frac{3x^2}{4z^2}$. | 8. $\frac{2a^2}{3bc}$. |
| 9. $\frac{4n}{5mp}$. | 10. $\frac{5m^2p^2}{6n^3}$. | 11. $\frac{c}{a^2b}$. | 12. $\frac{3xz}{5y^3}$. |
| 13. $\frac{yz^3}{2x}$. | 14. $\frac{a^2c^3}{3b^2}$. | 15. $\frac{nq}{mp^3}$. | 16. $\frac{2np^3}{3m}$. |
| 17. $\frac{3x^2}{5ay^4}$. | 18. $\frac{3}{4abc}$. | 19. $\frac{2p^2m^2}{3k}$. | 20. $\frac{2xyz}{3}$. |

XII. b. PAGE 77.

- | | | | |
|-----------------------------|-------------------------|------------------------------|---------------------------|
| 1. $\frac{2cd^2}{3b}$ | 2. $\frac{a^2}{bc}$ | 3. $\frac{9ax^2z^2}{bc}$ | 4. $\frac{14b^3}{15c^3y}$ |
| 5. $\frac{3mnz^2}{2x}$ | 6. $\frac{9mnp}{4k}$ | 7. $\frac{x}{2a^2}$ | 8. $\frac{400x}{441y^3}$ |
| 9. $\frac{y^3z^2}{nx^4}$ | 10. 3. | 11. $\frac{7b}{4a}$ | 12. $\frac{d^2}{4a^2c}$ |
| 13. $\frac{7acy}{8bdx}$ | 14. $\frac{6x^2yz}{5a}$ | 15. $\frac{9b^2cz^2}{4x^3y}$ | 16. 8. |
| 17. $\frac{p^2q^2y}{10x^2}$ | 18. y^2 | | |

XII. c. PAGE 78.

- | | | |
|-------------------------------|-------------------------------|----------------------------------|
| 1. $\frac{4x, y}{2a}$ | 2. $\frac{4x^3, 3y^2}{3x^2y}$ | 3. $\frac{ac, 2b^2}{2bc}$ |
| 4. $\frac{ad, bc, 2bd}{bd}$ | 5. $\frac{6ac, b^2}{3bc}$ | 6. $\frac{5m, 4p}{20n}$ |
| 7. $\frac{3k, 2p}{6x}$ | 8. $\frac{2m, n}{6x}$ | 9. $\frac{a^2, b^2}{abc}$ |
| 10. $\frac{ax, b}{x^2}$ | 11. $\frac{2y, 3x}{xy}$ | 12. $\frac{x^2, y^2, 3x^2y}{xy}$ |
| 13. $\frac{4x^2, 9y^2}{6xy}$ | 14. $\frac{8ac, 3ab}{10bc}$ | 15. $\frac{9ac, 5b^2}{21bc}$ |
| 16. $\frac{18, 3ab, a^2}{9a}$ | | |

XII. d. PAGE 79.

- | | | | |
|-------------------------------|-----------------------------|-----------------------------|-----------------------------|
| 1. $\frac{5x}{6}$ | 2. $\frac{y}{20}$ | 3. $\frac{a}{12}$ | 4. $\frac{2x^2 - 15}{3x}$ |
| 5. $\frac{5x + 2y}{10}$ | 6. $\frac{3a - 2b}{12}$ | 7. $\frac{3m - 2n}{24}$ | 8. $\frac{2m - 3n}{15}$ |
| 9. $\frac{3x - y}{21}$ | 10. $\frac{3a + b}{39}$ | 11. $\frac{3p - q}{48}$ | 12. $\frac{15m - n}{36}$ |
| 13. $\frac{22x}{15}$ | 14. $\frac{9x}{20}$ | 15. $\frac{x}{4}$ | 16. $\frac{6a - 4b}{15}$ |
| 17. $\frac{11a}{30}$ | 18. $\frac{5x}{24}$ | 19. $\frac{7x}{18}$ | 20. $\frac{5x}{4}$ |
| 21. $\frac{31x}{36}$ | 22. $\frac{17x}{24}$ | 23. $\frac{bx - ay}{ab}$ | 24. $\frac{9bx + 2ay}{3ab}$ |
| 25. $\frac{ac + b}{c}$ | 26. $\frac{xz - y}{z}$ | 27. $\frac{a^2 - 3b^2}{3a}$ | 28. $\frac{a^3 + b^3}{a}$ |
| 29. $\frac{x^3 - 2y^3}{2x^2}$ | 30. $\frac{p^5 - k^5}{p^2}$ | | |

Miscellaneous Examples II. PAGE 80.

4. $\frac{14b^3}{15c^2y}$
 8. $\frac{400x}{441y^3}$
 $\frac{7^2}{c^2}$ 13. $\frac{7acy}{8bdx}$
 $\frac{q^2y}{0x^2}$ 18. y^2
- $\frac{ac, 2b^2}{2bc}$
 $\frac{5m, 4p}{20n}$
 10. $\frac{ax, b}{x^2}$
 $\frac{4x^2, 9y^2}{6xy}$
 18. $\frac{3ab, a^2}{9a}$
1. $3x^2 + 7x - 8$. 2. $13x$. 3. 20.
 4. $a^2 + b^2 + c^2$. 5. $x^5 - 11x - 10$. 6. $(1) \frac{1}{2}$; (2) -3.
 7. $x^2 + 2x - 3$. 8. $-4a + 5b$. 9. $5x$.
 10. $4x^2 - 6x - 1$. 11. $(1) x^2 + 14x - 51$; (2) $24x^2 - 55x - 24$.
 12. $(1) \frac{1}{4}$; (2) 1. 13. $-ab$. 14. $\frac{8}{5}$.
 15. $16a^2 + 2ab$. 16. $x^5 + 4x^4 + 48x - 32$.
 17. $29a$. 18. $(1) -2$; (2) 41. 19. $3p^3 - 5p^2 + 2p$.
 20. $6a + 2c - 2d$. 21. $2x^3 - x^2 - x$. 22. 1.
 23. 1935. 24. 4. 25. $4m - 5n$.
 26. $3x - 9$. 27. 0. 28. $(1) -15$; (2) 4.
 29. $3y^3 - 9y^2 + 2y - 1$. 30. $A \text{ £}800, B \text{ £}320$.
 31. 14. 32. $6m^4 - 96$. 33. $x - 2$.
 34. $ap + bq$ miles; $\frac{ap + bq}{c}$ hours. Numerically, 55 miles; 5 hours.
 35. $(1) \frac{1}{7}$; (2) $7\frac{1}{13}$. 36. 4320.

XIII. a. PAGE 86.

1. $x=2, y=1$. 2. $x=3, y=5$. 3. $x=2, y=3$.
 4. $x=4, y=-1$. 5. $x=1, y=2$. 6. $x=3, y=4$.
 7. $x=5, y=6$. 8. $x=1, y=2$. 9. $x=3, y=1$.
 10. $x=2, y=1$. 11. $x=1, y=3$. 12. $x=1, y=1$.
 13. $x=7, y=5$. 14. $x=10, y=3$. 15. $x=5, y=12$.
 16. $x=7, y=8$. 17. $x=6, y=8$. 18. $x=5, y=8$.
 19. $x=-7, y=-3$. 20. $x=17, y=-19$. 21. $x=1, y=2$.

XIII. b. PAGE 87.

1. $x=12, y=8$. 2. $x=10, y=6$. 3. $x=18, y=12$.
 4. $x=20, y=15$. 5. $x=45, y=35$. 6. $x=51, y=17$.
 7. $x=20, y=60$. 8. $x=14, y=15$. 9. $x=-2, y=4$.
 10. $x=3, y=5$. 11. $x=7, y=3$. 12. $x=5, y=4$.
 13. $x=3, y=-4$. 14. $x=19, y=3$. 15. $x=12, y=-4$.
 16. $x=13, y=7$.

XIII. c. PAGE 90.

- | | |
|------------------------|--|
| 1. $x=1, y=2, z=3.$ | 2. $x=-2, y=4, z=1.$ |
| 3. $x=2, y=3, z=1.$ | 4. $x=1, y=2, z=3.$ |
| 5. $x=9, y=2, z=-4.$ | 6. $x=3, y=2, z=1.$ |
| 7. $x=5, y=6, z=7.$ | 8. $x=1, y=2, z=3.$ |
| 9. $x=2, y=-2, z=5.$ | 10. $x=4, y=-3, z=2.$ |
| 11. $x=8, y=10, z=14.$ | 12. $x=3, y=9, z=15.$ |
| 13. $x=6, y=8, z=5.$ | 14. $x=\frac{3}{2}, y=\frac{2}{3}, z=\frac{5}{6}.$ |
| 15. $x=6, y=2, z=1.$ | 16. $x=35, y=30, z=25.$ |

XIII. d. PAGE 92.

- | | | |
|--|--|-------------------------------------|
| 1. $x=5, y=3.$ | 2. $x=2, y=7.$ | 3. $x=3, y=2.$ |
| 4. $x=\frac{1}{3}, y=\frac{1}{4}.$ | 5. $x=7, y=6.$ | 6. $x=\frac{1}{3}, y=\frac{1}{5}.$ |
| 7. $x=2, y=-3.$ | 8. $x=-5, y=4.$ | 9. $x=\frac{2}{3}, y=\frac{3}{4}.$ |
| 10. $x=9, y=25.$ | 11. $x=\frac{1}{4}, y=\frac{1}{3}.$ | 12. $x=\frac{1}{5}, y=\frac{1}{6}.$ |
| 13. $x=\frac{1}{2}, y=\frac{1}{3}, z=\frac{1}{4}.$ | 14. $x=\frac{1}{8}, y=\frac{1}{12}, z=\frac{1}{16}.$ | |
| 15. $x=3, y=-2, z=1.$ | | |

XIV. PAGE 95.

- | | | | |
|---------------------------------|---|------------------------|------------------------------|
| 1. 22, 12. | 2. 55, 18. | 3. 25, 17. | 4. 53, 23. |
| 5. 23, 17. | 6. Tea 3s. 4d., Sugar 4d. | 7. Horse £23, Cow £16. | |
| 8. A £140, B £60, C £70, D £20. | 9. A £99, B 115, C £33, D £23. | | |
| 10. A 36 years, B 14 years. | 11. A 55 years, B 21 years. | | |
| 12. A 5 miles, B 4 miles. | 13. C $3\frac{1}{2}$ miles, D $4\frac{1}{4}$ miles. | | |
| 14. $\frac{13}{25}$ | 15. $\frac{15}{26}$ | 16. $\frac{2}{15}$ | 17. $\frac{3}{14}$ |
| 18. 28, 82. | 19. 85, 58. | 20. 27. | 21. 72. |
| 22. £5. 11s. | 23. 8 white, 12 black. | 24. 860. | 25. Man 2s. 6d., Boy 1s. 6d. |
| 26. 20 lbs., 40 lbs. | 27. 15 miles. | 28. 8 hours. | |
| 29. 6 miles, 3 miles an hour. | 30. 10s., 1s. 6d. | | |
| 31. £500. | 32. 3 miles, $4\frac{2}{7}$ miles an hour. | | |

XV. a. PAGE 99.

$$\begin{aligned} &= 4, z = 1. \\ &= 2, z = 3. \\ &= 2, z = 1. \\ &= 2, z = 3. \\ &= -3, z = 2. \\ &= 9, z = 15. \\ &= \frac{2}{3}, z = \frac{5}{6}. \\ &= 30, z = 25. \end{aligned}$$

$$\begin{aligned} 3. \quad &x=3, y=2. \\ 6. \quad &x=\frac{1}{3}, y=\frac{1}{5}. \\ 9. \quad &x=\frac{2}{3}, y=\frac{3}{4}. \\ 2. \quad &x=\frac{1}{5}, y=\frac{1}{6}. \\ 2. \quad &z=\frac{1}{16}. \end{aligned}$$

4. 53, 23.
orse £23, Cow £16
115, C £33, D £23.
rs, B 21 years.
es, D $4\frac{1}{4}$ miles.

18. 28, 82.

22. £5. 11s.

ed., Boy 1s. 6d.

ours.

3d.

$\frac{2}{7}$ miles an hour.

$$\begin{aligned} 1. \quad &9a^2b^6. & 2. \quad &a^6c^2. & 3. \quad &49a^2b^4. & 4. \quad &121b^4c^6. \\ 5. \quad &16a^8b^{10}x^4. & 6. \quad &25x^4y^{10}. & 7. \quad &4a^2b^2c^4. & 8. \quad &9c^2x^6. \\ 9. \quad &16x^2y^2z^6. & 10. \quad &\frac{4}{9}a^4b^0. & 11. \quad &\frac{4x^4}{9y^6}. & 12. \quad &\frac{16}{9x^4y^2}. \\ 13. \quad &\frac{49a^2b^2}{9}. & 14. \quad &\frac{9a^4b^6}{16c^{10}x^8}. & 15. \quad &\frac{1}{4x^2y^2}. & 16. \quad &4x^2y^4. \\ 17. \quad &\frac{25a^2b^6}{4x^2y^2}. & 18. \quad &169c^{10}x^6. & 19. \quad &\frac{1}{16a^8}. & 20. \quad &\frac{9a^{10}}{25x^6}. \\ 21. \quad &8a^3b^6. & 22. \quad &27x^9. & 23. \quad &64x^{12}. & 24. \quad &-27a^9b^3. \\ 25. \quad &-125a^3b^6. & 26. \quad &-b^9c^6x^3. & 27. \quad &-216a^{18}. & 28. \quad &-8a^{21}c^6. \\ 29. \quad &\frac{1}{27y^6}. & 30. \quad &-\frac{27x^{15}}{125a^9}. & 31. \quad &343x^3y^{12}. & 32. \quad &-\frac{8}{27}a^{15}. \\ 33. \quad &81a^9b^{12}. & 34. \quad &a^{12}x^6. & 35. \quad &-32x^{15}y^5. & 36. \quad &\frac{1}{128a^{14}}. \\ 37. \quad &\frac{243x^{20}}{32y^{15}}. & 38. \quad &\frac{256x^{24}}{6561y^8}. & 39. \quad &-\frac{x^{21}}{2187}. & 40. \quad &\frac{64x^{30}}{729a^{24}}. \end{aligned}$$

XV. b. PAGE 101.

$$\begin{aligned} 1. \quad &a^2+6ab+9b^2. & 2. \quad &a^2-6ab+9b^2. & 3. \quad &x^2-10xy+25y^2. \\ 4. \quad &4x^2+12xy+9y^2. & 5. \quad &9x^2-6xy+y^2. & 6. \quad &9x^2+30xy+25y^2. \\ 7. \quad &81x^2-36xy+4y^2. & 8. \quad &25a^2b^2-10abc+c^2. & 9. \quad &p^2q^2-2pqr+r^2. \\ 10. \quad &x^2-2abcx+a^2b^2c^2. & 11. \quad &a^2x^2+4abxy+4b^2y^2. \\ 12. \quad &x^4-2x^2+1. & 13. \quad &a^2+b^2+c^2-2ab-2ac+2bc. \\ 14. \quad &a^2+b^2+c^2+2ab-2ac-2bc. & 15. \quad &a^2+4b^2+c^2+4cb+2ac+4bc. \\ 16. \quad &4a^2+9b^2+16c^2-12ab+16ac-24bc. \\ 17. \quad &x^4+y^4+z^4-2x^2y^2-2x^2z^2+2y^2z^2. \\ 18. \quad &x^2y^2+y^2z^2+z^2x^2+2xy^2z+2x^2yz+2xyz^2. \\ 19. \quad &9p^2+4q^2+16r^2-12pq+24pr-16qr. \\ 20. \quad &x^3-2x^2+3x^2-2x+1. & 21. \quad &4x^4+12x^3+5x^2-6x+1. \\ 22. \quad &x^3+y^3+a^3+b^3-2xy+2ax-2bx-2ay+2by-2ab. \\ 23. \quad &4x^2+9y^2+a^3+4b^3+12xy+4ax-8bx+6ay-12by-4ab. \\ 24. \quad &m^2+n^2+p^2+q^2-2mn-2mp-2mq+2np+2nq+2pq. \\ 25. \quad &\frac{a^2}{4}+4b^2+\frac{c^2}{16}-2ab+\frac{ac}{4}-bc. & 26. \quad &\frac{a^3}{9}+9b^2+\frac{9}{4}-2ab-a+9b. \\ 27. \quad &\frac{4x^4}{9}-\frac{4x^3}{3}+3x^2-3x+\frac{9}{4}. \end{aligned}$$

XV. c. PAGE 101.

1. $x^3 + 3ax^2 + 3a^2x + a^3$.
2. $x^3 - 3ax^2 + 3a^2x - a^3$.
3. $x^3 - 6x^2y + 12xy^2 - 8y^3$.
4. $8x^3 + 12x^2y + 6xy^2 + y^3$.
5. $27x^3 - 135x^2y + 225xy^2 - 125y^3$.
6. $a^3b^3 + 3a^2b^2c + 3abc^2 + c^3$.
7. $8a^3b^3 - 36a^2b^2c + 54abc^2 - 27c^3$.
8. $125a^3 - 75a^2bc + 15ab^2c^2 - b^3c^3$.
9. $x^6 + 12x^4y^2 + 48x^2y^4 + 64y^6$.
10. $64x^6 - 240x^4y^2 + 300x^2y^4 - 125y^6$.
11. $8a^9 - 36a^6b^2 + 54a^3b^4 - 27b^6$.
12. $125x^{15} - 300x^{10}y^4 + 240x^5y^8 - 64y^{12}$.
13. $a^3 - 2a^2b + \frac{4}{3}ab^2 - \frac{8}{27}b^3$.
14. $\frac{1}{27}a^3 + \frac{2}{3}a^2 + 4a + 8$.
15. $\frac{1}{27}x^6 - x^5 + 9x^4 - 27x^3$.
16. $\frac{1}{216}a^3 + \frac{1}{6}a^2x + 2ax^2 + 8x^3$.

XVI. a. PAGE 103.

1. $2ab^2$.
2. $3x^3y$.
3. $5x^2y^3$.
4. $4a^2bc^3$.
5. $9a^3b^4$.
6. $10x^4$.
7. $a^{10}b^8c^2$.
8. a^4bc^6 .
9. $8x^3y^6$.
10. $\frac{6}{a^{18}}$.
11. $\frac{a^8b^4}{4}$.
12. $\frac{17y^2}{5}$.
13. $\frac{18x^6}{13y^3}$.
14. $\frac{9a^9}{6b^5}$.
15. $\frac{16xy^2}{17p^7}$.
16. $\frac{20a^{20}b^{10}}{9x^5y^6}$.
17. $3a^2bc$.
18. $-2a^4b^3$.
19. $4x^2yz^4$.
20. $-7a^4b^6$.
21. $-\frac{x^4y^3}{5}$.
22. $\frac{2x^3}{9y^5}$.
23. $\frac{5ab^2}{6x^2y^3}$.
24. $-\frac{3x^9}{4y^{21}}$.
25. a^2x^3 .
26. x^2y^3 .
27. $2xy^2$.
28. $3a^3b$.
29. $2ax^3$.
30. $-x^2y^3$.
31. $\frac{2}{a^9b^8}$.
32. $\frac{a^3x^5}{b^{10}}$.
33. $\frac{a^2}{b^8c^4}$.

XVI. b. PAGE 105.

1. $x + 2y$.
2. $3a + 2b$.
3. $x - 5y$.
4. $2x - 3y$.
5. $9x + y$.
6. $5x - 3y$.
7. $x^2 - y^2$.
8. $1 - a^3$.
9. $a^2 - a + 1$.
10. $2x^2 - 3x + 5$.
11. $3x^2 - 2x - 1$.
12. $x^2 - 2x + 1$.
13. $2a^2 + a - 2$.
14. $1 - 5x + x^2$.
15. $2x + 3y - 5z$.
16. $4x^3 + 2x^4 - x^5$.
17. $x^3 - 11x + 17$.
18. $5x^2 - 3ax + 4a^2$.
19. $2x^2 + y^2 - 3z^2$.
20. $ab - 2ac + 3bc$.
21. $2a^2 + b^2 - 3c^2$.
22. $2x^2 - xy + 3y^2$.
23. $3x^2 - 5x + 7$.
24. $1 - 2x + 3x^2 - 4x^3$.
25. $ax^5 - 2bx^2 + 3c$.

XVI. c. PAGE 106.

- $+3a^2x - a^3$
 $2y^2 + 6xy^2 + y^3$
 $b^2c + 3abc^2 + c^3$
 $5a^2bc + 15ab^2c^2 - b^3c^3$
1. $\frac{2}{3} - 3$. 2. $2 - \frac{x}{y}$. 3. $\frac{x}{5} + y$. 4. $\frac{x}{y} + 5$. 5. $\frac{x}{2y} - 2$.
 6. $\frac{x}{y} - \frac{a}{5}$. 7. $\frac{8x}{3y} + 2$. 8. $\frac{3x}{5} - \frac{5}{3x}$. 9. $\frac{a^2}{8} + \frac{a}{2} - 1$.
 10. $x^2 + x - \frac{1}{2}$. 11. $a^2 - \frac{3}{2}a + \frac{5}{3}$. 12. $x^2 - 3x + \frac{1}{3}$.
 13. $\frac{a^2}{2} + \frac{a}{x} - \frac{x}{a}$. 14. $x^2 - x + \frac{1}{4}$. 15. $\frac{x^2}{2} - 2x + \frac{a}{3}$.
 16. $\frac{3a}{x} - \frac{1}{5} + \frac{2x}{3a}$. 17. $4m^2 + \frac{2}{3}n + 1$. 18. $2x^2 + 8 + \frac{8}{x^2}$.

$$\frac{1}{6}a^2x + 2ax^2 + 8x^3.$$

XVI. d. PAGE 109.

1. $a + 1$. 2. $x + 2$. 3. $ax - y^2$.
 4. $2m - 1$. 5. $4a - 3b$. 6. $1 + x + x^2$.
 7. $1 - 2x + 3x^2$. 8. $a + 2b - c$. 9. $2a^2 - 3a + 1$.
 10. $y^2 - y + 1$. 11. $2x^2 + x - 3$. 12. $3x^2 - 2xa + 3a^2$.
 13. $3x^2 - x - 1$. 14. $x^2 - 2xy + 4y^2$. 15. $3x^2 - x + 6$.

XVI. e. PAGE 110.

1. $\frac{x}{2} - 1$. 2. $\frac{x}{3} + 2$. 3. $2x - \frac{y^2}{3}$.
 4. $\frac{3x}{4y} - 2$. 5. $x - \frac{3}{x}$. 6. $\frac{x^2}{y} - 2y^2$.
 7. $\frac{x}{y} + 2 - \frac{y}{x}$. 8. $\frac{x}{3} - 1 + \frac{3}{x}$. 9. $\frac{x}{a} - 4 + \frac{2a}{x}$.
 10. $\frac{4a}{x} - 4 + \frac{x}{a}$. 11. $\frac{a}{b} - 1 + \frac{b}{a}$. 12. $\frac{2x^2}{y^2} + \frac{4x}{y} - 3$.

XVII. a. PAGE 112.

1. $a(a^2 - x)$. 2. $x^2(x - 1)$. 3. $2a(1 - a)$. 4. $a(a - b^2)$.
 5. $p(7p + 1)$. 6. $2x(4 - x)$. 7. $5ax(1 - a^2x)$.
 8. $x^2(3 + x^2)$. 9. $x(x + y)$. 10. $x^2(x - y)$.
 11. $5x(1 - 5xy)$. 12. $5(3 + 5x^2)$. 13. $16x(1 + 4xy)$.
 14. $15a^2(1 - 15a^2)$. 15. $27(2 - 3x)$. 16. $5x^2(2 - 5xy)$.
 17. $x(3x^2 - x + 1)$. 18. $2x^3(3 + x + 2x^2)$. 19. $x(x^2 - xy + y^2)$.
 20. $3a^2(a^2 - ab + 2b^2)$. 21. $2xy^2(xy - 3x + y)$.
 22. $3x(2x^2 - 3xy + 4y^2)$. 23. $5x^3(x^2 - 2a^2 - 3a^3)$.
 24. $7a(1 - a^2 + 2a^3)$. 25. $19a^2x^2(2x^3 + 3a)$.

E, A,

2 D

XVII. b. PAGE 113.

- | | | |
|------------------------|--------------------------|------------------------|
| 1. $(a+b)(a+c)$. | 2. $(a-c)(a+b)$. | 3. $(ac+d)(ac+b)$. |
| 4. $(a+3)(a+c)$. | 5. $(2+c)(x+c)$. | 6. $(x-a)(x+5)$. |
| 7. $(5+b)(a+b)$. | 8. $(a-y)(b-y)$. | 9. $(a-b)(x-z)$. |
| 10. $(p+q)(r-s)$. | 11. $(x-y)(m-n)$. | 12. $(x-a)(m+n)$. |
| 13. $(2x+y)(a+b)$. | 14. $(3a-b)(x-y)$. | 15. $(2x+y)(3x-a)$. |
| 16. $(x-2y)(m-n)$. | 17. $(ax-3by)(x-y)$. | 18. $(x+my)(x-4y)$. |
| 19. $(a+b)(x^2+2)$. | 20. $(x-3)(x-y)$. | 21. $(2x-1)(x^3+2)$. |
| 22. $(3x+5)(x^2+1)$. | 23. $(x+1)(x^3+2)$. | 24. $(y-1)(y^2+1)$. |
| 25. $(a+bc)(xy-z)$. | 26. $(f^2+g^2)(x^2-a)$. | 27. $(2x+3y)(ax-by)$. |
| 28. $(ax+by)(mx-ny)$. | 29. $(a-b-c)(x-y)$. | 30. $(a+b)(ax+by+c)$. |

XVII. c. PAGE 115.

- | | | |
|--------------------------|------------------------------|--------------------------------|
| 1. $(a+1)(a+2)$. | 2. $(a+1)(a+1)$. | 3. $(a+3)(a+4)$. |
| 4. $(a-4)(a-3)$. | 5. $(x-5)(x-6)$. | 6. $(x-7)(x-8)$. |
| 7. $(x-9)(x-10)$. | 8. $(x+6)(x+7)$. | 9. $(x-10)(x-11)$. |
| 10. $(x-9)(x-12)$. | 11. $(x-5)(x-16)$. | 12. $(x+6)(x+15)$. |
| 13. $(x-7)(x-12)$. | 14. $(x-6)(x-13)$. | 15. $(x-3)(x-15)$. |
| 16. $(x+8)(x+12)$. | 17. $(x-11)(x-15)$. | 18. $(x-13)(x-8)$. |
| 19. $(x+17)(x+6)$. | 20. $(a-19)(a-5)$. | 21. $(a-16)(a-16)$. |
| 22. $(a+15)(a+15)$. | 23. $(a+27)(a+27)$. | 24. $(a-19)(a-19)$. |
| 25. $(a-7b)(a-7b)$. | 26. $(a+2b)(a+3b)$. | 27. $(m-5n)(m-8n)$. |
| 28. $(m-7n)(m-15n)$. | 29. $(x-11y)(x-12y)$. | 30. $(x-13y)(x-13y)$. |
| 31. $(x^2+1)(x^2+7)$. | 32. $(x^2+2y^2)(x^2+7y^2)$. | 33. $(xy-3)(xy-13)$. |
| 34. $(x+24y)(x+25y)$. | 35. $(xy+17)(xy+17)$. | 36. $(a^2b^2+25)(a^2b^2+12)$. |
| 37. $(a-5bx)(a-15bx)$. | 38. $(x+13y)(x+30y)$. | 39. $(a-2b)(a-27b)$. |
| 40. $(x^2+81)(x^2+81)$. | 41. $(4-x)(3-x)$. | 42. $(5+x)(4+x)$. |
| 43. $(12-x)(11-x)$. | 44. $(8+x)(11+x)$. | 45. $(26+xy)(5+xy)$. |
| 46. $(13-xa)(11-xa)$. | 47. $(17-x^2)(12-x^2)$. | 48. $(27+x)(8+x)$. |

XVII. d. PAGE 116.

- | | | |
|----------------------|---------------------|----------------------|
| 1. $(x+1)(x-2)$. | 2. $(x+2)(x-1)$. | 3. $(x+2)(x-3)$. |
| 4. $(x+3)(x-2)$. | 5. $(x+1)(x-3)$. | 6. $(x+3)(x-1)$. |
| 7. $(x+8)(x-7)$. | 8. $(x+8)(x-5)$. | 9. $(x+2)(x-6)$. |
| 10. $(a+4)(a-5)$. | 11. $(a+3)(a-7)$. | 12. $(a+5)(a-4)$. |
| 13. $(a+9)(a-13)$. | 14. $(x+12)(x-3)$. | 15. $(x+13)(x-12)$. |
| 16. $(x+11)(x-10)$. | 17. $(x+6)(x-15)$. | 18. $(x+15)(x-16)$. |

- $(ac+d)(ac+b)$.
 $(x-a)(x+5)$.
 $(a-b)(x-z)$.
 $(x-a)(m+n)$.
 $(2x+y)(3x-a)$.
 $(x+my)(x-4y)$.
 $(2x-1)(x^3+2)$.
 $(y-1)(y^2+1)$.
 $(2x+3y)(ax-by)$.
 $(a+b)(ax+by+c)$.
19. $(a+5)(a-17)$. 20. $(a+8)(a-19)$. 21. $(xy+3)(xy-8)$.
 22. $(x+12y)(x-5y)$. 23. $(x+7a)(x-6a)$. 24. $(x+3y)(x-35y)$.
 25. $(a+14y)(a-15y)$. 26. $(x+23)(x-5)$. 27. $(x+4y)(x-24y)$.
 28. $(x+26)(x-10)$. 29. $(a+2)(a-13)$. 30. $(ay+24)(ay-10)$.
 31. $(a^2+7b^2)(a^2-8b^2)$. 32. $(x^2+3)(x^2-17)$. 33. $(y^2+9x^2)(y^2-3x^2)$.
 34. $(ab+2c)(ab-5c)$. 35. $(a+14bx)(a-2bx)$.
 36. $(a+9xy)(a-27xy)$. 37. $(x^2+25a^2)(x^2-12a^2)$.
 38. $(x^2+11a^2)(x^2-12a^2)$. 39. $(x^2+21a^2)(x^2-22a^2)$.
 40. $(x^2+30)(x^2-29)$. 41. $(1+x)(2-x)$. 42. $(2+x)(3-x)$.
 43. $(11+x)(10-x)$. 44. $(20+x)(19-x)$. 45. $(15+ax)(8-ax)$.
 46. $(5+xy)(13-xy)$. 47. $(14+x)(7-x)$. 48. $(17+x)(12-x)$.

XVII. e. PAGE 119.

- $(a+3)(a+4)$.
 $(x-7)(x-8)$.
 $(x-10)(x-11)$.
 $(x+6)(x+15)$.
 $(x-3)(x-15)$.
 $(x-13)(x-8)$.
 $(a-16)(a-16)$.
 $(a-19)(a-19)$.
 $(m-5n)(m-8n)$.
 $(x-13y)(x-13y)$.
 $(xy-3)(xy-13)$.
 $(a^2b^2+25)(a^2b^2+12)$.
 $(a-2b)(a-27b)$.
 $(5+x)(4+x)$.
 $(26+xy)(5+xy)$.
 $(27+x)(8+x)$.
1. $(x+1)(2x+1)$. 2. $(x+1)(3x+2)$. 3. $(x+2)(2x+1)$.
 4. $(x+3)(3x+1)$. 5. $(x+4)(2x+1)$. 6. $(x+2)(3x+2)$.
 7. $(x+2)(2x+3)$. 8. $(x+5)(2x+1)$. 9. $(x+3)(3x+2)$.
 10. $(x+2)(5x+1)$. 11. $(x+2)(2x-1)$. 12. $(x+1)(3x-2)$.
 13. $(x+3)(4x-1)$. 14. $(x+5)(3x-1)$. 15. $(x+8)(2x-1)$.
 16. $(2x+1)(x-1)$. 17. $(x+3)(3x-2)$. 18. $(x+4)(2x-7)$.
 19. $(x+6)(3x-5)$. 20. $(2x+3)(3x-1)$. 21. $(3x+1)(2x-3)$.
 22. $(3x+4)(x+1)$. 23. $(x+7)(3x+2)$. 24. $(2x+5)(x-3)$.
 25. $(x+7)(3x-2)$. 26. $(x-7)(3x+2)$. 27. $(3x-5)(2x-7)$.
 28. $(4x-7)(x+2)$. 29. $(x-2)(3x-7)$. 30. $(x+13)(3x+2)$.
 31. $(x+5)(4x+3)$. 32. $(2x+y)(x-3y)$. 33. $(2x-7)(4x-5)$.
 34. $(3x-2y)(4x-5y)$. 35. $(15x-1)(x+15)$. 36. $(15x-2)(x-5)$.
 37. $(12x+5)(x-3)$. 38. $(12x-7)(2x+3)$. 39. $(8x-9)(9x-8)$.
 40. $(8x+y)(3x-4y)$. 41. $(2+x)(1-2x)$. 42. $(3-x)(1+4x)$.
 43. $(2+3x)(3-2x)$. 44. $(4+3x)(1-2x)$. 45. $(1+7x)(5-3x)$.
 46. $(7+3x)(1+x)$. 47. $(6-x)(3-5x)$. 48. $(4+5x)(2-x)$.
 49. $(5+4x)(4-5x)$. 50. $(8-9x)(3+8x)$.

XVII. f. PAGE 120.

- $(x+2)(x-3)$.
 $(x+3)(x-1)$.
 $(x+2)(x-6)$.
 $(a+5)(a-4)$.
 $(x+13)(x-12)$.
 $(x+15)(x-16)$.
1. $(x+2)(x-2)$. 2. $(a+9)(a-9)$. 3. $(y+10)(y-10)$.
 4. $(c+12)(c-12)$. 5. $(3+a)(3-a)$. 6. $(7+c)(7-c)$.
 7. $(11+x)(11-x)$. 8. $(20+a)(20-a)$. 9. $(x+3a)(x-3a)$.
 10. $(y+5x)(y-5x)$. 11. $(6x+5b)(6x-5b)$. 12. $(3x+1)(3x-1)$.
 13. $(6p+7q)(6p-7q)$. 14. $(2k+1)(2k-1)$. 15. $(7+10k)(7-10k)$.
 16. $(1+5x)(1-5x)$. 17. $(a+2b)(a-2b)$. 18. $(3x+y)(3x-y)$.

19. $(pq+6)(pq-6)$. 20. $(ab+2cd)(ab-2cd)$.
 21. $(x^2+3)(x^2-3)$. 22. $(3a^2+11)(3a^2-11)$.
 23. $(5x+8)(5x-8)$. 24. $(9a^2+7x^2)(9a^2-7x^2)$.
 25. $(x^3+5)(x^3-5)$. 26. $(1+6a^3)(1-6a^3)$. 27. $(3x^2+a)(3x^2-a)$.
 28. $(9x^2+5a)(9x^2-5a)$. 29. $(x^2a+7)(x^2a-7)$.
 30. $(a+8x^3)(a-8x^3)$. 31. $(ab+3x^3)(ab-3x^3)$. 32. $(x^3y^3+2)(x^3y^3-2)$.
 33. $(1+ab)(1-ab)$. 34. $(2+x)(2-x)$. 35. $(3+2a)(3-2a)$.
 36. $(3a^2+5b^2)(3a^2-5b^2)$. 37. $(x^2+4b)(x^2-4b)$.
 38. $(x+5y)(x-5y)$. 39. $(1+10b)(1-10b)$. 40. $(5+8x)(5-8x)$.
 41. $(11a+9x)(11a-9x)$. 42. $(pq+8a^2)(pq-8a^2)$.
 43. $(8x+5z^3)(8x-5z^3)$. 44. $(7x^2+4y^2)(7x^2-4y^2)$.
 45. $(9p^2z^3+5b)(9p^2z^3-5b)$. 46. $(4x^3+3y^3)(4x^3-3y^3)$.
 47. $(6x^{18}+7a^7)(6x^{18}-7a^7)$. 48. $(1+10a^3b^2c)(1-10a^3b^2c)$.
 49. $(5x^5+4a^4)(5x^5-4a^4)$. 50. $(ab^2c^3+x^8)(ab^2c^3-x^8)$.
 51. $1000 \times 150 = 150000$. 52. $241 \times 1 = 241$.
 53. $1000 \times 500 = 500000$. 54. $658 \times 20 = 13160$.
 55. $1008 \times 500 = 503000$. 56. $200 \times 2 = 400$.
 57. $2000 \times 1446 = 2892000$. 58. $2378 \times 900 = 2140200$.
 59. $2500 \times 1122 = 2805000$. 60. $3000 \times 2462 = 7386000$.
 61. $16264 \times 2 = 32528$. 62. $10002 \times 10000 = 100020000$.

XVII. g. PAGE 121.

1. $(a+b+c)(a+b-c)$. 2. $(a-b+c)(a-b-c)$.
 3. $(x+y+2z)(x+y-2z)$. 4. $(x+2y+a)(x+2y-a)$.
 5. $(a+3b+4x)(a+3b-4x)$. 6. $(x+5a+3y)(x+5a-3y)$.
 7. $(x+5c+1)(x+5c-1)$. 8. $(a-2x+b)(a-2x-b)$.
 9. $(2x-3a+3c)(2x-3a-3c)$. 10. $(a+b-c)(a-b+c)$.
 11. $(x+y+z)(x-y-z)$. 12. $(2a+y-z)(2a-y+z)$.
 13. $(3x+2a-3b)(3x-2a+3b)$. 14. $(1+a-b)(1-a+b)$.
 15. $(c+5a-3b)(c-5a+3b)$. 16. $(a+b+c+d)(a+b-c-d)$.
 17. $(a-b+x+y)(a-b-x-y)$. 18. $(7x+y+1)(7x+y-1)$.
 19. $(a+b+m-n)(a+b-m+n)$. 20. $(a-n+b+m)(a-n-b-m)$.
 21. $(b-c+a-x)(b-c-a+x)$. 22. $(4a+x+b+y)(4a+x-b-y)$.
 23. $(a+2b+3x+4y)(a+2b-3x-4y)$. 25. $(a-b+x-y)(a-b-x+y)$.
 24. $(1+7a-3b)(1-7a+3b)$. 27. $(2a-5x+1)(2a-5x-1)$.
 26. $(a-3x+4y)(a-3x-4y)$. 30. $y(2x+y)$.
 28. $(a+b-c+x-y+z)(a+b-c-x-y-z)$.
 29. $(3a+2b+c+x-2y)(3a+2b-c-x+2y)$.

31. $y(2x - y)$. 32. $(x + 5y)(x + y)$. 33. $47x(x + 2y)$.
 34. $(8x + y)(2x + 3y)$. 35. $5y(6x - 5y)$. 36. $(12x - 1)(2x + 7)$.
 37. $5a(a + 2)$. 38. $(7a + 1)(a - 1)$. 39. $3a(a + 2b - 2c)$.
 40. $x(x - 14y + 2z)$. 41. $y(2x + y - 16)$. 42. $a(4x + a - 6)$.

XVII. h. PAGE 122.

1. $(x + y + a)(x + y - a)$. 2. $(a - b + x)(a - b - x)$.
 3. $(x - 3a + 4b)(x - 3a - 4b)$. 4. $(2a + b + 3c)(2a + b - 3c)$.
 5. $(x + a + y)(x + a - y)$. 6. $(a + y + x)(a + y - x)$.
 7. $(x + a + b)(x - a - b)$. 8. $(y + c - x)(y - c + x)$.
 9. $(1 + x + y)(1 - x - y)$. 10. $(c + x - y)(c - x + y)$.
 11. $(x + y + 2xy)(x + y - 2xy)$. 12. $(a - 2b + 3ac)(a - 2b - 3ac)$.
 13. $(x + y + a + b)(x + y - a - b)$. 14. $(a - b + c + d)(a - b - c - d)$.
 15. $(x - 2a + b - y)(x - 2a - b + y)$. 16. $(y + b + a + 3x)(y + b - a - 3x)$.
 17. $(x - 1 + a + 2b)(x - 1 - a - 2b)$. 18. $(3a - 1 + x + 4d)(3a - 1 - x - 4d)$.
 19. $(x - y + a - b)(x - y - a + b)$. 20. $(a - b + c + d)(a - b - c - d)$.
 21. $(2x - 3a + c + k)(2x - 3a - c - k)$.
 22. $(a - 5b + 3bx - 1)(a - 5b - 3bx + 1)$.
 23. $(a^2 + 4x^2 + 5x^3 - 3)(a^2 + 4x^2 - 5x^3 + 3)$.
 24. $(x^2 - a^2 + x - 3)(x^2 - a^2 - x + 3)$.
 25. $(a^2 + ab + b^2)(a^2 - ab + b^2)$. 26. $(x^2 + 2xy + 4y^2)(x^2 - 2xy + 4y^2)$.
 27. $(p^2 + 3pq + 9q^2)(p^2 - 3pq + 9q^2)$. 28. $(c^2 + cd + 2d^2)(c^2 - cd + 2d^2)$.
 29. $(x^2 + 3xy - y^2)(x^2 - 3xy - y^2)$.
 30. $(2m^2 + 3mn + n^2)(2m^2 - 3mn + n^2)$.

XVII. k. PAGE 123.

1. $(x - y)(x^2 + xy + y^2)$. 2. $(x + y)(x^2 - xy + y^2)$.
 3. $(x - 1)(x^2 + x + 1)$. 4. $(1 + a)(1 - a + a^2)$.
 5. $(2x - y)(4x^2 + 2xy + y^2)$. 6. $(x + 2y)(x^2 - 2xy + 4y^2)$.
 7. $(3x + 1)(9x^2 - 3x + 1)$. 8. $(1 - 2y)(1 + 2y + 4y^2)$.
 9. $(ab - c)(a^2b^2 + a^2c + c^2)$. 10. $(2x + 3y)(4x^2 - 6xy + 9y^2)$.
 11. $(1 - 7x)(1 + 7x + 49x^2)$. 12. $(4 + y)(16 - 4y + y^2)$.
 13. $(5 + a)(25 - 5a + a^2)$. 14. $(6 - a)(36 + 6a + a^2)$.
 15. $(ab + 8)(a^2b^2 - 8ab + 64)$. 16. $(10y - 1)(100y^2 + 10y + 1)$.
 17. $(x + 4y)(x^2 - 4xy + 16y^2)$. 18. $(3 - 10x)(9 + 30x + 100x^2)$.
 19. $(ab + 6c)(a^2b^2 - 6abc + 36c^2)$. 20. $(7 - 2x)(49 + 14x + 4x^2)$.
 21. $(a + 3b)(a^2 - 3ab + 9b^2)$. 22. $(3x - 4y)(9x^2 + 12xy + 16y^2)$.
 23. $(5x - 1)(25x^2 + 5x + 1)$. 24. $(6p - 7)(36p^2 + 42p + 49)$.
 25. $(xy + z)(x^2y^2 - xyz + z^2)$. 26. $(abc - 1)(a^2b^2c^2 + abc + 1)$.

27. $(7x + 10y)(49x^2 - 70xy + 100y^2)$.
 28. $(9a - 4b)(81a^2 + 36ab + 16b^2)$.
 29. $(2ab + 5x)(4a^2b^2 - 10abx + 25x^2)$.
 30. $(xy - 6z)(x^2y^2 + 6xyz + 36z^2)$.
 31. $(x^2 - 3y)(x^4 + 3x^2y + 9y^2)$.
 32. $(4x^2 + 5y)(16x^4 - 20x^2y + 25y^2)$.
 33. $(2x - z^2)(4x^2 + 2xz^2 + z^4)$.
 34. $(6x^2 - b)(36x^4 + 6x^2b + b^2)$.
 35. $(a + 7b)(a^2 - 7ab + 49b^2)$.
 36. $(a^2 + 9b)(a^4 - 9a^2b + 81b^2)$.
 37. $(2x - 9y^2)(4x^2 + 18xy^2 + 81y^4)$.
 38. $(pq - 3x)(p^2q^2 + 3pqx + 9x^2)$.
 39. $(z - 4y^2)(z^2 + 4zy^2 + 16y^4)$.
 40. $(xy - 8)(x^2y^2 + 8xy + 64)$.

XVII. 1. PAGE 125.

1. $(a + y + z)(a - y - z)$.
 2. $(x + yz)(x^2 - xyz + y^2z^2)(x - yz)(x^2 + xyz + y^2z^2)$.
 3. $(3x - 11)(2x + 7)$.
 4. $(3y + 2x)(9y^2 - 6xy + 4x^2)(3y - 2x)(9y^2 + 6xy + 4x^2)$.
 5. $(x + 4)(x^2 - 4x + 16)(x - 4)(x^2 + 4x + 16)$.
 6. $(m + n + x - y)(m + n - x + y)$.
 7. $(11x + 13)(3x - 5)$.
 8. $(a^2 + b^2 + c^2 + d^2)(a^2 + b^2 - c^2 - d^2)$.
 9. $(m - n)(m^2 + mn + n^2)(x + y)$.
 10. $4a(b + c)$.
 11. $(x + 2 + a - y)(x + 2 - a + y)$.
 12. $(x - 17)(x + 7)$.
 13. $(a - d + b - c)(a - d - b + c)$.
 14. $(x - y + a)(x - y - a)$.
 15. $(a + x + y + z)(a + x - y - z)$.
 16. $(7x - 3)(3x + 13)$.
 17. $(1 + ax - by)(1 - ax + by)$.
 18. $(cd - 1)(c^2d^2 + cd + 1)(c + a)(c - a)$.
 19. $(a + b)(a - b)(x + y)(x^2 - xy + y^2)(x - y)(x^2 + xy + y^2)$.
 20. $(x - 19)(x + 13)$.
 21. $(x + y)(x - y)(a - c)(a^2 + ac + c^2)$.
 22. $(cx + d)(ax - b)$.
 23. $(a + b)(a - b)(x + y)$.
 24. $(x^2 + 2y^2z^2)(x^2 + 2y^2z^2)$.
 25. $(ab + 8)(a^2b^2 - 8ab + 64)$.
 26. $(2x + 7)(x + 5)$.
 27. $20y(5x + y)(5x - y)$.
 28. $a^2(a - 2b)(a^2 + 2ab + 4b^2)$.
 29. $x^3(a + 4y)(a - 4y)$.
 30. $(b - c + a)(b - c - a)$.
 31. $5x^2(x - 6)(x + 3)$.
 32. $7a^2x(2x - a)(x - 2a)$.
 33. $(x^4 + 1)(x^2 + 1)(x + 1)(x - 1)$.
 34. $(1 + m - n)(1 - m + n)$.
 35. $3(5x^2 + 4a^2)(5x^2 - 4a^2)$.
 36. $5ab(ab - 1)(a^2b^2 + ab + 1)$.
 37. $4y(2x + 3)(x + 5)$.
 38. $(3xy + 5a)(xy + 7a)$.
 39. $ab(3a + b)(9a^2 - 3ab + b^2)(3a - b)(9a^2 + 3ab + b^2)$.
 40. $a^2(ax + 2y)(a^2x^2 - 2axy + 4y^2)(ax - 2y)(a^2x^2 + 2axy + 4y^2)$.
 41. $(a^2 + b^2)(a^4 - a^2b^2 + b^4)(a + b)(a^2 - ab + b^2)(a - b)(a^2 + ab + b^2)$.

42. $6y^2(4x+3y)(x-2y)$. 43. $\{(a+b)^2+1\}(a+b+1)(a+b-1)$.
 44. $\{a^2+(b+c)^2\}(a+b+c)(a-b-c)$.
 45. $(c+d-1)\{(c+d)^2+c+d+1\}$. 46. $(1-x+y)\{1+x-y+(x-y)^2\}$.
 47. $2\{5(a-b)+1\}\{25(a-b)^2-5(a-b)+1\}$. 48. $2c(c^2+3d^2)$.
 49. $9y(4x^2+2xy+y^2)$. 50. $(x-2y)(x+2y+1)$.
 51. $(a-b)(a+b+1)$. 52. $(a+b)(a+b+1)$.
 53. $(a+b)(a^2-ab+b^2+1)$. 54. $(a+3b)(a-3b+1)$.
 55. $(x-y)\{2(x-y)+1\}\{2(x-y)-1\}$. 56. $xy(x+y)(x-y)(x-y)$.

Miscellaneous Examples III. PAGE 126.

1. x^3-2x . 2. $42a-40b+10c$. 3. a^6-c^6 .
 4. (1) 12; (2) $x=5$, $y=6$. 5. x^3+4x-1 .
 6. 72. 7. $\frac{109}{210}$. 8. $x^2+\frac{3}{4}x+\frac{5}{4}$.
 9. $2x^2-x$. 10. (1) $(ax-5)(ax+3)$; (2) $(2m^2+9pq)(2m^2-9pq)$.
 11. (1) $x=-2$, $y=4$; (2) $x=5$, $y=-2$. 12. $\frac{am}{pb}$ miles.
 13. $84x^2+25x^3+101x-30$. 14. (1) 7; (2) $-1\frac{1}{2}$.
 15. $x^4+14x^3+27x^2-154x+121$.
 16. (1) $(x+2a)(x-b)$; (2) $(x^2+14y)(x^2-4y)$.
 17. L.C.F. = 7; L.C.M. = $3528a^3b^2c^3$. 18. £14. 19. 3p.
 21. (1) $m-n=a+c$; (2) $3a^2b^2+c^3=p(m+n)$.
 22. 8. 23. $6x^2-xy-y^2$.
 24. Apples 4d. a dozen; eggs 1s. 4d. a score.
 25. $(x-5)(2x-3)(x+1)$. 26. 33. 27. $2x^2+9xy-7y^2$.
 28. $x=2$, $y=3$, $z=0$.
 29. (1) $xy(x+2y)(x-2y)$; (2) $(m+n)(m-n)(2m^2+3n^2)$.
 30. $\frac{bc}{am}$ days, $2\frac{5}{8}$.

XVIII. a. PAGE 129.

1. $a+b$. 2. $x+y$. 3. $x(x-y)$. 4. $2x-3y$.
 5. $x+y$. 6. $ab(a-b)$. 7. $a(a-x)$. 8. $a+2x$.
 9. $b(a+b)$. 10. $x-3y$. 11. $a-x$. 12. $2x+y$.
 13. $2(5x-1)$. 14. $3x+2y$. 15. $x+1$. 16. $y(x-1)$.
 17. $(x-y)^2$. 18. x^2+a^2 . 19. $x+2y$. 20. $x-3a$.
 21. $x+2$. 22. $x-5$. 23. $x-3$. 24. $x-3$.
 25. $3x+1$. 26. $x-1$. 27. $cx+d$. 28. x^2+y .
 29. $x(a-3b)$. 30. $2x+1$. 31. $x^2(3x+2)$.

XVIII. b. PAGE 133.

- | | | |
|-------------------------|-----------------------------|-------------------------|
| 1. $x^3 - 3x + 2.$ | 2. $x^3 - 13x + 5.$ | 3. $x^3 - 8.$ |
| 4. $x^3 - 5.$ | 5. $x^3 + 2x + 1.$ | 6. $x + 3.$ |
| 7. $a^3 - 2ax + x^3.$ | 8. $x + 1.$ | 9. $x^3 - 3x + 7.$ |
| 10. $2x^3 - 7.$ | 11. $3x^2 + 1.$ | 12. $2x^3 - 3.$ |
| 13. $3x^2 + 2a^3.$ | 14. $x^3 - ax + a^3.$ | 15. $x^2 + 2ax - a^2.$ |
| 16. $3a^3 - ax - 2x^2.$ | 17. $xy(2x^2 + xy - 3y^2).$ | 18. $2x^2a^3(2x - 3a).$ |
| 19. $2x^2(2x + 7).$ | 20. $6(3x - 5a).$ | 21. $3x^2 - 2xy + y^2.$ |
| 22. $x^4 + x^3 - 1.$ | 23. $1 + x^3 - x^4.$ | 24. $1 + a.$ |
| 25. $x(3 + 4x).$ | 26. $x^3 - 2x + 1.$ | 27. $2x^3 - 7.$ |

XIX. a. PAGE 137.

- | | | | |
|---------------------------------|---------------------------|-----------------------------|--------------------------------|
| 1. $\frac{3}{2b}.$ | 2. $\frac{b}{c}.$ | 3. $\frac{1}{ax-1}.$ | 4. $\frac{3b^2c}{20(a-b)}.$ |
| 5. $\frac{2x-3y}{2x}.$ | 6. $4(x-y).$ | 7. $\frac{1}{2a+3x}.$ | 8. $\frac{x}{x^3-2y^3}.$ |
| 9. $\frac{x-3y}{x^2+3xy+9y^2}.$ | 10. $\frac{x}{x+1}.$ | 11. $\frac{3x}{x+2}.$ | 12. $\frac{5a}{3b}.$ |
| 13. $\frac{xy}{x-2}.$ | 14. $\frac{3(a+b)}{a-b}.$ | 15. $\frac{x^2-17}{x^2-5}.$ | 16. $\frac{x+2y}{x^2+xy+y^2}.$ |
| 17. $\frac{2x+3}{3x+5}.$ | 18. $\frac{a(x-4)}{x+5}.$ | 19. $\frac{x+7}{x+13}.$ | 20. $\frac{3+a}{2}.$ |

XIX. b. PAGE 139.

The expression in [] is in each case the H.C.F. of the numerator and the denominator.

- | | |
|--|--|
| 1. $\frac{a-2b}{a+2b} [(a^2+ab+b^2)].$ | 2. $\frac{x-3}{x+2} [(x-1)^2].$ |
| 3. $\frac{a+5}{a+4} [(a-1)(a-2)].$ | 4. $\frac{x^2+4xy-9y^2}{2x^2+3xy+7y^2} [2x-3y].$ |
| 5. $\frac{2a+5b}{3a+5b} [(2a+3b)(a-b)].$ | 6. $\frac{1-x+2x^2}{1-x+3x^2} [1+x+x^2].$ |
| 7. $\frac{x-1}{3x^2+3x+10} [x-1].$ | 8. $\frac{3a^2+b^2}{4a-b} [a-b].$ |
| 9. $\frac{4x^2-ax+a^2}{x^3+a^3} [x+a].$ | 10. $\frac{2(2x^2-3x-1)}{3x^3+x^2+x-2} [x-1].$ |
| 11. $(2x-3a)^2 [(2x+3a)^2].$ | 12. $\frac{3x^2-x-2}{3x^2+x-2} [2x+1].$ |

$$\begin{aligned}
 &x^2 - 8, \\
 &x + 3, \\
 &x^2 - 3x + 7, \\
 &2x^2 - 3, \\
 &x^2 + 2ax - a^2, \\
 &2x^2a^2(2x - 3a), \\
 &3x^2 - 2xy + y^2, \\
 &1 + a, \\
 &2x^2 - 7.
 \end{aligned}$$

$$13. \frac{5x+2}{7x-4} [x^2 - 3].$$

$$14. \frac{2x^2+3x+5}{2x^2+3x-5} [2x^2 - 3x + 5].$$

$$15. \frac{3(x-3a)(x-4a)}{2(x+3a)(x+4a)} [x-2a].$$

$$16. \frac{a(x+8a)}{x(x+7a)} [x^2 - 13ax + 5a^2].$$

XIX. c. PAGE 142.

$$1. \frac{7}{12},$$

$$2. \frac{ab}{2a-1},$$

$$3. 2,$$

$$4. \frac{a-11}{a-2},$$

$$5. \frac{4x+3a}{x+2},$$

$$6. \frac{5a-b}{x(3a-2)},$$

$$7. \frac{x+2}{x-1},$$

$$8. \frac{x+1}{x+5},$$

$$9. \frac{x}{x-2},$$

$$10. \frac{2x-1}{2x-3},$$

$$11. \frac{x+1}{x+5},$$

$$12. \frac{x-1}{4x+7},$$

$$13. b^2 + 3b + 9,$$

$$14. \frac{1}{x+7},$$

$$15. 8pq - z^2,$$

$$16. x,$$

$$17. \frac{x+1}{x-1},$$

$$18. \frac{x-5}{x-1},$$

$$19. \frac{x}{y},$$

$$20. x,$$

$$21. \frac{2x-1}{2x-5},$$

$$22. 1,$$

$$23. \frac{1}{b},$$

$$24. \frac{a+b-c}{b-c-a},$$

$$25. \frac{1}{x-8},$$

$$26. \frac{a-x}{a+x},$$

$$27. \frac{m}{n},$$

$$28. x(2+x),$$

$$29. \frac{x+4}{x(x-4)},$$

$$30. 1,$$

$$31. a+x$$

$$32. \frac{a^2}{16a^2+4ab+b^2}.$$

XX. a. PAGE 144.

$$1. x(x+1),$$

$$2. x^2(x-3),$$

$$3. 12x^2(x+2),$$

$$4. 21x^3(x+1),$$

$$5. x(x+1)(x-1),$$

$$6. ab(a+b),$$

$$7. xy(2x+1)(2x-1),$$

$$8. 6x(3x-1),$$

$$9. x(x+1)(x+2),$$

$$10. (x+1)(x-1)(x-2),$$

$$11. (x+2)^2(x+3),$$

$$12. (x-1)(x-2)(x-4),$$

$$13. (x-3)(x-1)(x+2),$$

$$14. (x+5)(x-4)(x-6),$$

$$15. (x+7)(x-6)(x-5),$$

$$16. (x+1)(x+2)(2x+1),$$

$$17. (x+2)(x+3)(3x+2),$$

$$18. (x+2)(x+3)(5x+1),$$

$$19. (x+2)(x+8)(2x-1),$$

$$20. (x+2)(x-2)(3x-7),$$

$$21. 12x(x+2)(2x+1)(4x-7),$$

$$22. 6x^2(x+7)(3x+5)(3x-2),$$

$$23. 20x^2y(3x+1)(5x+1)(4x-1),$$

$$24. (x+y)(2x-7y)(4x-5y),$$

$$25. (x-y)(3x-2y)(4x-5y),$$

$$26. 3a^2x(3x-a)(2x+3a)(x+5a),$$

$$27. 2axy^3(x+3)(4x-1)(3x-2),$$

$$28. x^2(3-5x)^2(2+x)^2,$$

$$29. 42a^4b^2(a-b)^3(a+b)(a^2+ab+b^2),$$

$$30. m^2n(m^6-n^6)(m-n)^2,$$

$$31. 8c^2(2c-3d)^2(8c^3-27d^3),$$

of the numerator

1)].

$$\frac{9y^2}{7y^2} [2x-3y],$$

$$[1+x+x^2]$$

-b].

$$\frac{-1}{-2} [x-1],$$

$$[2x+1].$$

XX. b. PAGE 146.

1. H.C.F. $x-2$. L.C.M. $(x+1)^2(x+2)(x-2)(x-3)$.
2. $(ax+b)(ax-b)(bx+a)$. 3. $xy(x-a)(y-b)(y-2b)$.
4. H.C.F. $x(x+3)$. L.C.M. $x(x-1)(x+3)(2x-1)$.
5. $(1+x)^3(1-x)^2$. 6. $(x-2)(x-4)(x-6)$.
7. H.C.F. $2x+1$. L.C.M. $(2x+1)(x+1)(x-1)(3x+2)(3x-2)$.
8. $ab^2c^2(c+a)^2(c-a)^2$.
9. L.C.M. $y^2(x-y)^2(x^2+xy+y^2)$. H.C.F. $x-y$.
10. H.C.F. $2x-3$. L.C.M. $(2x-3)(3x-2)(x+4)(3x+4)$.
11. $(x+a)^2(x^2+ax+a^2)(x^2-ax+a^2)$.
12. H.C.F. $3x-y$. L.C.M. $(3x-y)(x+y)^2(x-y)^2$.
13. $x-1$. 14. $(a+b)(a-b)(a-2b)(a^2+ab+b^2)$.
15. H.C.F. a^2+xy . L.C.M. $(a^2+xy)(2x+3y)(2x-3y)$.
16. H.C.F. $(x-3)(x-4)$. L.C.M. $(x-2)(x-3)(x-4)(x-5)$.
17. $x-8a$. 18. $105x^2y^2(x+y)^2(x-y)^2$.

XXI. a. PAGE 150.

1. $\frac{4(x+1)}{5}$. 2. $\frac{13(x-2)}{12}$. 3. $\frac{25x-61}{56}$. 4. $\frac{17x}{36}$.
5. $\frac{19x-201}{225}$. 6. $\frac{12x^2+28x-27}{8x^2}$. 7. 0. 8. $\frac{3(a+3b)}{8a}$.
9. $\frac{6b^2c+6bc^2+3ac^2+3a^2c-4a^2b+4ab^2}{12abc}$. 10. $\frac{a^2+3x^2}{2ax}$.
11. $\frac{5x+31}{102x}$. 12. $\frac{a^4b^2-b^4c^2+a^2c^4}{a^3b^2c^3}$. 13. $\frac{11x^3-18x^2-27x-16}{30x^3}$.
14. $\frac{x^3+y^3}{x^2y^3}$. 15. $\frac{3y+2z}{yz}$. 16. $\frac{a^3+b^3+c^3-3abc}{abc}$.

XXI. b. PAGE 151.

1. $\frac{2x+5}{(x+2)(x+3)}$. 2. $\frac{x+5}{(x+3)(x+4)}$. 3. $\frac{1}{(x-4)(x-5)}$.
4. $\frac{2(x+6)}{(x-6)(x+2)}$. 5. $\frac{(a-b)x}{(x+a)(x+b)}$. 6. $\frac{(a+b)x-2ab}{(x-a)(x-b)}$.
7. $\frac{2}{(x+2)(x+4)}$. 8. $\frac{4ax}{a^2-x^2}$. 9. $\frac{8x}{x^2-4}$.
10. $\frac{6}{(x-2)(x-5)}$. 11. $\frac{ax}{x^2-a^2}$. 12. $\frac{5x+9}{x^2-9}$.
13. $\frac{x+2y}{4x^2-9y^2}$. 14. $\frac{3ax}{x^2-4a^2}$. 15. $\frac{4ab}{4a^2-b^2}$.

$$\begin{aligned} & x-3), \\ & y-b)(y-2b), \\ & -1), \\ & 4)(x-6), \\ &)(3x+2)(3x-2), \end{aligned}$$

$$y, \\ 4)(3x+4).$$

$$\begin{aligned} & y)^2, \\ & b)(a^2+ab+b^2), \\ & 2x-3y), \\ & (x-4)(x-5), \\ & y)^2. \end{aligned}$$

$$\begin{aligned} & \hat{61}, \quad 4. \frac{17x}{36}, \\ & 8. \frac{3(a+3b)}{8a}, \\ & 10. \frac{a^2+3x^2}{2ax}, \\ & -18x^2-27x-16 \\ & \frac{30x^3}{abc} \\ & +c^3-3abc \end{aligned}$$

$$\begin{aligned} & 3. \frac{1}{(x-4)(x-5)}, \\ & 6. \frac{(a+b)x-2ab}{(x-a)(x-b)}, \\ & 9. \frac{8x}{x^2-4}, \\ & 12. \frac{5x+9}{x^2-9}, \\ & 15. \frac{4ab}{4a^2-b^2}, \end{aligned}$$

$$16. \frac{2xy}{x^2-y^2},$$

$$19. \frac{5x^2}{25x^2-y^2},$$

$$22. \frac{2x^3}{x^2-y^2},$$

$$25. \frac{4(x-1)}{(x-2)^2(x+2)},$$

$$17. \frac{2x^3}{1-x^4},$$

$$20. \frac{x^4+y^4}{xy(x^4-y^4)},$$

$$23. -\frac{2ax}{a^3-8x^3},$$

$$26. \frac{x^2+a^2}{ax(x-a)(x+a)^2},$$

$$18. \frac{x^2+y^2}{xy(x^2-y^2)},$$

$$21. \frac{4a^2}{x(x+2a)},$$

$$24. 2b.$$

XXI. c. PAGE 153.

$$1. \frac{2}{x+y},$$

$$2. \frac{x}{4x^2-y^2},$$

$$3. \frac{1-6x^2}{1-4x^2},$$

$$4. \frac{4a^2+b^2}{4a^2-9b^2},$$

$$5. \frac{1+a}{9-a^3},$$

$$6. \frac{4x-5}{6(x^2-1)},$$

$$7. 0,$$

$$8. \frac{12a^2-4a+7}{3(4a^2-9)},$$

$$9. \frac{2(13x+7)}{3(x^2-4)},$$

$$10. \frac{x^2+y^2}{x^4+x^2y^2+y^4},$$

$$11. \frac{2}{(x-4)(x-6)},$$

$$12. \frac{2}{(x-2)(x-3)(x-4)},$$

$$13. \frac{2}{(x-1)(2x+1)(2x+3)},$$

$$14. \frac{1}{(x-1)(2x+1)(3x-2)},$$

$$15. \frac{17a}{(1-2a)(4+a)(3+5a)},$$

$$16. \frac{23x}{(1+2x)(2+x)(5-9x)},$$

$$17. \frac{x+2}{(x+1)(x+3)},$$

$$18. \frac{1}{x+1},$$

$$19. \frac{1}{a+b},$$

$$20. \frac{3x+2}{(x-2)(x-1)(x+1)},$$

$$21. \frac{1}{x+2},$$

$$22. \frac{8x^2+4x-3}{(x-1)(x+1)(2x+1)},$$

$$23. \frac{1}{2x+1},$$

$$24. \frac{x^2+11}{(x-1)(x+2)(x+3)},$$

$$25. \frac{2x+13}{(x+3)(x+4)(x-4)},$$

$$26. -\frac{32a^2}{(1-2a)^2(1+2a)},$$

$$27. \frac{96x^2}{(3-2x)^2(3+2x)},$$

$$28. \frac{4x^3}{81-x^4},$$

$$29. \frac{72a}{16a^4-81},$$

$$30. \frac{1}{1-x^4},$$

$$31. \frac{a(a^2+2ax+3x^2)}{4(a^4-x^4)},$$

$$32. \frac{16x}{16-x^4},$$

$$33. \frac{x(37+172x^2)}{6(1-16x^4)},$$

$$34. \frac{2a(a^2+32x^2)}{3(a^4-256x^4)},$$

$$35. \frac{7a^2+45}{6(a^4-81)},$$

$$36. \frac{x^5}{1-x^3},$$

$$37. \frac{36a^4}{a^3-6561},$$

$$38. \frac{2}{x^2(x^2-4)},$$

$$39. \frac{1}{(3x-y)(x-3y)},$$

$$40. \frac{2}{(x-1)(x+1)^2},$$

$$41. 1,$$

$$42. 0,$$

$$43. \frac{4x}{x^2-1},$$

XXI. d. PAGE 156.

1. $\frac{x-11}{20(x^2-1)}$
2. $\frac{1}{1-a^2}$
3. $\frac{x+3a}{x+a}$
4. $\frac{2x-a}{x+a}$
5. 0.
6. $\frac{7x}{1-x^2}$
7. $\frac{1}{x-3}$
8. $\frac{12(2x+1)}{4x^2-9}$
9. $\frac{61-21b}{12(1-b^2)}$
10. $\frac{2}{3(1-a^2)}$
11. $\frac{y^5}{x^6-y^6}$
12. $\frac{x}{y}$
13. $\frac{2x}{x+y}$
14. $\frac{2x^3}{x^2-4}$
15. $\frac{a}{4a^2-25b^2}$
16. $\frac{b(a+b)}{x^2-b^2}$
17. $\frac{2bx}{4x^2-1}$
18. $\frac{x+c}{(x-a)(x-b)}$
19. $\frac{x-c}{(x-a)(x-b)}$
20. $\frac{2a}{(x-a)(x-b)}$
21. 0.
22. $\frac{4a^3}{x^4-a^4}$
23. $\frac{48a^3}{(x^2-a^2)(x^2-9a^2)}$
24. $\frac{x^4}{a^5-b^5}$
25. 0.
26. $\frac{a^3}{a^5-b^5}$
27. $\frac{a-x}{a+x}$
28. $\frac{a^3}{(a-b)(a^2+b^2)}$
29. $\frac{2+x+3x^2}{2(1-x^4)}$
30. $\frac{2(x^2+1)}{x(x^2-1)}$
31. 0.
32. $\frac{4ab}{a^3-b^3}$

XXI. e. PAGE 159.

1. 0.
2. $\frac{bc+ca+ab-a^2-b^2-c^2}{(a-b)(b-c)(c-a)}$
3. $\frac{x^2+y^2+z^2-yz-zx-xy}{(x-y)(y-z)(z-x)}$
4. 0.
5. $\frac{2(bc+ca+ab-a^2-b^2-c^2)}{(a-b)(b-c)(c-a)}$
5. 0.
7. 0.
8. 0.
9. $\frac{2(qr+rp+pq-p^2-q^2-r^2)}{(p-q)(q-r)(r-p)}$
10. 0.
11. 0.
12. $\frac{p(y-z)+q(z-x)+r(x-y)}{(y-z)(z-x)(x-y)}$

XXII. a. PAGE 163

1. $\frac{m^2-nl}{na-mb}$
2. $\frac{x+y}{y-x}$
3. $\frac{ad+b}{dx-y}$
4. $\frac{x+c}{b-x}$
5. $\frac{3}{4b}$
6. $\frac{a}{c}$
7. $\frac{x^2-y^2}{x^2+y^2}$
8. $\frac{c}{ac+b}$
9. $\frac{ad}{bd+c}$
10. $\frac{nx}{nx-m}$
11. $\frac{pn(ad+bc)}{bd(pm+kn)}$
12. $x-1$
13. $\frac{x(x+3)}{x+4}$
14. $-\frac{x+1}{x^2(x+3)}$
15. $-\frac{x^2(2x+3)}{x+2}$
16. $\frac{1}{x}$

$$4. \frac{2x-a}{x+a}.$$

$$8. \frac{12(2x+1)}{4x^2-9}.$$

$$12. \frac{x}{y}.$$

$$16. \frac{b(a+b)}{x^2-b^2}.$$

$$\frac{x-c}{(x-a)(x-b)}.$$

$$\frac{48a^3}{(x^2-a^2)(x^2-9a^2)}.$$

$$\frac{a-x}{a+x}.$$

$$30. \frac{2(x^2+1)}{x(x^2-1)}.$$

$$y^2+z^2-yz-zx-xy$$

$$-y)(y-z)(z-x).$$

$$6. 0.$$

$$\frac{pq-p^2-q^2-r^2}{(q-r)(r-p)}.$$

$$\frac{q(z-x)+r(x-y)}{(z-x)(x-y)}.$$

$$4. \frac{x+c}{b-x}.$$

$$8. \frac{c}{ac+b}.$$

$$12. x-1.$$

$$16. \frac{1}{x}.$$

$$17. \frac{a^2-b^2}{2}.$$

$$18. 2.$$

$$19. \frac{y^4}{x^2+y^2}.$$

$$20. \frac{1}{2x^2-1}.$$

$$21. \frac{2(a+b)}{a-b}.$$

$$22. a+x.$$

$$23. \frac{4}{x^2}.$$

$$24. \frac{1+x}{1+x^2}.$$

$$25. \frac{a^2-a+1}{2a-1}.$$

$$26. \frac{6+x+2y}{8x(y+6)}.$$

$$27. \frac{a(yz+n)}{xyz+nx+mz}.$$

$$28. 1-x.$$

$$29. \frac{x^2-3x+1}{x^2-4x+1}.$$

$$30. \frac{2}{x^2}.$$

$$31. \frac{a-c}{1+ac}.$$

$$32. \frac{b}{a}.$$

$$33. \frac{1}{a+x}.$$

$$34. 4.$$

$$35. 8x^2-1.$$

$$36. 2x^2.$$

XXII. b. PAGE 167.

$$1. \frac{x}{3} + \frac{y}{9} - \frac{y^2}{9x}.$$

$$2. \frac{a^2}{4} - \frac{ax}{3} + \frac{x^2}{2}.$$

$$3. \frac{a^2}{2b} - \frac{3a}{2} + \frac{3b}{2} + \frac{b^2}{2a}.$$

$$4. \frac{1}{bc} + \frac{1}{ca} + \frac{1}{ab}.$$

$$5. \frac{1}{a} + \frac{1}{b} + \frac{1}{c}.$$

$$6. \frac{a^2}{6} - \frac{b^2}{2} + \frac{1}{3}.$$

$$7. x-x^2+x^3-x^4; \text{ Rem. } x^5.$$

$$8. 1 + \frac{b}{a} + \frac{b^2}{a^2} + \frac{b^3}{a^3}; \text{ Rem. } \frac{b^4}{a^3}.$$

$$9. 1+2x+2x^2+2x^3; \text{ Rem. } 2x^4.$$

$$10. 1+x-x^3-x^4; \text{ Rem. } x^5.$$

$$11. x-3+\frac{9}{x}-\frac{27}{x^2}; \text{ Rem. } \frac{81}{x^3}.$$

$$12. 1+2x+3x^2+4x^3; \text{ Rem. } 5x^4-4x^5.$$

$$17. \frac{x-3}{x-4}.$$

$$18. 3(a-2x)^2.$$

$$19. \frac{b^2-3b-2}{b-6}.$$

$$20. \frac{a^2-4b^2}{a+3b}.$$

$$21. \frac{(2x-3)(2x+7)}{6}.$$

XXII. c. PAGE 168.

$$1. \frac{4(c-x)}{3(a+x)}.$$

$$2. \frac{x(x+a)}{2}.$$

$$3. \frac{1}{a^2+ab-2b^2}.$$

$$4. \frac{8xy(x^2+y^2)}{(x^2-y^2)^2}.$$

$$5. \frac{4x(2-x)}{(x-1)(x^2+1)}.$$

$$6. \frac{x^4}{1-x^8}.$$

$$7. \frac{1}{x(x+1)^2(1+x+x^2)}.$$

$$8. \frac{1-x+x^2}{1+x+x^2}.$$

$$9. \frac{2x+3}{3(x+b)}.$$

$$10. \frac{bx+a}{ax+b}.$$

$$11. \frac{ax^3(x^2+a^2)}{x^3+a^3}.$$

$$12. \frac{a^6}{(a-x)(a+x)^2}.$$

$$13. \frac{(x+1)^2}{3x^3+6x^2-x-8}.$$

$$14. \frac{x+y}{a^2-y}.$$

$$15. \frac{2(a^2+a+1)}{a(a+1)(a+2)}.$$

$$16. \frac{1}{2(3-2x)}.$$

$$17. \frac{4}{1-x^4}.$$

18. 1. 19. $\frac{(2x-1)(x+1)}{(x+2)(x-1)}$ 20. $\frac{x-2}{4x^2-5x-6}$
 21. $\frac{x}{(x-2a)^2}$ 22. $\frac{a(a^2+x^2)}{(x-a)(a+x)^2}$ 23. 1. 24. 1.
 25. $9x - \frac{1}{x}$ 26. $\frac{a}{2x^2}$ 27. $\frac{1}{2x(2x-1)}$ 28. $\frac{b^4}{b^2+a^2}$
 29. $\frac{ab}{a+b}$ 30. x 31. x 32. bx 33. $\frac{a^2-b^2}{2}$
 34. 1. 35. $\left(x - \frac{1}{x}\right)^2$ 36. 1. 37. 1.
 38. $\frac{x(x+1)}{x^2+4x+1}$ 39. $\frac{12}{(a^4-4)(a^4-1)}$ 40. $\frac{3m^2}{(3m+2n)(9m^2-n^2)}$
 41. $\frac{1+x+x^2}{(1+x)(1+x^2)(1-x)^2}$ 42. $\frac{2x}{(x-2)(x+1)^2}$ 43. $\frac{1}{x+y}$
 44. 1. 45. 1. 46. 1. 47. 0.
 48. $\frac{a^2+b^2+c^2-bc-ca-ab}{(b-c)(c-a)(a-b)}$ 49. 1. 50. 1. 51. 0.
 52. $2y+a+b$ 53. $\frac{(2a^2+x^2)(a-x)}{a^2x}$ 54. $\frac{28(x+4)}{9(x+3)}$
 55. $\frac{7(x-4)}{4(x-1)}$ 56. $x+3$ 57. $1+a-a^3$ 58. $-\frac{c}{e}$

Miscellaneous Examples IV. PAGE 172.

1. $-\frac{1}{22}$ 2. 6. 3. $abc(b-c)$; -6. 4. 7. 5. $\frac{5}{3}$
 6. (1) 232; (2) -29. 7. (1) -19; (2) 0. 8. 1. 9. $-\frac{3}{10}$
 10. (1) -12; (2) 1. 11. 1. 12. $8\frac{1}{2}$. 13. $98x-2y$; $19\frac{1}{3}$.
 14. (1) 1; (2) 21. 15. $(x+9)(x+12)$. 16. $(a-7)(a+13)$.
 17. $(x-8y)(x-12y)$. 18. $(ab-17)(ab+3)$. 19. $c(c+13)(c-12)$.
 20. $n(m-3n)(m-3n)$. 21. $(p^2+7q^2)(p^2-8q^2)$.
 22. $(d^2+5c^2)(d+3c)(d-3c)$. 23. $xy(x+6y)(x-7y)$.
 24. $(m+13)(m+15)$. 25. $(14-a)(15+a)$. 26. $(19-pq)(3+pq)$.
 27. $(x^2+16)(x^2+11)$. 28. $(a^2+14)(a^2-7)$. 29. $(c+27)(c+27)$.
 30. $(9-xy)(8+xy)$. 31. $(a^2+2x^2)(a^2+7x^2)$.
 32. $(p-12q)(p+9q)$. 33. $2(a^3+12)(a^3-11)$.
 34. $x^2(x-9)(x+7)$. 35. $(bc+12)(bc-7)$. 36. $(z+17)(z+17)$.
 37. $(a-3c)(a-19c)$. 38. $yz(y-7)(y+13)$.
 39. $(2+3x^3)(1-x)(1+x+x^3)$. 40. $(2ab-5)(ab+3)$.

$$20. \frac{x-2}{4x^2-5x-5}.$$

$$1. \quad 24. \quad 1.$$

$$28. \frac{b^4}{b^3+a^2}.$$

$$b.x. \quad 33. \frac{a^2-b^2}{2}.$$

$$1. \quad 37. \quad 1.$$

$$\frac{3n^2}{(3m+2n)(9m^2-n^2)}.$$

$$43. \frac{1}{x+y}.$$

$$1. \quad 51. \quad 0.$$

$$28. \frac{(x+4)}{9(x+3)}.$$

$$-a^3. \quad 58. \quad -\frac{c}{e}.$$

GE 172.

$$7. \quad 5. \quad \frac{5}{3}.$$

$$1. \quad 9. \quad -\frac{3}{10}.$$

$$98x-2y; \quad 19\frac{1}{3}.$$

$$(a-7)(a+13).$$

$$c(c+13)(c-12).$$

$$p^2-8q^2.$$

$$(x-7y).$$

$$(19-pq)(3+pq).$$

$$(c+27)(c+27).$$

$$x^2+7x^2).$$

$$(z+17)(z+17).$$

$$(2ab-5)(ab+3).$$

$$41. (3p-4)(3p-4). \quad 42. (5+mn)(7+mn). \quad 43. (17+c)(7-c).$$

$$44. x^2(2-x)(3-x). \quad 45. (2m+3)(3m-1). \quad 46. (2a-5b)(2a+b).$$

$$47. (6p-q)(p-2q). \quad 48. (5x+4z)(4x-5z). \quad 49. (2x^2+3)(4x^2-5).$$

$$50. 6(2y-1)(y-2). \quad 51. (3ab+4)(4ab-3).$$

$$52. (2a^2b-5)(a^2b-2). \quad 53. (7x+8y)(3x-2y).$$

$$54. (9m-5n)(2m+3n). \quad 55. (c+a-b)(c-a+b).$$

$$56. (a+b-c)(a-b+c). \quad 57. (5x+3y)(25x^2-15xy+9y^2).$$

$$58. (ab+7)(a^2b^2-7ab+49). \quad 59. (8b-a^2)(64b^2+8ba^2+a^4).$$

$$60. (a+2x-2y)(a-2x+2y). \quad 61. (m+n+1)(m+n-1).$$

$$62. 2c^2(3c+d)(c-d). \quad 63. (a^2b^2-1+x-y)(a^2b^2-1-x+y).$$

$$64. (1+2m)(1-2m)(1-2m+4m^2)(1+2m+4m^2).$$

$$65. p^3(1+10q)(1-10q+100q^2). \quad 66. (81+a^2)(9+a)(9-a).$$

$$67. (x^2-1+y-z)(x^2-1-y+z). \quad 68. (a+4b-4c)(a-4b+4c).$$

$$69. (c-d)(1+2c-2d)(1-2c+2d). \quad 70. (p-4q)(p+4q+1).$$

$$71. 2[1+4a+4b][1-4(a+b)+16(a+b)^2].$$

$$72. (x+3y)(1+x^2-3xy+9y^2). \quad 73. (x+y)(x^2+y^2).$$

$$74. (cx-d)(ax+b). \quad 75. (7+a)(2-a).$$

$$76. (14x^2+y^2)(7x^2-y^2). \quad 77. (17+a)(3-a).$$

$$78. (1+m+p)(1-m-p). \quad 79. (bx-a)(ax-b).$$

$$80. (3b-c+4)(3b-c-4). \quad 81. (c+1)(c^2-c+1)(x+1)(x-1).$$

$$82. (3x-b)(x+2a). \quad 83. (m-n)(m+n+x)(m+n-x).$$

$$84. (a+b)(c+a-b)(c-a+b).$$

$$85. (x+2)(x^2-2x+4)(x^2+1)(x+1)(x-1). \quad 86. (x+1)(x+7)(2x-3).$$

$$87. (2x+5y)(x-3y)(2x-5y). \quad 88. 325a^3b^3(x^2-a^2)^3(x+2a).$$

$$89. 2x^2-9x+9. \quad 90. 2x^3(x^2-4)(x^2-16).$$

$$91. H.C.F. = a+b+c, \quad L.C.M. = (a+b+c)(a-b)(b-c)(c-a).$$

$$92. a+b-c. \quad 93. (a-b)^2(a+b). \quad 95. (a^4-b^4)(a+b-2c).$$

$$97. H.C.F. = (x-7)(x-3),$$

$$L.C.M. = (x-1)(x-2)(x-3)(x-4)(x-5)(x-7).$$

$$98. \frac{1}{(1-x)^2} \quad 99. \frac{x-9}{(x^2-9)(x-3)} \quad 100. \frac{6x+1}{(2x+1)^2(2x-1)}$$

$$101. \frac{2}{x} \quad 102. \frac{4}{(1-x^2)^2} \quad 103. \quad 1. \quad 104. \quad 0. \quad 105. \quad \frac{x}{9}$$

$$106. \frac{2x-y}{x^2-y^2} \quad 107. \quad y-x. \quad 108. \quad ab. \quad 109. \quad 2(ac+bd)(ad+bc).$$

$$110. \quad 1. \quad 111. \quad 1. \quad 112. \quad \frac{(x^2+2)(x^4+1)}{x} \quad 113. \quad \frac{1}{f-g}.$$

$$114. \quad \frac{1}{x+1} \quad 115. \quad x(1+x-x^2). \quad 116. \quad \frac{3abc}{a+b} \quad 117. \quad a+b. \quad 118. \quad 1.$$

XXIII. a. PAGE 180.

- | | | | | |
|---------------------|----------------------|-----------------------|-----------------------|-----------------------|
| 1. 6. | 2. $1\frac{3}{10}$. | 3. $\frac{1}{5}$. | 4. 1. | 5. 20. |
| 6. 2. | 7. $\frac{7}{17}$. | 8. 0. | 9. 2. | 10. $-6\frac{5}{8}$. |
| 11. 5. | 12. 6. | 13. $-\frac{8}{11}$. | 14. $-\frac{7}{18}$. | 15. 1. |
| 16. -10. | 17. -4. | 18. $3\frac{3}{8}$. | 19. 3. | 20. 4. |
| 21. 6. | 22. 13. | 23. -7. | 24. 2. | 25. $2\frac{1}{2}$. |
| 26. 4. | 27. $1\frac{1}{2}$. | 28. 14. | 29. $\frac{1}{4}$. | 30. $2\frac{1}{4}$. |
| 31. $\frac{1}{6}$. | 32. 3. | 33. 20. | | |

XXIII. b. PAGE 182.

- | | | | |
|---------------------------|---------------------------|---------------------------|--------------------------------|
| 1. $\frac{2b-3a}{a-5b}$. | 2. $a+b$. | 3. $\frac{b^2-a^2}{2b}$. | 4. $\frac{a^3-ab+b^2}{a-b}$. |
| 5. 3. | 6. $m-n$. | 7. $-\frac{ab}{a+b+c}$. | 8. $\frac{a^2-2ab+bc}{c-b}$. |
| 9. a . | 10. $\frac{7bc}{9b+4c}$. | 11. $\frac{2ab}{a+b}$. | 12. $17a$. |
| 13. $\frac{1}{c}$. | 14. $3a+2b$. | 15. $\frac{a+b}{2}$. | 16. $\frac{a^2-2b^2}{3a-4b}$. |
| 17. $\frac{a}{17}$. | 18. $\frac{a^3}{b}$. | 19. $\frac{bc^2}{a^2}$. | 20. a . |
| 21. $a+b$. | 22. $\frac{2a}{21}$. | 23. $\frac{a+2b}{2}$. | 24. $\frac{a}{3}$. |
| | | | 25. $\frac{b(2a-b)}{a}$. |

XXIII. c. PAGE 185.

- | | |
|---|---|
| 1. $x = \frac{al-bm}{a^2-b^2}, y = \frac{am-bl}{a^2-b^2}$. | 2. $x = \frac{nq-mr}{lq-mp}, y = \frac{lr-np}{lq-mp}$. |
| 3. $x = \frac{bc}{a^2+b^2}, y = \frac{ac}{a^2+b^2}$. | 4. $x = \frac{a^2+ab+b^2}{a+b}, y = -\frac{ab}{a+b}$. |
| 5. $x = \frac{a'^2-a}{a'-a^2}, y = \frac{1-aa'}{a'-a^2}$. | 6. $x = \frac{q^2-pr}{qr-p^2}, y = \frac{pq-r^2}{qr-p^2}$. |
| 7. $x = \frac{a+a'}{a'b+ab'}, y = \frac{b'-b}{a'b+ab'}$. | 8. $x=2a, y=2b$. |
| 9. $x = \frac{2}{3}a, y = \frac{1}{2}b$. | 10. $x = \frac{pa}{q}, y = \frac{rb}{p}$. |

5. 20.

10. $-6\frac{5}{8}$.

15. 1.

20. 4.

25. $2\frac{1}{2}$.

30. $2\frac{1}{4}$.

11. $x = \frac{mm'(m+m')}{m^2+m'^2}, y = \frac{mm'(m-m')}{m^2+m'^2}$.

12. $x = \frac{qn}{ql-pm}, y = \frac{pm}{mp-lq}$.

14. $x = a+b, y = a-b$.

16. $x = \frac{2aa'b}{ab'+a'b}, y = \frac{2abb'}{ab'+a'b}$.

18. $x = m+l, y = m+l$.

20. $x = a+b, y = a-b$.

13. $x = \frac{c(a+b)}{2a}, y = \frac{c(a-b)}{2a}$.

15. $x = 3a, y = -2b$.

17. $x = a, y = 0$.

19. $x = \frac{a}{b}, y = \frac{b}{c}$.

21. $x = a^3 - b^3, y = a^3 + b^3$.

XXIV. PAGE 188.

1. 40.

2. 60.

3. 55.

4. £2. 12s.

5. Silk 9s. Calico 9d. per yard.

6. 54.

7. 42.

8. 48, 23.

9. $21\frac{9}{11}$ past one.

10. $17\frac{5}{11}$ past three.

11. $32\frac{8}{11}$ past six.

12. $51\frac{9}{11}$ past two.

13. 378, 216.

14. 15 persons; 5 shillings.

15. 8 yards at 4s. 6d.; 16 yards at 4s.

16. 17, 15.

17. 3 miles per hour.

18. 54.

19. $2\frac{1}{2}$ miles per hour.

20. $21\frac{9}{11}$ and $54\frac{6}{11}$ past seven. At $5\frac{5}{11}$ past.

21. $\frac{8}{12}$.

22. 10 P.M.; half way.

23. $1\frac{1}{3}$ hours.

24. £200.

25. 30 miles.

26. £36000.

27. £200.

28. 4 and 3 gallons.

29. $\frac{3}{5}$ and $\frac{2}{5}$ of a pint.

30. $\frac{pa}{p+q}$ miles.

31. 111 and 126 miles.

32. Coffee to chicory as 7 to 2.

33. $c-b$ and $a-c$ lbs.

34. $\frac{c}{2a}, \frac{2c}{b}$ yards.

36. 60 miles.

XXV. a. PAGE 194.

1. ± 5 .

2. ± 4 .

3. 3, -25.

4. 1, -25.

5. 3, 7.

6. ± 8 .

7. 3, -6.

8. 2, -7.

9. 9, -4.

10. 9, -8.

11. 31, -11.

12. 20, -11.

13. 4, -17.

14. 13, -12.

15. 11, -17.

16. 8, 15.

17. 7, 6.

18. 23, -1.

19. 6, $-\frac{16}{3}$.

20. $\frac{1}{3}, -\frac{3}{5}$.

21. $\frac{3}{2}, -\frac{1}{3}$.

22. $\frac{1}{5}, -4$.

23. ± 9 .

XXV. b. PAGE 197.

- | | | | |
|------------------------------------|-----------------------------------|-----------------------------------|-------------------------------------|
| 1. $\frac{11}{5}, -5.$ | 2. $11, \frac{11}{3}.$ | 3. $3, \frac{7}{6}.$ | 4. $\frac{15}{8}, -2.$ |
| 5. $\frac{7}{3}, 5.$ | 6. $2, -\frac{11}{6}.$ | 7. $\frac{3}{4}, -5.$ | 8. $\frac{7}{2}, -3.$ |
| 9. $\frac{13}{3}, -\frac{11}{3}.$ | 10. $\frac{7}{4}, \frac{2}{3}.$ | 11. $\frac{3}{4}, -\frac{4}{5}.$ | 12. $\frac{5}{8}, -3.$ |
| 13. $-\frac{5}{7}, -\frac{1}{3}.$ | 14. $\frac{9}{10}, -\frac{3}{5}.$ | 15. $\frac{13}{6}, -\frac{2}{3}.$ | 16. $3, -\frac{7}{5}.$ |
| 17. $\frac{a}{3}, -\frac{a}{5}.$ | 18. $\frac{3a}{7}, -\frac{a}{3}.$ | 19. $\frac{7k}{3}, -\frac{k}{2}.$ | 20. $-\frac{5k}{4}, -\frac{2k}{3}.$ |
| 21. $\frac{4c}{3}, -\frac{5c}{4}.$ | 22. $3, -\frac{4}{3}.$ | 23. $5, -\frac{5}{2}.$ | 24. $4, \frac{7}{5}.$ |
| 25. $3, -1.$ | 26. $2, \frac{1}{3}.$ | 27. $4, \frac{11}{2}.$ | 28. $7, 2.$ |
| 29. $11, 2.$ | 30. $4, \frac{4}{3}.$ | 31. $13, \frac{2}{3}.$ | 32. $6, \frac{40}{13}.$ |
| 33. $2, \frac{39}{8}.$ | 34. $3, -\frac{1}{2}.$ | 35. $12, -2.$ | 36. $5, \frac{23}{7}.$ |
| 37. $3a, \frac{3a}{2}.$ | 38. $2c, \frac{11c}{14}.$ | 39. $a, \frac{ab}{a-2b}.$ | |

XXV. c. PAGE 201.

- | | | | |
|------------------------------------|------------------------------------|------------------------------------|-----------------------------------|
| 1. $\frac{5}{3}, -3.$ | 2. $\frac{3}{2}, -5.$ | 3. $1, \frac{7}{2}.$ | 4. $\frac{3 \pm \sqrt{29}}{2}.$ |
| 5. $-4, -\frac{1}{5}.$ | 6. $\frac{7 \pm \sqrt{5}}{2}.$ | 7. $1, -\frac{7}{8}.$ | 8. $\frac{17 \pm \sqrt{89}}{10}.$ |
| 9. $7, -\frac{5}{2}.$ | 10. $\frac{1 \pm \sqrt{13}}{6}.$ | 11. $\frac{1}{3}, -2.$ | 12. $3, -\frac{11}{2}.$ |
| 13. $\frac{7}{6}, -1.$ | 14. $\frac{7}{4}, \frac{3}{2}.$ | 15. $\frac{7}{11}, -3.$ | 16. $-\frac{6}{5}, -4.$ |
| 17. $\frac{9}{10}, -\frac{5}{6}.$ | 18. $\frac{8}{3}, -\frac{3}{4}.$ | 19. $\frac{2}{7}, -14.$ | 20. $\frac{5}{12}, -\frac{3}{8}.$ |
| 21. $\frac{3}{5}, -\frac{2}{5}.$ | 22. $\frac{5}{2}, -\frac{7}{2}.$ | 23. $\frac{9a}{4}, -\frac{4a}{3}.$ | 24. $\frac{9a}{4}, \frac{4a}{3}.$ |
| 25. $\frac{5b}{3}, -\frac{7b}{3}.$ | 26. $\frac{7b}{6}, -\frac{5b}{6}.$ | 27. $2a, 2b.$ | 28. $2a, -8.$ |

29. $0, \frac{2a+b}{3}$ 30. $0, \frac{b-2}{a}$ 31. $\pm 2, \pm 1$ 32. $\pm 2, \pm 3$
 33. $1, -2$ 34. $3, -2$ 35. $\pm 4, \pm \frac{1}{4}$ 36. $\pm a, \pm b$
 37. $2, -3$ 38. $\pm 3, \pm 4$ 39. $3, -2, 4, -3$ 40. $4a, -2a, a$

XXVI. a. PAGE 203.

4. $\frac{15}{8}, -2$
 8. $\frac{7}{2}, -3$
 12. $\frac{5}{8}, -3$
 16. $3, -\frac{7}{5}$
 20. $-\frac{5k}{4}, -\frac{2k}{3}$
 24. $4, \frac{7}{5}$
 28. $7, 2$
 32. $6, \frac{40}{13}$
 36. $5, \frac{23}{7}$
 1. $x=17, 11; y=11, 17$ 2. $x=37, 14; y=14, 37$
 3. $x=53, 21; y=21, 53$ 4. $x=14, -9; y=9, -14$
 5. $x=27, -19; y=19, -27$ 6. $x=43, -25; y=25, -43$
 7. $x=71, 13; y=13, 71$ 8. $x=33, -41; y=41, -33$
 9. $x=52, -74; y=74, -52$ 10. $x=43, -51; y=-51, 43$
 11. $x=29, -47; y=47, -29$ 12. $x=22, -87; y=-87, 22$
 13. $x=\pm 8, \pm 5; y=\pm 5, \pm 8$ 14. $x=\pm 13, \pm 1; y=\pm 1, \pm 13$
 15. $x=\pm 4, \pm 7; y=\pm 7, \pm 4$ 16. $x=13, 3; y=3, 13$
 17. $x=10, 5; y=5, 10$ 18. $x=9, -5; y=5, -9$
 19. $x=12, -6; y=6, -12$ 20. $x=11, -8; y=8, -11$
 21. $x=9, 4; y=4, 9$ 22. $x=5, 4; y=4, 5$
 23. $x=7, -4; y=4, -7$ 24. $x=10, 4; y=4, 10$
 25. $x=12, -2; y=2, -12$ 26. $x=1; y=1$
 27. $x=4, 3; y=3, 4$ 28. $x=\frac{1}{a}; y=\frac{1}{b}$
 29. $x=\pm 1; y=\pm 1$

XXVI. b. PAGE 205.

4. $\frac{3 \pm \sqrt{29}}{2}$
 8. $\frac{17 \pm \sqrt{89}}{10}$
 12. $3, -\frac{11}{2}$
 16. $-\frac{6}{5}, -4$
 20. $\frac{5}{12}, -\frac{3}{8}$
 24. $\frac{9a}{4}, \frac{4a}{3}$
 28. $2a, -8$
 1. $x=7, 4; y=4, 7$ 2. $x=8, 5; y=5, 8$
 3. $x=14, 9; y=9, 14$ 4. $x=7, -5; y=5, -7$
 5. $x=11, -7; y=7, -11$ 6. $x=13, 0; y=0, -13$
 7. $x=\pm 6, \pm 4; y=\pm 4, \pm 6$ 8. $x=\pm 7, \pm 3; y=\pm 3, \pm 7$
 9. $x=\pm 9, \pm 5; y=\pm 5, \pm 9$ 10. $x=\pm 9, \pm 3; y=\pm 3, \pm 9$
 11. $x=\frac{6}{5}, \frac{8}{3}; y=\frac{8}{3}, \frac{6}{5}$ 12. $x=\pm 6, \pm 5; y=\pm 5, \pm 6$
 13. $x=4, 2; y=2, 4$ 14. $x=7, -3; y=3, -7$
 15. $x=5, 3; y=3, 5$ 16. $x=4, -2; y=2, -4$
 17. $x=8, -2; y=2, -8$ 18. $x=5, 1; y=1, 5$
 19. $x=5, 1; y=1, 5$ 20. $x=\frac{1}{6}, -\frac{1}{5}; y=\frac{1}{5}, -\frac{1}{6}$

XXVI. c. PAGE 208.

1. $x=4, -\frac{3}{5}$; $y=3, -20$.
2. $x=\pm 3$; $y=\pm 2$.
3. $x=12, 8$; $y=2, -2$.
4. $x=2, \frac{10}{3}$; $y=5, 3$.
5. $x=4, 7$; $y=1, 10$.
6. $x=4, -3$; $y=1, -\frac{4}{3}$.
7. $x=1, -\frac{71}{17}$; $y=4, \frac{112}{17}$.
8. $x=\pm 2, \pm \frac{4}{\sqrt{5}}$; $y=\pm 1, \pm \frac{3}{\sqrt{5}}$.
9. $x=2, \frac{5}{8}$; $y=-7, -\frac{1}{8}$.
10. $x=\pm 4, \pm 6$; $y=\pm 2, \pm 4$.
11. $x=\pm 3, \pm 4$; $y=\pm 2, \pm 5$.
12. $x=\pm \frac{3}{2}, \pm \frac{1}{2}$; $y=\pm \frac{1}{2}, \pm \frac{3}{2}$.
13. $x=\pm 2, \pm 1$; $y=\pm 1, \pm 2$.
14. $x=\pm 2, \pm 5$; $y=\pm 3, \pm 6$.
15. $x=\pm 7, \pm \sqrt{3}$; $y=\pm 2, \mp 3\sqrt{3}$.
16. $x=\pm 3, \pm 36$; $y=\pm 5, \mp \frac{23}{2}$.
17. $x=5, 3$; $y=3, 5$.
18. $x=7, -6$; $y=6, -7$.
19. $x=6, -2$; $y=2, -6$.
20. $x=7, 1, 4\pm\sqrt{28}$; $y=1, 7, 4\mp\sqrt{28}$.
21. $x=4, 3, 6, 2$; $y=\frac{3}{2}, 2, 1, 3$.
22. $x=2, \frac{2}{3}, 4, \frac{1}{3}$; $y=2, 6, 1, 12$.

XXVII. PAGE 212.

1. 13.
2. 45, 9.
3. 7, 8.
4. 3.
5. 15, 12.
6. 9.
7. 7 hours.
8. 7, 5.
9. 90 yards, 160 yards.
10. 55 feet, 30 feet.
11. 36', 60'.
12. 6.
13. 5 shillings.
14. 12.
15. Ninepence.
16. 3 feet.
17. 4 inches.
18. 121 square feet.
19. Fourpence.
20. 40, 12; 30, 16 yards.
21. 56.
22. 50.
23. 25.
24. $6\frac{2}{3}$ miles.
25. 75.
26. 20, 30 miles an hour.
27. 40 and 45 miles an hour.
28. 10 gallons.
29. A, 16; B, 14.
30. Distance, 12 miles; rate, 8 miles an hour.

XXVIII. a. PAGE 216.

1. $(x^2+4x+16)(x^2-4x+16)$.
2. $(9a^2+3ab+b^2)(9a^2-3ab+b^2)$.
3. $(x^2+3xy+y^2)(x^2-3xy+y^2)$.
4. $(m^2+4mn-n^2)(m^2-4mn-n^2)$.
5. $(x^2+2xy-y^2)(x^2-2xy-y^2)$.

- $y = \pm 2$.
 $y = 5, 3$.
 $y = 1, -\frac{4}{3}$.
 $\frac{4}{\sqrt{5}}; y = \pm 1, \pm \frac{3}{\sqrt{5}}$.
 $6; y = \pm 2, \pm 4$.
 $\frac{1}{2}; y = \pm \frac{1}{2}, \pm \frac{3}{2}$.
 $5; y = \pm 3, \pm 6$.
 $y = 6, -7$.
5. 15, 12.
 yards, 160 yards.
 12. 6.
 e. 16. 3 feet.
 19. Fourpence.
 23. 25.
 30 miles an hour.
 gallons.
 ate, 8 miles an hour.
6. $(2x^2 + 9xy - 3y^2)(2x^2 - 9xy - 3y^2)$.
 7. $(2m^2 + 6mn + 3n^2)(2m^2 - 6mn + 3n^2)$.
 8. $(3x^2 + xy + 2y^2)(3x^2 - xy + 2y^2)$. 9. $(x^2 + 3xy - 5y^2)(x^2 - 3xy - 5y^2)$.
 10. $(4a^2 - 6ab + b^2)(4a^2 + 6ab + b^2)$.
 11. $\left(\frac{3}{ab} - 1\right)\left(\frac{9}{a^2b^2} + \frac{3}{ab} + 1\right)$. 12. $\left(6a - \frac{b}{2}\right)\left(36a^2 + 3ab + \frac{b^2}{4}\right)$.
 13. $\left(\frac{x}{5} + y\right)\left(\frac{x^2}{25} - \frac{xy}{5} + y^2\right)$. 14. $\left(\frac{mn}{9} - 1\right)\left(\frac{m^2n^2}{81} + \frac{mn}{9} + 1\right)$.
 15. $\left(\frac{ab}{5} + 10\right)\left(\frac{a^2b^3}{25} - 2ab + 100\right)$. 16. $\left(\frac{x}{8} - \frac{4}{x}\right)\left(\frac{x^3}{64} + \frac{1}{2} + \frac{16}{x^3}\right)$.
 17. $(y - 3x)(x + y)(x - y)$. 18. $(m - 5n)(2n + 3m)(2n - 3m)$.
 19. $(ax + b)(bx + a)$. 20. $(x^2z^2 + y^2)(xy + z)(xy - z)$.
 21. $(a^2 + bx)(a + x)$. 22. $(mn - p)(pm - n)$.
 23. $(3ab - 2x)(2ax - 3b)$. 24. $(2x + 3y)(a^2 + xy)$.
 25. $(2x - 3y)(a^2 + xy)$. 26. $\{ax + (a + 1)\}\{(a - 1)x + a\}$.
 27. $(x - a)(3x - a - 2b)$.
 28. $\{ax + 2(b - c)y\}\{2ax - (3b - 4c)y\}$.
 29. $\{(a - 1)x + a\}\{(a - 2)x + (a - 1)\}$.
 30. $\{(a + 1)x - (b - 1)y\}(ax + by)$.
 31. $(b + c - 1)(b^2 + c^2 + 1 - bc + c + b)$.
 32. $(a + 2c + 1)(a^2 + 4c^2 + 1 - 2ac - a - 2c)$.
 33. $(a + b + 2c)(a^2 + b^2 + 4c^2 - ab - 2bc - 2ca)$.
 34. $(a - 3b + c)(a^2 + 9b^2 + c^2 + 3ab + 3bc - ca)$.
 35. $(a - b - c)(a^2 + b^2 + c^2 + ab - bc + ca)$.
 36. $(2a + 3b + c)(4a^2 + 9b^2 + c^2 - 6ab - 3bc - 2ca)$.
 37. $(x^4 - 9x^2 + 81)(x^2 + 3x + 9)(x^2 - 3x + 9)$.
 38. $(a^4 - 4a^2b^2 - b^4)(a^2 + b^2)(a + b)(a - b)$.
 39. $(a + b + c - d)(a + b - c + d)(c + d + a - b)(c + d - a + b)$.
 40. $\left(x^4 + \frac{1}{16}\right)\left(x^2 + \frac{1}{4}\right)\left(x + \frac{1}{2}\right)\left(x - \frac{1}{2}\right)$.
 41. $(x^3 + y^3)(x^4 + y^4)(x^2 + y^2)(x + y)(x - y)$.
 42. $(x^6 + x^2y^3 + y^6)(x^6 - x^2y^3 + y^6)(x^2 + xy + y^2)(x^2 - xy + y^2)(x + y)(x - y)$.
 43. $\left(\frac{1}{x} + 1\right)\left(\frac{1}{x} - 1\right)(a - 2x)(a^2 + 2ax + 4x^2)$.
 44. $(x^2 + y^2)(x^4 - x^2y^2 + y^4)(x - 2y)(x^2 + 2xy + 4y^2)$.
 45. $(x^2 + 4)(x^4 - 4x^2 + 16)(x + 1)(x^2 - x + 1)$.
 46. $\left(\frac{2}{a} + \frac{3}{b}\right)\left(\frac{2}{a} - \frac{3}{b}\right)(a + b)(a^3 - ab + b^3)$.

47. $\left(\frac{1}{3x} + \frac{y}{2}\right)\left(\frac{1}{3x} - \frac{y}{2}\right)\left(\frac{xy}{2} - 1\right)\left(\frac{x^2y^2}{4} + \frac{xy}{2} + 1\right).$
 48. $(x^2 + 5)(x^2 - 5)\left(x + \frac{1}{2}\right)\left(x - \frac{1}{2}\right).$
 49. $(x + 1)(x^2 - x + 1)(x^2 + 4)(x + 2)(x - 2).$
 50. $(x - 1)(x^2 + x + 1)(4x^2 + 9)(2x + 3)(2x - 3).$

XXVIII. b. PAGE 220.

1. $4x^2 - 49y^2 + 42yz - 9z^2.$ 2. $9x^4 + 26x^2y^2 + 49y^4.$
 3. $25x^4 - 115x^2y^2 + 81y^4.$ 4. $49x^4 - 64x^2y^2 + 48xy^3 - 9y^4.$
 5. $x^5 - y^6.$ 6. $(x + y)^4 + 4(x + y)^3 + 16.$ 7. $16x^2(1 - 4x^2).$
 8. $48a^2(a^4 - 1).$ 9. $x^6 - 64.$ 10. $x^6 - 729a^6.$
 11. $\frac{a^4}{x^2} - 3a^2 - x^2 - \frac{x^4}{a^2}.$ 12. $64x^4(9x^2 - 1).$ 13. $x^8 + a^4x^4 + a^8.$
 14. $1 + x^8 + x^{16}.$ 15. $a^{12} - 3a^8x^4 + 3a^4x^8 - x^{12}.$
 16. $1 - 2x^8 + x^{16}.$ 17. $x^6 - 14x^4 + 49x^2 - 36.$ 18. $x^6 - 14x^4 + 49x^2 - 36.$
 19. $x^6 - 64.$ 20. $a^4 - 18a^2b^2 + 81b^4.$
 21. $a^3 + b^3 + c^3 - 3abc.$ 22. $7x + y + z.$
 23. $(x^4 - 4a^2x^2 + 16a^4)(x^2 - 2ax + 4a^2).$ 24. $5x + 7y - 6z.$
 25. $x + 5.$ 26. $2x(x + 1).$ 27. $5(x - 13).$
 28. $(x + 3)(x^2 + 2x + 4).$ 29. $(7x - 3)(x - 1).$ 30. $a - b.$
 31. $x^3 - 3x^2y - 3xy^2 + y^3.$ 32. $x^4 - 4x^2yz + 7y^2z^2.$
 33. $1 + 9x^2 + 4y^2 + 6xy + 2y - 3x.$ 34. $(x + 1)(x - 3).$
 35. $(2a - 5)(2a - 7).$ 36. $(x - a)(x - b).$
 37. $a^2 + 9x^2 + 4y^2 - 6xy + 2ay + 3ax.$
 38. $9 + 4x^2 + 16y^2 - 8xy + 12y + 6x.$
 45. $\frac{m(m^2 + 3n^2)}{4}.$ 47. $(3a^2 + b^2)(a^2 + 3b^2).$
 49. $\frac{1}{16}(9p^2 - 5q^2)(9q^2 - 5p^2).$ 50. $16ab^3.$

XXIX. a. PAGE 223.

1. $x + c.$ 2. $x^2 - ax + b.$ 3. $x^2 + 2bx - ax - 2ab.$
 4. $x^2 - (p + q)x + 2q(p - q).$ 5. $x^2 - (m + n)x + m(m - n).$
 6. $ax + a + 1.$ 7. $x^2 + bx + a^2.$ 8. $2lx - (3m - 4n)y.$
 9. $(a + 2)x + (a + 1)y.$ 10. $x + b.$
 11. $(x + 1)^6 + 3(x + 1)^4 + 3(x + 1)^2 + 1.$
 12. $(m + 1)b^2x^2 + (n + 1)(m + 1)abx + (n + 1)a^2.$

13. $(m-1)x+m$. 14. $mx-n$. 15. $ap-bq$.
 16. $ax+b$. 17. $2ax-3$. 18. $x+2ab$.
 19. $(x^3-1)(x^2-px+q)(x^2-qx+p)$.
 20. $\{px-(p-1)\}\{(p+1)x+p\}\{(p+2)x+p+1\}$.
 21. $\{(a-3)x+a+1\}\{(a-2)x-a\}\{ax-(a+4)\}$.

XXIX. b. PAGE 227.

1. $x-7$. 2. $2-\frac{1}{m}$. 3. $a+2x$.
 4. $1+x-x^2$. 5. $1+2x-x^2$. 6. $1+x$.
 7. $x-2a$. 8. $a-3x$. 9. $x-y$.
 10. $x^2+(p-1)x-1$. 11. $1+\frac{x}{2}-\frac{x^2}{8}+\frac{x^3}{16}$. 12. $1-x-\frac{x^2}{2}-\frac{x^3}{2}$.
 13. $2+\frac{x}{2}-\frac{x^2}{16}+\frac{x^3}{64}$. 14. $1-\frac{x}{2}-\frac{5x^2}{8}-\frac{5x^3}{16}$.
 15. $a-\frac{x}{2a}-\frac{x^2}{8a^3}-\frac{x^3}{16a^5}$. 16. $x+\frac{a^2}{2x}-\frac{a^4}{8x^3}+\frac{a^6}{16x^5}$.
 17. $a^2-\frac{3x^2}{2a^2}-\frac{9x^4}{8a^6}-\frac{27x^6}{16a^{10}}$. 18. $3a+2x-\frac{2x^2}{3a}+\frac{4x^3}{9a^3}$.
 19. $x-\frac{a^3}{3x^3}-\frac{a^6}{9x^6}$. 20. $2+\frac{x}{12}-\frac{x^2}{288}$.
 21. $\frac{1}{a}+3a^2x-9a^5x^2$. 22. $1-2x+3x^2$.
 23. $3x^2-x-1$. 24. $4-x-\frac{x^2}{16}$.

XXIX. c. PAGE 230.

19. 0. 20. 0. 26. 0.

XXIX. d. PAGE 232.

1. 0. 2. 1. 3. 1. 4. $a+b+c$.
 5. 1. 6. $\frac{1}{abc}$. 7. $\frac{1}{abc}$. 8. 1.
 9. d . 10. $\frac{1}{(x-a)(x-b)(x-c)}$. 11. $\frac{x^2}{(x+a)(x+b)(x+c)}$.
 12. $(a+b+c)^2$. 13. $-\frac{a+b+c}{3}$. 14. $\frac{1}{3}$. 15. $a+b+c$.
 16. $bc+ca+ab$. 17. abc . 18. $(b+c)(c+a)(a+b)$.

XXIX. e. PAGE 238.

- | | | | |
|--|---------------------------------|-------------------------------|----------------------|
| 1. 5. | 2. 10. | 3. $\frac{b}{a}$. | 4. $\frac{p}{16q}$. |
| 5. $\frac{5c}{b}$. | 6. $\frac{d-a^4}{2a^3-c}$. | 7. $a=c, b=\frac{a^2}{4}+2$. | 8. 6. |
| 9. $\pm 3a$. | 10. $\pm \sqrt{\frac{2n}{m}}$. | 11. $b^3=27c^2$. | |
| 12. $c=a(b-a^2)^2, d=(b-a^2)^3$, whence $c^3=a^3d^2$. | 13. 32. | | |
| 15. $(x-1)(x-2)(x-3)$. | 16. $(x+2)(x-3)(x-4)$. | | |
| 17. $(x+2)(x+3)(x+4)$. | 18. $(x-3)(x-5)(x+7)$. | | |
| 19. $(x-2)(x-5)(x+7)$. | 20. $(x+1)(x+2)(x-11)$. | | |
| 21. $(x+1)(3x+2)(2x-1)$. | 22. $(x+2)(3x-1)(2x-3)$. | | |
| 23. $x^6-x^5y+x^4y^2-x^3y^3+x^2y^4-xy^5+y^6$. | | | |
| 24. $x^7-x^6y+x^5y^2-x^4y^3+x^3y^4-x^2y^5+xy^6-y^7$. | | | |
| 25. $x^6+x^4y+x^3y^2+x^2y^3+xy^4+y^5$. | | | |
| 26. $x^9+x^7y+x^6y^2+x^5y^3+x^4y^4+x^3y^5+x^2y^6+xy^7+y^8$. | | | |
| 27. $x^2+(a-2)x+a$. | 28. $(a+1)x^2+ax+a-3$. | 29. 6 or $\frac{2}{3}$. | |
| 30. 13. | 34. 3005. | 35. $-37a^3$. | |

XXX. a. PAGE 244.

- | | | | | |
|----------------------------------|-----------------------------------|-------------------------------------|---|-----------------------------------|
| 1. $\frac{2}{x^{\frac{1}{4}}}$. | 2. $\frac{3}{a^{\frac{1}{3}}}$. | 3. $\frac{4a^3}{x^{\frac{1}{2}}}$. | 4. $3a^2$. | 5. $\frac{a^2}{4}$. |
| 6. $\frac{x^{\frac{1}{3}}}{5}$. | 7. $\frac{3c^4x^2}{5a^3y^3}$. | 8. $\frac{x^ab^a}{y^b}$. | 9. $\frac{6}{x^{\frac{1}{3}}}$. | 10. $\frac{a^{\frac{1}{3}}}{2}$. |
| 11. y^2 . | 12. $\frac{1}{3a^2x^2}$. | 13. $\frac{1}{x^{\frac{3}{2}}}$. | 14. $\frac{x^{\frac{3}{2}}}{4}$. | 15. $2y^{\frac{3}{2}}$. |
| 16. $x^{\frac{5}{4}}$. | 17. $\frac{a}{x^{\frac{1}{3}}}$. | 18. $\frac{1}{a^{\frac{2}{3}}}$. | 19. $\frac{1}{a^2}$. | 20. $\sqrt[5]{x^3}$. |
| 21. $\frac{1}{\sqrt{a}}$. | 22. $\frac{5}{\sqrt{x}}$. | 23. $\frac{2}{\sqrt[3]{a}}$. | 24. $\frac{1}{2\sqrt[3]{a}}$. | 25. $2\sqrt[4]{b^3}$. |
| 26. $\frac{1}{2\sqrt[3]{c}}$. | 27. $\sqrt[3]{x}$. | 28. $\frac{2}{\sqrt[6]{a^5}}$. | 29. $\frac{\sqrt{a}}{2\sqrt[3]{x^2}}$. | 30. $\frac{21}{\sqrt{a^3}}$. |
| 31. $\frac{2}{\sqrt{a}}$. | 32. $\frac{1}{3\sqrt{a^3}}$. | 33. $\frac{4}{\sqrt[3]{x^2}}$. | 34. $\frac{1}{\sqrt[3]{x^{a+2}}}$. | 35. $\sqrt[6]{a^{13}}$. |

36. $\sqrt[5]{a^2}$. 37. $\sqrt[2]{a/x^5}$. 38. $\frac{1}{\sqrt[2]{x}}$. 39. $\frac{1}{\sqrt[2]{a}}$. 40. $\sqrt[6]{a^n}$.
 41. 8. 42. $\frac{1}{32}$. 43. 25. 44. $\frac{1}{4}$. 45. $\frac{1}{216}$.
 46. 625. 47. 9. 48. $\frac{3}{2}$. 49. $\frac{27}{8}$. 50. $\frac{2187}{128}$.

XXX. b. PAGE 247.

1. a^6b^9 . 2. $\frac{x^{\frac{3}{4}}}{y}$. 3. $\frac{1}{y^{2a+35}}$. 4. $\frac{1}{2x^{\frac{1}{2}}y^{\frac{1}{2}}}$. 5. $\frac{4}{9a^2x^2}$.
 6. $16ac^4$. 7. $\frac{x^{\frac{1}{2}}}{y^{\frac{1}{2}}}$. 8. $x^{\frac{1}{6}}$. 9. $\frac{3ax}{2}$. 10. x^{n-1} .
 11. $\frac{1}{x^{n+1}}$. 12. $\frac{1}{x^{\frac{1}{2}}}$. 13. $\frac{1}{a^{\frac{2}{3}}b^{\frac{1}{2}}}$. 14. $a^{\frac{1}{3}}$. 15. x^{b+1} .
 16. $\frac{1}{x^{\frac{1}{2}}}$. 17. $\frac{1}{x^2}$. 18. ab^2 . 19. $a+b$. 20. $\frac{1}{(x^2-y^2)^{3n}}$.
 21. $\frac{1}{a^5}$. 22. $b^{\frac{3}{2}}$. 23. $x^{\frac{1}{6}}$. 24. $\frac{a+b}{(a-b)^{\frac{1}{2}}}$. 25. $c^{\frac{7}{2}}$.
 26. $\frac{x^2}{a^3}$. 27. $\frac{1}{a^{\frac{8}{3}}}$. 28. $ab(b^6-a^6)^{\frac{1}{3}}$. 29. $a^{n(n-1)}+a$.
 30. $x^{n(n-1)}+x^{n-1}$. 31. $a^{4n(p-q)}$. 32. x^b . 33. $\frac{x^7}{y^7}$.
 34. $x^{\frac{1}{2}}y^{\frac{5}{6}}$. 35. 2^{n^2} . 36. $\frac{1}{4}$. 37. 4. 38. 1.

XXX. c. PAGE 250.

1. $12x^{\frac{2}{3}}-20x^{\frac{1}{3}}+41-15x^{-\frac{1}{3}}+24x^{-\frac{2}{3}}$.
 2. $9a^{\frac{1}{2}}-9a^{\frac{3}{2}}-25+23a^{-\frac{2}{3}}+6a^{-\frac{1}{3}}$.
 3. $2c^{2x}-9c^x-34+31c^{-x}-6c^{-2x}$. 4. $8x^{3a}+14x^a-3x^{-a}-9x^{-3a}$.
 5. $7x^{\frac{2}{3}}-2x^{\frac{1}{3}}+1$. 6. $3a^{\frac{1}{2}}-3a^{-\frac{1}{2}}+2a^{-1}$.
 7. $8a^{-2}+7a^{-1}+6$. 8. $5b^{\frac{1}{2}}+4b^{\frac{1}{3}}+3b^{-\frac{1}{2}}+2b^{-\frac{1}{3}}$.
 9. $7a^{2x}+3a^x-4$. 10. $c^{2n}-1+c^{-2n}$.
 11. $3x^{\frac{1}{2}}-2+x^{-\frac{1}{2}}$. 12. $5a^{\frac{2}{3}}-3a^{\frac{1}{3}}+4$.

13. $2x^{\frac{2}{3}} - 4 + 3x^{-\frac{2}{3}}$. 14. $a^{2x} - 3a^x - 2$.
 15. $a^3 + 2a - 16a^{-2} - 32a^{-3}$. 16. $1 - x^{\frac{1}{3}} - 2x^{\frac{1}{3}} + 2x^{\frac{2}{3}}$.
 17. $4a^{\frac{2}{3}} - 8a^{\frac{1}{3}} - 5 + 10a^{-\frac{1}{3}} + 3a^{-\frac{2}{3}}$. 18. $x^{\frac{1}{2}} - 2x^{\frac{1}{4}} + 4x^{-\frac{1}{4}} - 8x^{-\frac{1}{2}}$.
 19. $1 - 2a - 2a^{\frac{2}{3}}$. 20. $2x^{\frac{1}{4}} - 3x^{-\frac{1}{4}} - x^{-\frac{3}{4}}$.
 21. $3x^{-2} - 3x^{-1}y^{\frac{1}{2}} + y$. 22. $2x^{\frac{2}{3}} - 3y^{\frac{1}{2}} + 4x^{-\frac{2}{3}}y^{\frac{1}{2}}$.
 23. $9x^{\frac{2}{3}}y^{-1} + 2x^{\frac{1}{3}}y^{-\frac{1}{2}} - 9$. 24. $\frac{1}{4}x^{-1} + 1 - 3y^{\frac{1}{2}}$.

XXX. d. PAGE 252.

1. $x - 4x^{\frac{1}{2}} - 21$. 2. $16x^2 - 8 - 15x^{-2}$. 3. $49x^2 - 81y^{-2}$.
 4. $x^my^{-n} - x^{-m}y^n$. 5. $a^{2x} - 4 + 4a^{-2x}$. 6. $a^{2x} + 2a^{x+\frac{1}{x}} + a^{\frac{2}{x}}$.
 7. $x^a - x^{-\frac{a}{2}} + \frac{1}{4}x^{-2a}$. 8. $20x^{2a}y^{2b} + 13 - 15x^{-2a}y^{-2b}$.
 9. $\frac{1}{9}a^{\frac{2}{3}} - \frac{2}{3} + a^{-\frac{2}{3}}$. 10. $9x^{2a} + 15y^{2b} - 15y^{-2b} - 25x^{-2a}$.
 11. $a^{2x} - (x^2 - \frac{7}{4} + a^{-x} + a^{-2x})$. 12. $x^{\frac{2}{a}} + x^{-\frac{2}{a}} + x^2 - 2 + 2x^{1+\frac{1}{a}} - 2x^{1-\frac{1}{a}}$.
 13. $2a + 2(a^2 - b^2)^{\frac{1}{2}}$. 14. $a + b + (a - b)^{-1} - 2(a + b)^{\frac{1}{2}}(a - b)^{-\frac{1}{2}}$.
 15. $x^{\frac{1}{2}} - 3a^{\frac{1}{2}}$. 16. $x + 3x^{\frac{1}{2}} + 9$. 17. $a^x + 4$.
 18. $x^{2a} - 2x^a + 4$. 19. $c^x + c^{-\frac{x}{2}}$. 20. $1 + 2a^{-1} + 4a^{-2}$.
 21. $a^{2x} - x^3$. 22. $x^{-3} - x^{-2} + x^{-1} - 1$.
 23. $x^{\frac{4}{3}} + x + x^{\frac{2}{3}} + x^{\frac{1}{3}} + 1$. 24. $x^{4n} - 2x^{2n} + 4x^{2n} - 8x^n + 16$.
 25. $x^2 + 2x^{\frac{3}{2}} + x - 16$. 26. $4x^{\frac{2}{3}} + 16x^{\frac{1}{3}} + 16 - 9x^{-\frac{2}{3}}$.
 27. $4 - x^{\frac{2}{3}} + 4x + x^2$. 28. $a^{2x} - 49 - 42a^{-x} - 9a^{-2x}$. 29. $a^{\frac{1}{2}}(a^{\frac{1}{2}} - 2b^{\frac{1}{2}})$.
 30. 1. 31. $\frac{x^{\frac{3}{2}} - 2}{x^{\frac{3}{2}} + 2}$. 32. $\frac{a^{\frac{1}{2}}}{b}$.

XXXI. a. PAGE 255.

1. $\sqrt[12]{x^4}$. 2. $\frac{1}{\sqrt[12]{a^6}}$. 3. $\sqrt[12]{\frac{x}{a}}$. 4. $\sqrt[12]{a^9}$.
 5. $\sqrt[12]{a^{21}}$. 6. $\sqrt[12]{a^4}$. 7. $\sqrt[12]{x^{\frac{21}{8}}}$. 8. $\sqrt[n]{x^{an}}$.

$$-2.$$

$$2x^{\frac{1}{2}} + 2x^{\frac{3}{2}}.$$

$$+4x^{-\frac{1}{2}} - 8x^{-\frac{3}{2}}.$$

$$1^{\frac{1}{2}} - x^{-\frac{5}{2}}.$$

$$1 + 4x^{-\frac{3}{2}}y^{\frac{1}{2}}.$$

$$-3y^{\frac{1}{2}}.$$

$$49x^2 - 81y^{-2}.$$

$$a^{2x} + 2a^{x+\frac{1}{x}} + a^{\frac{2}{x}}.$$

$$y^{-20}.$$

$$25x^{-2a}.$$

$$2 + 2x^{1+\frac{1}{a}} - 2x^{1-\frac{1}{a}}.$$

$$-b)^{\frac{1}{2}}(a-b)^{-\frac{1}{2}}.$$

$$a^x + 4.$$

$$1 + 2a^{-1} + 4a^{-2}.$$

$$+4x^{2n} - 8x^n + 16.$$

$$1 + 16 - 9x^{-\frac{2}{3}}.$$

$$29. a^{\frac{1}{2}}(a^{\frac{1}{2}} - 2b^{\frac{1}{2}}).$$

$$32. \frac{a^{\frac{1}{2}}}{b}.$$

$$9. \sqrt[n]{a^{\frac{3}{2}}}.$$

$$10. \frac{1}{\sqrt[n]{a^{\frac{1}{2}}}}.$$

$$11. \sqrt[n]{x^{\frac{n-1}{2}}y^{\frac{1}{2}}}. \quad 12. \sqrt[n]{a^n}.$$

$$13. \frac{1}{\sqrt[n]{x^{\frac{2}{3}}y^{2n}}}.$$

$$14. \sqrt[n]{a^{\frac{2}{3}}x^{n^2}}.$$

$$15. \sqrt[10]{a^9}, \sqrt[10]{a^{10}}. \quad 16. \sqrt[10]{a^6}, \sqrt[10]{a^5}.$$

$$17. \sqrt[24]{x^9}, \sqrt[24]{x^{18}}, \sqrt[24]{x^9}.$$

$$18. \sqrt[12]{x^3}, \sqrt[12]{x^{10}}. \quad 19. \sqrt[21]{a^3b^4}, \sqrt[21]{a^3b^3}.$$

$$20. \sqrt[26]{a^{13}x^{26}}, \sqrt[26]{a^3x^4}.$$

$$21. \sqrt[6]{125}, \sqrt[6]{121}, \sqrt[6]{13}.$$

$$22. \sqrt[3]{64}, \sqrt[3]{81}, \sqrt[3]{6}.$$

$$23. \sqrt[3]{2}, \sqrt[3]{2}, \sqrt[3]{2}.$$

XXXI. b. PAGE 256.

$$1. 12\sqrt{2}.$$

$$2. 7\sqrt{3}.$$

$$3. 4\sqrt[3]{4}.$$

$$4. 6\sqrt[3]{2}.$$

$$5. 15\sqrt{6}.$$

$$6. 24\sqrt{5}.$$

$$7. 35\sqrt{5}.$$

$$8. 7\sqrt[3]{3}.$$

$$9. 5\sqrt[4]{5}.$$

$$10. -9\sqrt[3]{3}.$$

$$11. 6a\sqrt{a}.$$

$$12. 3ab^2\sqrt{3ab}.$$

$$13. -3xy\sqrt[3]{4x}. \quad 14. x^3y^2\sqrt[3]{y^5}.$$

$$15. xy^2\sqrt[3]{x^a}.$$

$$16. (a+b)\sqrt{a}.$$

$$17. 2(x-y)\sqrt[3]{xy}.$$

$$18. \sqrt[3]{242}.$$

$$19. \sqrt[3]{980}.$$

$$20. \sqrt[3]{864}.$$

$$21. \sqrt[3]{750}.$$

$$22. \sqrt[3]{\frac{14}{11}}.$$

$$23. \sqrt[3]{5b}.$$

$$24. \sqrt{9a^2y}.$$

$$25. \sqrt[3]{\frac{3a}{x}}.$$

$$26. \sqrt[3]{8ax}.$$

$$27. \sqrt[3]{2a}.$$

$$28. \sqrt[3]{a^3b^3}.$$

$$29. \sqrt[3]{ab}.$$

$$30. \sqrt{\frac{x}{y}}.$$

$$31. \sqrt{x^2-y^2}.$$

$$32. \sqrt{\frac{a+x}{a-x}}.$$

$$33. 14\sqrt{5}.$$

$$34. \sqrt{7}.$$

$$35. -12\sqrt{11}.$$

$$36. -15\sqrt{3}.$$

$$37. 7\sqrt[3]{7}.$$

$$38. 11\sqrt[3]{3}.$$

$$39. 0.$$

$$40. 17\sqrt[3]{2}.$$

$$41. 20\sqrt{3} - 13\sqrt{2}.$$

$$42. 3\sqrt[3]{6}.$$

$$43. 6\sqrt{7} - 15\sqrt{6}.$$

$$44. \frac{181\sqrt{3}}{9}.$$

XXXI. c. PAGE 259.

$$1. 14\sqrt{6}.$$

$$2. 12\sqrt{3}.$$

$$3. 10\sqrt{3a}.$$

$$4. 30\sqrt{3}.$$

$$5. 288\sqrt{2}.$$

$$6. \sqrt[3]{x^3-4}.$$

$$7. 3\sqrt{3}.$$

$$8. \frac{5\sqrt{2}}{4}.$$

$$9. -\sqrt{13}.$$

$$10. 14\sqrt[3]{9}.$$

$$11. 240\sqrt[3]{4}.$$

$$12. \sqrt{6}.$$

$$13. ab^3\sqrt{ab}.$$

$$14. \frac{33}{10}.$$

$$15. \frac{1}{10}\sqrt{2}.$$

$$16. \frac{2\sqrt{2}}{a}.$$

$$17. \sqrt[3]{a^3}.$$

$$18. \sqrt[3]{a^3}.$$

$$19. \sqrt[3]{a^3}.$$

$$20. \sqrt[3]{a^3}.$$

17. $\frac{a-b}{x}$. 18. 9·89947. 19. 11·18035. 20. 3·77964.
 21. 19·59592. 22. 26·83284. 23. 58·78776. 24. ·81649.
 25. ·28867. 26. ·04472. 27. ·2566. 28. 1·57485.
 29. ·40316.

XXXI. d. PAGE 260.

1. $6x - 10\sqrt{x}$. 2. $2x - 2\sqrt{ax}$. 3. $a\sqrt{b} + b\sqrt{a}$.
 4. $x + y - \sqrt{x+y}$. 5. $30 + 12\sqrt{6}$. 6. $6\sqrt{21} - 46$.
 7. $6 + \sqrt{10}$. 8. $6a - 6x + 5\sqrt{ax}$. 9. $x - 1 + \sqrt{x^2 - x}$.
 10. $x + a - \sqrt{x^2 - a^2}$. 11. $5a + x - 4\sqrt{a^2 + ax}$.
 12. $1 + 8a - 4\sqrt{a + 4a^2}$. 13. $2a - 2\sqrt{a^2 - x^2}$.
 14. $a + x + 2 - 3\sqrt{a+x}$. 15. $2\sqrt{6}$. 16. $16 + 6\sqrt{10}$.
 17. $4x - 2\sqrt{4x^2 - a^2}$. 18. $2x^2 + 2\sqrt{x^4 - 4y^4}$.
 19. $2m + 2\sqrt{m^2 - n^2}$. 20. $13a^2 + 5b^2 - 12\sqrt{a^4 - b^4}$.
 21. $63 - 18x\sqrt{14 - 4x^2}$. 22. $8x^2 - 2\sqrt{16x^4 - 1}$.

XXXI. e. PAGE 262.

1. 113. 2. -166. 3. 172. 4. -6.
 5. $a - 4b$. 6. $9c^2 - 4x$. 7. x . 8. $2p - q$.
 9. $2x$. 10. $25(x^2 - 3y^2) - 49a^2$. 11. $\frac{11 - 3\sqrt{7}}{2}$.
 12. $\frac{3\sqrt{7} - 2\sqrt{3}}{3}$. 13. $\frac{19 - 6\sqrt{2}}{17}$. 14. $2 + \sqrt{6}$.
 15. $\frac{\sqrt{xy}}{y}$. 16. $\frac{\sqrt{5}}{5}$. 17. $\frac{\sqrt{ax}}{a-x}$. 18. $4 + \sqrt{15}$.
 19. $5 + \sqrt{6}$. 20. $8 - \sqrt{42}$. 21. $\frac{\sqrt{7} - \sqrt{2}}{5}$.
 22. $3\sqrt{2} - 2\sqrt{3}$. 23. $x - \sqrt{x^2 - y^2}$. 24. $\sqrt{x^2 + a^2} - a$.
 25. $\frac{1 - \sqrt{1 - x^4}}{x^2}$. 26. $\frac{7a + b + 8\sqrt{a^2 - b^2}}{3a + 5b}$.
 27. $\frac{18 + x^2 - 6\sqrt{9 + x^2}}{x^2}$. 28. $\sqrt{3}$.
 29. $2 - \sqrt{3} = \cdot 26795$. 30. $11 + 5\sqrt{5} = 22\cdot 18035$.
 31. $\sqrt{5} - \sqrt{3} = \cdot 50402$. 32. $\sqrt{5} + 2 = 4\cdot 23607$.
 33. $\frac{\sqrt{5}}{2} = 1\cdot 11803$. 34. $\frac{3\sqrt{3} - 5}{2} = \cdot 09807$.

20. $3.77964.$

24. $.81649.$

28. $1.57485.$

$a\sqrt{b} + b\sqrt{a}.$

$6\sqrt{21} - 46.$

$x - 1 + \sqrt{x^2 - x}.$

$\frac{2}{x^2} + ax.$

$\frac{2}{x^2}.$

16. $16 + 6\sqrt{10}.$

$4y^4.$

$2\sqrt{a^4 - b^4}.$

$-1.$

4. $-6.$

8. $2p - q.$

1. $\frac{11 - 3\sqrt{7}}{2}.$

4. $2 + \sqrt{6}.$

8. $4 + \sqrt{15}.$

1. $\frac{\sqrt{7} - \sqrt{2}}{5}.$

4. $\sqrt{x^2 + a^2} - a.$

$-\frac{b^2}{x}.$

18035.

07.

07.

XXXI. f. PAGE 266.

1. $\sqrt{5} - \sqrt{2}.$

4. $\sqrt{3} + \sqrt{2}.$

7. $4\sqrt{2} - 3.$

10. $\frac{1}{2}\sqrt{5} + 1.$

13. $\frac{4}{3}(\sqrt{2} + 1).$

16. $\sqrt{2} + 1.$

19. $\frac{1}{\sqrt{2}}(\sqrt{3} + 1).$

22. $\sqrt{2} - 1.$

25. $4 + \sqrt{3}.$

28. $2\sqrt{2} + \sqrt{3}.$

2. $\sqrt{10} + \sqrt{3}.$

5. $3\sqrt{7} + 2\sqrt{3}.$

8. $2\sqrt{5} + 3\sqrt{7}.$

11. $2 - \frac{1}{3}\sqrt{3}.$

14. $\frac{4}{2}(\sqrt{3} - 1).$

17. $\sqrt{5} + 1.$

20. $\sqrt{3} - \sqrt{2}.$

23. $\sqrt{3} + 1.$

26. $\sqrt{5} + \sqrt{3}.$

29. $2\sqrt{2} - \sqrt{7}.$

3. $\sqrt{7} - 1.$

6. $\sqrt{10} - 2\sqrt{2}.$

9. $2\sqrt{11} - \sqrt{3}.$

12. $5\sqrt{\frac{1}{2}} + \sqrt{\frac{7}{2}}.$

15. $\frac{4}{5}(\sqrt{2} + 1).$

18. $\frac{1}{2}(\sqrt{5} + 1).$

21. $\frac{4}{2}(\sqrt{5} + \sqrt{3}).$

24. $\sqrt{5} - 1.$

27. $\sqrt{7} - \sqrt{2}.$

30. $\sqrt{11} + 3\sqrt{2}.$

XXXI. g. PAGE 268.

1. 14.

2. 33.

3. 20.

4. 44.

5. 13.

6. $\frac{6}{5}.$

7. $\frac{17}{6}.$

8. 9.

9. 7.

10. $\frac{56}{5}.$

11. 144.

12. 2.

13. $\frac{121}{25}.$

14. $\frac{25}{16}.$

15. 5.

16. 12.

17. 1.

18. 9.

19. 8.

20. 12.

21. $\frac{1}{51}.$

22. 1.

23. 2.

24. $(b - a)^2.$

25. $\frac{(a - b)^2}{2a - b}.$

26. $0, a - b.$

27. 10.

28. $\frac{5}{2}.$

29. 2.

30. $\pm 1.$

XXXI. h. PAGE 269.

1. 49.

2. 4.

3. 49.

4. $\frac{121}{9}.$

5. 16.

6. 64.

7. $\frac{64}{9}.$

8. $\frac{1}{3}.$

9. $\frac{4}{3}.$

10. 9.

11. 4.

12. 1.

13. 4.

14. 50.

15. 11.

16. 3.

17. 6.

18. 25.

19. $\frac{1}{4}.$

20. $\frac{8}{5}.$

21. 6.

22. 361.

XXXII. a. PAGE 274.

- | | | | |
|----------------|----------------|------------|--------------------|
| 1. 6 : 1. | 2. 1 : 2. | 3. 1 : 5. | 4. 9 : 32. |
| 5. $2x : 3y$. | 6. $3b : 4a$. | 7. 6 : 1. | 8. $\frac{1}{5}$. |
| 9. 4 : 1. | 10. 17 : 7. | 11. 3 : 4. | 12. 5 : 4. |
| 16. 21, 28. | 17. 11. | 18. 27. | |

XXXII. b. PAGE 280.

- | | | | |
|--------------------|-----------------------|---------------|--------------------------|
| 1. bc . | 2. $\frac{6b^3}{a}$. | 3. $5y^2$. | 4. b . |
| 5. $4x$. | 6. $12xy^2$. | 7. x^2 . | 8. ab . |
| 9. $4x^2$. | 10. $6a^2x$. | 11. $9ab^2$. | 19. 8 or $\frac{2}{3}$. |
| 20. $x=17, y=11$. | 21. 2 or 0. | 22. 5 or 0. | |

XXXII. c. PAGE 285.

- | | | | | |
|------------------------|-----------------------------------|--------------------------|---------------|---------------|
| 1. 54. | 2. 27. | 3. 35. | 4. 21. | 5. 10. |
| 6. $\frac{1}{5}$. | 7. 16. | 8. 18. | 12. $3a=5b$. | 13. $5x=7y$. |
| 14. $25x^2=27y^2$. | 15. $a^2=b^2$. | 16. ± 6 . | 17. 20. | |
| 19. $y=6x-3x^2+x^3$. | 20. $346\frac{1}{2}$ square feet. | | | |
| 22. 20 miles per hour. | 23. 9 : 4. | 24. $1\frac{5}{9}$ feet. | 25. 4 feet. | |
| | 26. £1960. | | | |

XXXIII. a. PAGE 289.

- | | | | |
|--------------------------------------|---------------------------------------|------------------------------|------------------------|
| 1. 161, 245. | 2. 59, -37. | 3. 34, $89\frac{1}{2}$. | 4. 16, 9. |
| 5. $574\frac{1}{2}, 93\frac{1}{2}$. | 6. 98, $243\cdot6$. | 7. 43. | 8. -49. |
| 9. $-40\frac{1}{2}$. | 10. 7·2. | 11. 9·7. | 12. $25x$. |
| 13. $a+57d$. | 14. $80a-79b$. | 15. 964, 9780. | 16. 3·2, $25\cdot2$. |
| 17. -387, -18900. | 18. $-9\frac{3}{4}, -99\frac{3}{4}$. | 19. $-41\frac{1}{2}, -361$. | |
| 20. 544, 4864. | 21. 779. | 22. -483. | 23. $980\frac{1}{2}$. |
| 24. $-5569\frac{1}{2}$. | 25. 493. | 26. 140. | 27. p^3 . |
| 28. $a^2(4-a)$. | 29. $\frac{a^2(3-a)}{2}$. | 30. $pq(p-4)$. | 31. 30, 3. |
| 32. 25, -3. | 33. 16, -1. | 34. 24, $2\frac{1}{2}$. | 35. 14, -1. |
| 36. 20, 4. | 37. 7, $2a$. | 38. 20, $-2x$. | |

XXXIII. b. PAGE 293.

4. 9:32.
 8. $\frac{1}{5}$
 12. 5:4.
4. b.
 8. ab.
 19. 8 or $\frac{2}{3}$
 22. 5 or 0.
1. 4, 11, 18, ... 2. 13, 10, 7, ... 3. 3, 1, -1, ...
 4. 1, $-\frac{1}{2}$, -2, ... 5. 4, $5\frac{1}{2}$, 7, ... 6. -11, 4, 19, ...
 7. 43. 8. -95. 9. $-6\frac{1}{2}$.
 10. 68, 65, ... 26. 11. $91\frac{2}{3}$, $90\frac{1}{3}$, ... $70\frac{1}{3}$.
 12. $-6\frac{1}{16}$, $-6\frac{8}{16}$, ... $-2\frac{8}{16}$. 13. 6'4, 5'6, ... -5'6.
 14. $8\frac{1}{3}$, $8\frac{1}{8}$, ... $2\frac{1}{2}$. 15. 14 or 15. 16. 8 or 25.
 17. 9 or 86. 18. 13 or 20. 19. 7 or 8.
 20. 11 or 24. 21. 12, 13, 14. 22. 1, 4, 7.
 23. 7, 11, 15, 19, 23. 24. 2, 5, 8, 11, 14. 25. 131.

XXXIV. a. PAGE 297.

5. 10.
 5b. 13. $5x=7y$.
 17. 20.
 et. 25. 4 feet.
4. 16, 9.
 8. -49.
 12. $25x$.
 16. $3\cdot2$, $25\cdot2$.
 19. $-41\frac{1}{2}$, -361.
 23. $980\frac{1}{2}$.
 27. p^3 .
 31. 30, 3.
 35. 14, -1.
1. 48, 384. 2. $\frac{1}{2}$, $\frac{1}{128}$. 3. 1, $\frac{1}{16}$.
 4. $-\frac{1}{27}$, $-\frac{1}{2187}$. 5. 128, 1. 6. 1, 625.
 7. 512. 8. -4374. 9. $\frac{243}{16}$.
 10. -3^{2n} . 11. x^{2p-1} . 12. $\frac{1}{x^{28}}$.
 13. 162, 54, 18. 14. $\frac{1}{2}$, 2, 8, 32. 15. -28, 14, ... $\frac{7}{8}$.
 16. $\frac{16}{27}$, $\frac{8}{9}$, ... 3. 17. 384, 765. 18. -1458, -1092.
 19. $\frac{1}{8}$, $127\frac{7}{8}$. 20. $\frac{1}{90}$, $12\frac{1}{90}$. 21. $30\frac{3}{8}$, $45\frac{5}{8}$.
 22. $\frac{1}{1458}$, $6\frac{1093}{1458}$. 23. $2\frac{2}{81}$. 24. $1\frac{601}{1458}$.
 25. $\frac{1281}{2560}$. 26. $\frac{1365}{2048}$. 27. $5\frac{5}{81}$.
 28. $\frac{4369}{8192}$. 29. $\frac{1}{2}(3^p-1)$. 30. $\frac{2}{3}(1-2^{2p})$.
 31. $\frac{40(3+\sqrt{3})}{3}$. 32. $\frac{\sqrt{a}(a-1)}{a-1}$. 33. $\frac{585\sqrt{2}-292}{2}$.
 34. $364(\sqrt{6}+\sqrt{2})$.

XXXIV. b. PAGE 300.

1. 27. 2. 24. 3. 1. 4. $\frac{1}{3}$ 5. 1. 6. $\frac{64}{65}$
 7. $\frac{27}{29}$ 8. $\frac{8}{15}$ 9. $\frac{1}{3}$ 10. $\frac{1}{6}$ 11. $\frac{8}{33}$ 12. $\frac{25}{66}$
 13. $\frac{1}{27}$ 14. $\frac{5}{8}, \frac{5}{4}, \frac{5}{2}, \dots$ 15. $\frac{2187}{256}, \frac{729}{128}, \frac{243}{64}, \dots$
 16. $\frac{1}{25}, -\frac{1}{5}, 1, \dots$ 17. $\frac{9}{64}, -\frac{9}{32}, \frac{9}{16}, \dots$ 18. 75, 60, 48.
 20. $\frac{3}{4}, \frac{1}{4}, \frac{1}{12}, \dots$ 21. $\frac{y^2(y^{2n}-1)}{y^2-1} + bn(n+1)$ 22. $\frac{140+99\sqrt{2}}{8}$
 23. $\frac{9(3\sqrt{6}+2\sqrt{2})}{46}$ 24. $2n^2(2n+1) - \frac{3}{8}\left(1 - \frac{1}{3^{2n}}\right)$.

XXXV. PAGE 304.

1. $\frac{2}{3}$ 2. $\frac{5}{14}$ 3. -4. 4. $\frac{3}{n}$ 5. $-\frac{1}{3}, -1, 1, \dots$
 6. 4, 2, $1\frac{1}{3}, \dots$ 7. $-\frac{1}{27}, -\frac{1}{26}, -\frac{1}{25}, \dots$ 8. $2\frac{2}{3}$ 9. $1\frac{6}{7}$
 10. $\frac{1}{7}$ 11. $\frac{2}{a+b}$ 12. $\frac{1}{x}$ 13. $\frac{x^2-y^2}{x}$ 14. $5\frac{1}{7}, 7\frac{1}{5}$
 15. 3, 4, 6. 16. $1\frac{1}{8}, 1\frac{1}{2}, 2, 3$ 19. $36\frac{987}{1624}$ 20. $17\frac{1}{4}$
 21. $\frac{p}{2}\{(p+3)a - (p-3)x\}$ 22. $1\frac{46}{1215}$ 23. -18. 31. $\frac{a^{2n}-b^{2n}}{b^{2n-2}(a^2-b^2)^2}$

Miscellaneous Examples V. PAGE 307.

1. $\frac{c^{1\frac{1}{2}}}{a^{\frac{1}{2}}b^{\frac{1}{4}}}$; 1. 7. $4\sqrt{2}$. 8. 5. 10. 52, 78, 91 yards.
 12. (1) $x^{25} + x^{-25}$. (2) $(a+b)^{\frac{1}{2}}$. 13. $-\frac{32}{7a}$ 14. (1) $\frac{b^4}{a^2}$; (2) $\frac{7}{8}$
 15. $\frac{a+b}{x^2-y^2}$ 21. 2. 23. -5, -2, 1, 4, 7, 10, 13.
 24. 1, 4, 7, ... 25. 1. 26. (1) 275; (2) -1705.
 28. -2, 0, 2, 4, 6. 29. $18\left[1 - \left(\frac{5}{6}\right)^n\right]$. 30. 1. 31. $a+b$
 32. $\frac{3n^2}{2}$. $\frac{63}{2}, \frac{69}{2}, \frac{75}{2}, \frac{81}{2}, \frac{87}{2}$ 34. (1) $9(19a+64x)$. (2) $\frac{2315}{81}$
 37. 10. 38. (1) $s = \frac{1-3^{2n}}{4}$; $l = -3^{2n-1}$. (2) $s = -2n$; $l = 1-4n$.
 39. 1 and 9. 40. $1 + \frac{1}{3} + \frac{1}{9} + \dots$ 41. 8 and 2.

XXXVI. PAGE 317.

1. 6. $\frac{64}{65}$
 12. $\frac{25}{86}$
 37. $\frac{729}{128}$, $\frac{243}{64}$, ...
 60, 48.
 22. $\frac{140+99\sqrt{2}}{8}$
 $\frac{1}{3^{2n}})$
 5. $-\frac{1}{3}$, -1 , 1 , ...
 8. $2\frac{2}{3}$, 9. $1\frac{6}{7}$.
 14. $5\frac{1}{7}$, $7\frac{1}{8}$.
 20. $17\frac{1}{4}$.
 1. $\frac{a^{2n}-b^{2n}}{b^{2n-2}(a^2-b^2)^2}$
 307.
 52, 78, 91 yards.
 (1) $\frac{b^4}{a^2}$, (2) $\frac{7}{8}$
 , 7, 10, 13.
) -1705 .
 1. 31. $a+b$.
 64x). (2) $\frac{2315}{81}$.
 $-2n$; $l=1-4n$.
 3 and 2.
 1. Rational. 2. Rational. 3. Equal, but opposite in sign.
 4. Imaginary. 5. Imaginary. 6. Equal, but opposite in sign.
 7. $x^2-2x-15=0$. 8. $x^2+20x+99=0$.
 9. $x^3-2ax+a^3-b^3=0$. 10. $12x^2-28x+15=0$.
 11. $15x^2+2ax-8a^2=0$. 12. $8x^2-7x=0$. 13. $-\frac{1}{3}$.
 15. Sum $\frac{4}{3}$, difference $\frac{2\sqrt{7}}{3}$, sum of squares $\frac{22}{9}$. 17. $x^2-6x+4=0$.
 18. $x^2+4x+1=0$. 19. $30x^2+(6a-5b)x-ab=0$. 20. $4x^2-16x+9=0$.
 21. $(a^2-b^2)x^2-2(a^2+b^2)x+a^2-b^2=0$. 22. $4abx^2-2(a^2+b^2)x+ab=0$.
 23. $\frac{q^2-2pr}{p^2}$. 24. $\frac{q^2-4pr}{p^2}$. 25. $-\frac{qr}{p^2}$. 26. $\frac{q^4-4prq^2+2p^2r^2}{p^4}$.
 27. $\frac{qr^2(3pr-q^2)}{p^5}$. 28. $\frac{q(3pr-q^2)}{p^2r}$. 29. $P=p(p^2-3q)$, $Q=q^3$.
 30. $b^2x^2-(a^3-3ab)x+b=0$. 31. $2b^2=9ac$.
 32. $8x^2-20a^2x-a^6=0$. 35. $2px^2-(p^3+4q)x+2pq=0$.

XXXVII. a. PAGE 323.

1. 120, 5040, 56, 300. 2. (1) 2520. (2) 5040. 3. 8.
 4. 126. 5. 6. 6. 36. 7. 7 or 8. 8. 2100.
 9. 455, 816; ($r=15$). 10. 242880. 11. 1596000. 12. 504000.

XXXVII. b. PAGE 328.

1. (1) 9979200. (2) 151200. (3) 166320. 2. 420, 360. 3. 18.
 4. 1023. 5. m^n . 6. 168168. 7. 34650.
 8. 120, 144. 9. 1296. 10. 180. 11. 11520.
 13. 78. 14. $(n-2)(n-3)(n-2)$.

XXXVIII. a. PAGE 333.

1. $x^4+8x^2+24x^2+32x+16$.
 2. $x^5+15x^4+90x^3+270x^2+405x+243$.
 3. $a^7+7a^6x+21a^5x^2+35a^4x^3+35a^3x^4+21a^2x^5+7ax^6+x^7$.
 4. $a^5-5a^4x+10a^3x^2-10a^2x^3+5ax^4-x^5$.
 5. $1-10y+40y^2-80y^3+80y^4-32y^5$.
 6. $16x^4+16x^3y+6x^2y^2+xy^3+\frac{y^4}{16}$.

7. $64 - 96x + 60x^2 - 20x^3 + \frac{15x^4}{4} - \frac{3x^5}{8} + \frac{x^6}{64}$.
8. $a^7 - \frac{21a^6}{b} + \frac{189a^5}{b^2} - \frac{945a^4}{b^3} + \frac{2835a^3}{b^4} - \frac{5103a^2}{b^5} + \frac{5103a}{b^6} - \frac{2187}{b^7}$.
9. $a^3x^2 + 9a^7x^2y + 36a^5x^7y^2 + 84a^3x^5y^3 + 126a^5y^4$
 $+ \frac{126x^4y^5}{a} + \frac{84x^3y^6}{a^2} + \frac{36x^2y^7}{a^3} + \frac{9xy^8}{a^4} + \frac{y^9}{a^5}$
10. $220x^3$. 11. $-448y^5$. 12. $21875a^3b^4$.
13. $5440x^3$. 14. $\frac{210}{x^6}$. 15. $\frac{5103x^4a^5}{16}$.
16. -20 . 17. $\frac{2300b^{22}}{x^{16}}$. 18. $-24310x^{25}$.
19. $2x^4 + 36x^2 + 18$. 20. $32 - 40x^2 + 10x^4$. 21. 11520 .
22. $-\frac{1001}{256}a^9$. 23. 7920 . 24. $\frac{1025024}{81}$.

XXXVIII. b. PAGE 338.

1. The 8th. 2. The 9th. 3. The 2nd and 3rd.
4. The 7th. 5. The 11th. 6. The 6th and 7th.
7. $r=7$; excluding the value $r=4$, which makes the terms the same.
8. $n=40$. 9. $3r=3n+2$. 10. $\frac{|2m}{|m|}$. 11. $\frac{|2n}{|n|}x^n$.
12. 65536. 13. 262144.
14. $\frac{n(n-1)\dots(n-r+2)}{|r-1|}a^{n-r+1}(2x)^{r-1}$,
 $\frac{n(n-1)\dots(n-r+2)}{|r-1|}a^{r-1}(2x)^{n-r+1}$.
15. $a^6 + 6a^5 + 15a^4 + 20a^3 + 15a^2 + 6a + 1$
 $x^6 - 12x^5 + 54x^4 - 112x^3 + 108x^2 - 48x + 8$.
16. $1 + \frac{1}{3}x - \frac{1}{9}x^2 + \frac{5}{81}x^3 - \dots$ 17. $1 + \frac{3}{4}x - \frac{3}{32}x^2 + \frac{5}{128}x^3 - \dots$
18. $1 + \frac{2}{5}x - \frac{3}{25}x^2 + \frac{8}{125}x^3 - \dots$ 19. $1 - 6x + 27x^2 - 108x^3 + \dots$
20. $1 + 3x^2 + 6x^4 + 10x^6 + \dots$ 21. $1 - 12x + 90x^2 - 540x^3 + \dots$
22. $\frac{1}{8} - \frac{3}{16}x + \frac{3}{16}x^2 - \frac{5}{32}x^3 + \dots$ 23. $1 - x + \frac{3}{2}x^2 - \frac{5}{2}x^3 + \dots$
24. $\frac{1}{a^{\frac{1}{3}}}\left(1 + \frac{3x}{a} + \frac{15x^2}{2a^2} + \frac{35x^3}{2a^3} + \dots\right)$ 25. $\frac{315}{128}x^4 - \frac{230945}{65536}x^9$.

$$\frac{3a}{b^6} - \frac{2187}{b^7}.$$

$$\frac{2y^7}{b^6} + \frac{9xy^5}{a^7} + \frac{y^9}{a^8}$$

$$21875a^3b^4.$$

$$\frac{5103x^4a^5}{16}.$$

$$-24310x^{20}.$$

$$11520.$$

$$\frac{1025024}{81}.$$

The 2nd and 3rd.
The 6th and 7th.
the terms the same.

$$11. \frac{\frac{1}{2n}x^n}{\frac{1}{n}x^n}.$$

$$2x^2 + \frac{5}{28}x^3 - \dots$$

$$7x^2 - 108x^3 + \dots$$

$$90x^2 - 540x^3 + \dots$$

$$2 - \frac{5}{2}x^3 + \dots$$

$$\frac{230945}{65536}x^9.$$

$$26. \frac{99}{2}x^3, \frac{77}{256}x^{10}.$$

$$27. -4x^3, (-1)^r(r+1)x^r.$$

$$28. -\frac{21}{1024}x^6, -\frac{1 \cdot 3 \cdot 5 \dots (2r-3)}{2^r r}x^r.$$

$$29. \frac{b^r}{a^{r+1}}x^r, -\frac{(n-1)(2n-1) \dots \{(r-1)n-1\}}{r}x^r.$$

$$30. 4.95967.$$

$$31. 4.98998.$$

$$32. 1.98734.$$

$$33. 100504.$$

XXXIX. PAGE 347.

$$1. \frac{15}{2}, -15, -1, \frac{5}{2}.$$

$$2. \frac{9}{2}, -4, -\frac{3}{4}, \frac{3}{2}.$$

$$3. 5, \frac{1}{9}, 125000, 1, -\frac{1}{4}, 2.89, .01.$$

$$4. \frac{1}{2} \log a.$$

$$5. -5 \log y.$$

$$6. 3, 2, 0, -1, -1, -4, 1.$$

$$7. 1.5705780, \bar{5}.5705780, 8.5705780.$$

$$8. 7.623, .000007623, 76230000.$$

$$9. 2.8627278.$$

$$10. 3.9242793.$$

$$11. \bar{1}.4082400.$$

$$12. 7658178.$$

$$13. 8644286.$$

$$14. \bar{1}.4841414.$$

$$16. \log 7 + 4 \log 3 = 2.7535832. \quad 17. 6 \log 2 + \frac{43}{6} \log 3 - \frac{11}{6} = 3.3922160.$$

$$18. \frac{1}{3} \log 2 + \frac{1}{2} \log 3 + \frac{1}{6} \log 7 = .4797536.$$

$$19. \frac{1}{5} (7 \log 2 - 3 \log 3 - \log 7) = \bar{1}.9661496.$$

$$20. \text{Sixty-nine.}$$

$$22. 176.$$

$$23. .398742.$$

$$24. .500977.$$

$$25. 2 - \log 2 - \log 3 - \log 7 = .3767507.$$

$$26. \frac{1 + 2 \log 3 - \log 2}{\log 3} = 3.46.$$

$$27. \frac{5 - 7 \log 2}{3 - 2 \log 2} = 1.206.$$

$$28. \frac{3}{1 - \log 2} = 4.29.$$

$$29. \frac{2 \log 2 - 4 \log 3}{4 \log 2 - \log 3} = -1.8 \text{ very nearly.}$$

XL. PAGE 353.

$$1. 30215.$$

$$2. 25566556.$$

$$3. 244332343.$$

$$4. 3664111.$$

$$5. 3245.$$

$$6. 14320241.$$

$$7. 123807; 1101122.$$

$$8. 3e7580.$$

$$9. 32099.$$

$$10. 30523.$$

$$11. 10000011.$$

$$12. 2^9 + 2^8 + 2^7 + 2^6 + 2^5 + 2^4 + 1; 6e^4 + 9e^3 + e^2 + 4e + 1.$$

$$13. 1736; 145; 328108. \quad 14. 667.$$

$$15. 203.71.$$

10. 31·573.

17. 100133.

18. 50213; 404052.

19. $\frac{7}{9}$

20. Five.

21. Nine.

22. Nine.

23. Seven.

24. 444; 1425; 3333.

XLI. PAGE 359.

2. $\frac{1}{2} \left(x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \right)$

3. $b = a + \frac{a^2}{2} + \frac{a^3}{3} + \frac{a^4}{4} + \dots$

9. 1·6487.

10. $\frac{(-1)^{r-1} 2^r - 1}{r} x^r$

13. $\frac{2^r + 1}{r} x^r$

XLII. a. PAGE 362.

1. 2, -3, $\frac{-1 \pm \sqrt{-27}}{2}$

2. 2, $-\frac{1}{2}$, $\frac{1}{3}(1 \pm \sqrt{10})$

3. $\frac{-1 \pm \sqrt{-3}}{2}$, $\frac{5 \pm \sqrt{21}}{2}$

4. -2, $-\frac{1}{2}$

5. 16, $-\frac{4}{3}$

6. 3, 0.

7. 2, $\frac{15}{4}$

8. 9, $\frac{1}{9}$

9. 3, -2, 1, -6.

10. 1, $\frac{1}{81}$

11. 20, 11.

12. 2, -1.

13. -8, -1, 0.

14. 0.

15. 2, $\frac{3}{2}$, $\frac{7 \pm \sqrt{33}}{4}$

16. 1, $1 \pm 2\sqrt{15}$

17. 7, -1, $3 \pm 2\sqrt{2}$

18. 3, $-\frac{7}{2}$, $\frac{-3 \pm \sqrt{1357}}{12}$

19. $\frac{3 \pm \sqrt{5}}{2}$, $\frac{9 \pm \sqrt{-83}}{6}$

20. 2, $-\frac{1}{2}$, $\frac{3 \pm \sqrt{505}}{4}$

21. $\frac{2}{13}$, $\frac{8}{7}$

22. a^3 , $\frac{1}{a}$

23. 1, $\frac{c-a}{a-b}$

24. 1, $\frac{c(a-b)}{a(b-c)}$

25. a , b .

26. c , $\frac{a^2 + b^2 - ac - bc}{a + b - 2c}$

27. $p + q$

28. 8, -2.

29. $\frac{5}{2}$, $-\frac{13}{2}$

30. α , b .

XLII. b. PAGE 365.

1. $x=5$, $y=2$.

2. $x=4$, 3; $y=3$, 4.

3. $x=7$, -2; $y=2$, -7

4. $x=8$, 2; $y=2$, 8.

5. $x=12$, 3; $y=3$, 12.

6. $x=3$, $y=6$.

60213; 404052.

fine.

44; 1425; 3333.

$$\frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots$$

$$13. \frac{2^r+1}{r} x^r.$$

 $\pm\sqrt{10}$.

$$7. 2, \frac{15}{4}.$$

$$10. 1, \frac{1}{81}.$$

$$13. -8, -1, 0.$$

$$16. 1, 1 \pm 2\sqrt{15}.$$

$$21. \frac{2}{13}, \frac{8}{7}.$$

$$24. 1, \frac{c(a-b)}{a(b-c)}.$$

$$27. p+q.$$

$$30. a, b.$$

$$=3, 4.$$

$$=2, 8.$$

$$=6.$$

$$7. x = \frac{1}{7}, y = \frac{1}{3}. \quad 8. x = 4, 1, \frac{1}{2}(-5 \pm \sqrt{41}); y = 1, \frac{1}{2}(-5 \pm \sqrt{41}).$$

$$9. x = \pm 10, 0; y = \pm 1, \pm \frac{\sqrt{34}}{2}. \quad 10. x = 7, -\frac{35}{4}; y = 3, -\frac{15}{4}.$$

$$11. x = 5, 2, 1 \pm \sqrt{6}; y = -2, -5, -1 \pm \sqrt{6}.$$

$$12. x = \frac{3}{4}, 1, 0; y = \frac{3}{4}, 0, 1.$$

$$13. x = -1, 5 \pm \sqrt{6}; y = -1, 1 \pm \sqrt{\frac{2}{3}}. \quad [\text{It may be shewn that } (x+1)^2 = 27(y+1)^2.]$$

$$14. (1) 3, 2; 2, 3. \quad (2) x = 3, -1; y = 1, -3.$$

$$15. x = 16, 1; y = 1, 16. \quad 16. x = 17, y = \pm 8.$$

$$17. x = \pm 3, y = \pm 2, z = \pm 4. \quad 18. x = 5, y = 6, z = 1, u = 4.$$

$$19. x = \pm 6, y = \pm \frac{8}{3}, z = \pm \frac{3}{2}. \quad 20. x = \pm 5, y = \pm 7, z = \pm 11.$$

$$21. x = -a \pm 2, y = b \pm 1, z = -c \pm 3. \quad 22. x = \pm 5, y = \pm 2, z = \pm 4.$$

$$23. x = \pm 7, y = \pm 3, z = \pm 2. \quad 24. x = \pm 3, y = \mp 4, z = \pm 2.$$

$$25. x = 2, -4; y = 3, -5; z = 0, -2. \quad 26. x = 7, y = 2, z = 4.$$

$$27. x = \pm \frac{1}{2}, 0; y = \pm \frac{1}{3}, 0; z = \pm 1, 0. \quad 28. x = 8, 2; y = 2, 8; z = 4.$$

$$29. x = 2, -6; y = 5; z = 6, -2. \quad 30. x = -5; y = 3, 1; z = 1, 3.$$

$$31. x = 6; y = 9, 4; z = 4, 9. \quad 32. x = \frac{30}{11}, y = \frac{26}{11}.$$

$$33. x = 3, y = 9, z = 6.$$

$$34. x = \pm \frac{a(b^2+c^2)}{2bc}, y = \pm \frac{b(c^2+a^2)}{2ca}, z = \pm \frac{c(a^2+b^2)}{2ab}.$$

XLIII. PAGE 370.

$$1. \text{£}128855000.$$

$$2. 87 \text{ years nearly.}$$

$$3. \text{£}1287. 10s.$$

$$4. 20.63 \text{ per cent. nearly.}$$

$$5. \text{£}2001. 17s. 6d.$$

$$6. 4 \text{ per cent.}$$

$$7. \text{£}3360.$$

$$8. \text{£}11708. 8s.$$

$$9. \text{£}1604.$$

$$10. 20 \text{ years nearly.}$$

Miscellaneous Examples VI. PAGE 371.

1. 0. 2. $-6a - 2b - 4d$. 3. $\frac{1}{2}x^2 + \frac{1}{2}xy - \frac{2}{9}y^2$.
4. 3. 5. $4 - 12x + 13x^2 - 6x^3 + x^4$. 6. $4\frac{1}{3}$.
7. $a - 2$. 8. $\frac{a^2 + b^2}{a^2 - b^2}$. 9. $x = 15, y = 16$.
10. 1, 3. 11. $\frac{4}{9}$. 12. $\frac{13}{12}x^2 - \frac{11}{20}y^2 + \frac{5}{3}z^2$.
13. $x^3 + 24x^2y + 192xy^2 + 512y^3$. 14. $x^2 - y^2$. 15. 11.
16. H.C.F. $(x+2)(x-1)$. L.C.M. $(x-1)(x+2)^2(x^2+2)$.
17. $2a^2 - 3a + 3$. 18. $x = \frac{8b+7a}{9}, y = \frac{8a+7b}{9}$. 19. $\frac{x}{a-x}$.
20. $\frac{5}{3}$. 22. $x - \frac{3}{4}y$. 23. $-35x + 18y + 17z$.
24. (1) $(10x-1)(x+8)$.
(2) $(3x-y)(3x+y)(9x^2+3xy+y^2)(9x^2-3xy+y^2)$.
25. 13. 26. 2. 27. $x^2 - 1$.
28. $\frac{x-3y}{x+3y}$. 29. $x = \frac{2}{a}, y = 3b$.
30. $21\frac{9}{11}$ and $54\frac{6}{11}$ past 7. 31. 1.
32. $\frac{1}{8}x^6 - 8y^6$. 33. $2a^4 + 12a^2 + 2$. 34. 884.
35. $\frac{1}{x^2 - y^2}$. 36. $x = 14, y = 17$. 37. $2x^3 - 3x + 7$.
38. $x = \frac{2}{3}$. 39. $x(x^3 + y^3)(3x - y)$.
40. 21 crowns, 40 half-guineas. 41. $8ab$.
42. $2ab^3 + 3b^4$. 43. 36. 44. $4x - 5$.
45. $\frac{a^2 + b^2}{ab(a-b)^2}$. 46. $2x - \frac{y}{6}$.
47. $x = \frac{1}{2}, y = \frac{1}{3}$; or $x = 0, y = 0$. 48. $\frac{2(x-7)(2x-7)}{(x-2)(x-3)(x-4)(x-5)}$.
49. $(2x+3)(4x+5)(3x-5)(x+2)(x-2)$. 50. 3s. 9d.
51. $-5805x + 5589$. 52. $4a^2 - 9b^2 + 24bc - 16c^2$.
54. $6(x+1)(x-3)(x-4)$. 55. $2(a^2 + b^2)(x^2 + y^2)$.
56. $x = 3, y = 2, z = 1$. 57. 0. 58. $x = -5$.
59. $\frac{x^4 + 2y^4}{y^2(x-y)^2(x^2 + xy + y^2)}$. 60. 24 days. 61. $\frac{3}{2}x^3 - 5x^2 + \frac{x}{4} + 9$.

E 371.

$$\frac{1}{2}x^2 + \frac{1}{2}xy - \frac{2}{9}y^2.$$

$$4\frac{1}{3}.$$

$$x=15, y=16.$$

$$\frac{13}{12}x^2 - \frac{11}{20}y^2 + \frac{5}{3}z^2.$$

$$x^2 - y^2. \quad 15. \quad 11.$$

$$^2 + 2).$$

$$\frac{x}{a-x}.$$

$$-35x + 18y + 17z.$$

$$y^2).$$

$$x^2 - 1.$$

$$1.$$

$$884.$$

$$2x^3 - 3x + 7.$$

$$8ab.$$

$$4x - 5.$$

$$\frac{-7)(2x-7)}{-3)(x-4)(x-5)}.$$

$$3s. \quad 9d.$$

$$2(a^2 + b^2)(x^2 + y^2).$$

$$x = -5.$$

$$\frac{3}{2}x^3 - 5x^2 + \frac{x}{4} + 9.$$

$$62. \quad 94.$$

$$65. \quad x = 2\frac{1}{2}.$$

$$67. \quad x = 24, y = 9, z = 5.$$

$$70. \quad 3.$$

$$73. \quad 6.$$

$$75. \quad x = \frac{qr}{p^2 + q^2}, \quad y = \frac{pr}{p^2 + q^2}.$$

$$77. \quad x = 8.$$

$$80. \quad 640.$$

$$82. \quad \frac{1}{2}.$$

$$84. \quad 2a^2b^2(a-2b)^2(2a+b)^2.$$

$$86. \quad \frac{5x^2 - 4x - 8}{3x^2 + 4x + 24}.$$

$$89. \quad 3(2x-y)(5x+4y). \quad 90. \quad 25 \text{ shillings, } 30 \text{ half-crowns.} \quad 91. \quad 0.$$

$$92. \quad x^3 - x^2 + \frac{5}{3}x - \frac{23}{9}. \quad \text{Rem. } -\frac{163}{9}. \quad 93. \quad 60(p^6 - q^6).$$

$$94. \quad (1) (a - 2b^5)(a^2 + 2ab^5 + 4b^{10}). \quad (2) (x^2 + x - 1)(x^2 - x + 1).$$

$$95. \quad x = a - 2b. \quad 96. \quad (1) \frac{7bc}{13a^2}. \quad (2) \frac{y(y^2 - y + 2)}{2(y+5)}.$$

$$97. \quad x = 1, y = -1, z = 0.$$

$$99. \quad \frac{2a - 3b + c}{2a - 3c}.$$

$$100. \quad \text{Twelve minutes past four.}$$

$$101. \quad (1) 5\frac{1}{8}. \quad (2) -1. \quad 102. \quad (1) \frac{2(a+x)}{a^2 + ax + x^2}. \quad (2) 1 + x - x^2.$$

$$103. \quad a^3 - 3a + \frac{3}{a} - \frac{1}{a^3}; \quad a - \frac{1}{a}.$$

$$104. \quad 1 - 5x + 15x^2 - 45x^3.$$

$$105. \quad £20.$$

$$106. \quad (2a - 3b)(a + b).$$

$$107. \quad (1) 2 \text{ or } \frac{7}{5}. \quad (2) \frac{5}{2} \text{ or } \frac{4}{3}.$$

$$109. \quad 7x^2 - \frac{x}{5} + 3.$$

$$110. \quad \frac{3}{2a^3}.$$

$$111. \quad \frac{3x^2 + 7x - 12}{(x^2 - 9)(x^2 - 16)}; \quad \frac{1}{(x-3)(x-4)}.$$

$$112. \quad \text{H.C.F. } x - 5b. \quad \text{L.C.M. } 6(x+3a)(x-3a)(x-5b).$$

113. (1) $\frac{ab}{2c}$ or $2d$. (2) 9 or -3. 114. 177. 115. 18 miles.
116. $(3x+2y)^2 + (3x+2y)(2x+3y) + (2x+3y)^2 = 19x^2 + 37xy + 19y^2$.
118. (1) $(x+y)(x+y)(x+y)$. (2) $mn(m-n)$.
119. (1) $\begin{matrix} x=3, & 1 \\ y=-1, & -3 \end{matrix}$. (2) $\begin{matrix} x=7, & -5 \\ y=2, & -2 \end{matrix}$. 120. a^3+b^3 .
121. (1) 1. (2) $\frac{x-1}{x^2}$. 122. $a-b$. 123. 435.
124. (1) $x=a \pm b$. (2) $x=3, y=2$.
125. (1) $\frac{x(x+y+z)}{z(x-y+z)}$. (2) $\frac{3x^2+1}{4x(x^2+1)}$. 126. $x^2+(a+2)x+3$.
127. (1) $(x-3y)(x+8y+1)$. (2) $x\left(x+\frac{2}{x}\right)\left(x-\frac{2}{x}\right)$.
128. $x - \frac{3q}{2p} - \frac{9q^2}{8p^3}$. 129. (1) $x=4\frac{1}{2}$. (2) $x = \frac{b}{a^2-ab+b^2}, y = \frac{a}{a^2-ab+b^2}$.
130. $x+a$. 131. (1) x^{a+b+c} . (2) $x^{\frac{5}{12}}y^{\frac{1}{12}}$.
132. 3 shillings. 134. $x-4+\frac{2}{x}$.
135. H.C.F. x^2+a^2 . L.C.M. $(x^2+a^2)(x^2-4a^2)$.
136. (1) $-2y$. (2) 3. 137. $(x-2a)(x^2+2ax+4a^2)(2a+3b)(2a-3b)$.
138. (1) 3. (2) $x=105, y=210, z=420$. 139. The difference is 3.
140. $x^3-4y^3-9z^3-12y^{\frac{2}{3}}z^{\frac{1}{3}}$. 141. 3 hrs. 36 min.
142. (1) $\frac{3}{2}$. (2) $\frac{8}{9}$. 143. $\sqrt{\frac{x}{y}} + \sqrt{\frac{y}{x}} - 1$.
144. (1) $\frac{1}{x^3+1}$. (2) $\frac{a^2+y^2}{a-y}$. 145. (1) $4(\sqrt{2}+\sqrt{3})$. (2) $\sqrt{21}+\sqrt{14}$.
146. (1) $x = \frac{a^2+b^2}{a+b}$. (2) $x=2\frac{1}{2}$ or $-1\frac{3}{4}$, $y=-1\frac{1}{6}$ or $1\frac{2}{3}$.
148. $a^{\frac{1}{2}}-2a^{\frac{1}{4}}x^{\frac{1}{4}}+x^{\frac{1}{4}}$. 149. a^2+1 . 150. Six Shillings.
152. 0. 153. $\frac{x^{11}}{(7x+4)(4x-3)}$.
155. (1) $x=7$ or $-\frac{77}{2}$. (2) $x=\pm 5$ or $\pm 2\sqrt{3}$, $y=\pm 3$ or $\pm \frac{\sqrt{3}}{3}$.
156. (1) 0. (2) $x^{\frac{7}{12}}y^{\frac{5}{6}}$. 157. $(p+1)x-(p-1)$. 158. $\frac{3a}{5b(x^2y^2-1)}$.
159. (1) $1-2^n$. (2) 3^n-2^n . 160. 5 hrs. 57', 47 $\frac{1}{2}$ ".
161. 12. 162. (1) $\left(\frac{ab}{2a+b}\right)^2$. (2) $\frac{5}{5}$.

115. 18 miles.

$$+37xy+19y^2.$$

$$^2+b^2.$$

35.

$$^2+(a+2)x+3.$$

$$\frac{2}{x}.$$

$$y = \frac{a}{a^2-ab+b^2}.$$

$$y^{\frac{1}{2}}.$$

$$(2a+3b)(2a-3b).$$

the difference is 3.

hrs. 36 min.

$$(2) \sqrt{21} + \sqrt{14}.$$

$$\text{or } 1\frac{2}{3}.$$

Six Shillings.

$$= \pm 3 \text{ or } \pm \sqrt{\frac{3}{3}}.$$

$$\frac{3a}{5b(x^2y^3-1)}.$$

$$5 \text{ hrs. } 57', 47\frac{1}{2}''.$$

163. 1.

164. 1, $5+\sqrt{7}$.

165. $(3y+2x)(3y-2x)(x+2)(x^2-2x+4)(x-2)(x^2+2x+4).$

166. (1) 1. (2) $\frac{32}{3}$.

167. (1) 1. (2) $\sqrt{11}-\sqrt{3}$.

168. H.C.F. $5x^2-1$. L.C.M. $=(5x^2-1)^2(4x^2+1)(5x^2+x+1).$

169. (1) $\frac{(a-b)^2}{2b}$. (2) $-2\frac{5}{8}$, $-8\frac{5}{8}$.

170. 9s. & 12s. a dozen.

172. 0.

173. (1) $\frac{x^3+y^3+z^3}{3xyz}$. (2) 1.

174. 11.

175. (1) n. (2) $2-\sqrt{3}$.

176. (1) $x = \frac{69}{20}a$; (2) $x = \pm \frac{1}{2}$, $\pm \frac{3}{2}$; $y = \pm \frac{3}{2}$, $\pm \frac{1}{2}$.

177. $\frac{ax+b^2x^2}{a^m+x^n}$.

178. (1) $c^{\frac{1}{2}}$. (2) 27.

179. 4 miles an hour.

181. $\frac{ac-b^2}{a+c-2b}$.

182. (1) x^2+y^2+xy-1 . (2) $\frac{1}{2x-1}$.

183. $3-2x^2$.

184. (1) $2x$. (2) 10.

185. (1) 5, 1. (2) $x=6, 2, 4$; $y=2, 6, 4$.

186. 0.

187. (1) $20\frac{3}{4}$ or $16\frac{3}{4}$. (2) $x=3$ or $\frac{1}{2}$, $y=-1$ or $\frac{2}{3}$.

189. $(2a+1)x-a$.

190. $2x - \frac{1}{2x}$.

191. $(b+c)(c+a)(a+b)$.

192. (1) $\frac{3x-4}{(x+1)(x^2-1)}$. (2) $\frac{2a}{\sqrt{x+a}}$.

194. Began at $16\frac{4}{11}$ past 3, and ended $27\frac{3}{11}$ past 5; walked 2 hours, $10\frac{10}{11}$ minutes.

195. $\left(\frac{p}{q}\right)^{p+q}$.

197. (1) 47. (2) b.

199. 20.

200. $9x^2+y^2$.

202. 17 years.

203. $q=4$.

The other root is 1.

204. $1\frac{5}{8}$ hours.

205. $a=\frac{5}{3}$, $r=\frac{3}{5}$; or $a=\frac{20}{3}$, $r=-\frac{3}{5}$.

207. $(-1)^m \left(\frac{x+y}{x-y}\right)^{m-n}$.

208. 30240.

209. £1. 15s. 6d.; £377. 10s.

211. Ue0121.

212. $\frac{x}{2} = \frac{y}{3} = \frac{z}{4}$ [zero values are excluded].

214. 9979200; 7560.

215. $32a^5 - 240a^4x + 720a^3x^2 - 1080a^2x^3 + 810ax^4 - 243x^5$. The 3rd and 4th terms.

216. 1.

217. -1.

218. (1) 2, 3, $\frac{5 \pm \sqrt{37}}{2}$; (2) ± 1 , $\pm \sqrt{-3 \pm 2\sqrt{2}}$.

220. 13104000.

222. 2400; 4032000.

223. Number of terms=6; common difference=2.

224. (1) $32 - 60a + 45a^2 - \frac{135}{8}a^3 + \frac{405}{128}a^4 - \frac{243}{1024}a^5$;

(2) $1 - x + \frac{1}{6}x^2 + \frac{1}{54}x^3 + \frac{1}{216}x^4$.

225. Senary. 226. $4\frac{2}{3}$; 1.412. 227. 5.039684. 228. 315.

229. (1) $\sqrt{5x}$; (2) $x - \frac{1}{x}$. 230. $x^2 - 10x + 10 = 0$.

231. 2.71405; *tel.* 232. 1.75; 1.75; -2; $\bar{5}.2375439$.

233. *B* overtakes *A* at the end of the 8th day; then *A* overtakes *B* at the end of the 15th day.

234. (1) $x=21$, $y=6$; (2) $x=4$, $\frac{56}{9}$; $y=3$, $-\frac{11}{3}$; $z=9$, $\frac{121}{9}$.

235. $a=2d$. 236. The 4th and 5th terms. 237. 120; 60.

238. $\bar{3}.698970$; $\cdot 799340$; $\bar{1}.785248$; $x=\frac{22}{17}$. 239. $8x=\sqrt{yz}+2\sqrt[3]{yz}$.

240. $\log 2$. 241. 14. 244. (1) 3; (2) $\bar{1}.36564$; (3) 22.

245. $\frac{5 \cdot 2 \cdot 1 \cdot 4 \cdot 7 \dots (3r-8)}{3^r r} (2a)^{\frac{1}{3}-r} x^{\frac{1}{3}+r}$. 246. 2880.

247. $x^2 - 2(a+b)x + 2ab = 0$. 248. $2(n-1)$ hours.

249. (1) a , $\frac{a(a-b)}{b-c}$, [one root is evidently a , and the product of the roots is $\frac{a^2(a-b)}{b-c}$]; (2) q , $p-q$.

250. The series is the expansion of $\left(1 - \frac{1}{2}\right)^{-\frac{1}{2}}$.

MATHEMATICAL WORKS

BY

Messrs. HALL and KNIGHT,

PUBLISHED BY

Macmillan and Co., Limited.

SEVENTH EDITION, Revised and Enlarged. Now Ready.

ELEMENTARY ALGEBRA FOR SCHOOLS. By H. S.

HALL, M.A., formerly Scholar of Christ's College, Cambridge, Master of the Military Side, Clifton College; and S. R. KNIGHT, B.A., M.B., Ch.B., formerly Scholar of Trinity College, Cambridge, sometime Assistant Master at Marlborough College. Globe 8vo. (bound in maroon-coloured cloth), 3s. 6d. With Answers (bound in green-coloured cloth), 4s. 6d.

The distinctive features of the Seventh Edition are:—

- (1) Greater prominence has been given to *the fundamental Laws of Algebra* (see Arts. 22, 29-32, 46-48). With this object parts of the chapters on Multiplication and Division have been re-written.
- (2) A short section on the use of *Detached Coefficients* has been given on page 37.
- (3) A fuller treatment of the *Remainder Theorem* and its applications will be found on pages 236, 237.
- (4) Five new sets of *Miscellaneous Examples* have been added at convenient intervals, beginning with one on page 32, which replaces Examples IV. c. With the exception of this change and a combination of Examples XI. b. and XI. c, which now appear as one exercise, all the original examples will be found under their old numbers and with the answers unaltered, even where the examples themselves have been improved.

MACMILLAN AND CO., LIMITED, LONDON.

Works by H. S. HALL, M.A., and S. R. KNIGHT, B.A.

It will be found that the pagination is very little altered, the only difference being that the introduction of miscellaneous examples has placed the beginning of some chapters a few pages further on in the book than in previous editions. It is believed that the few alterations in the text will all be justified by usage, and that they will cause no inconvenience to those who are familiar with the book in its old form.

OPINIONS OF THE PRESS.

SCHOOLMASTER—" . . . Has so many points of excellence as compared with its predecessors, that no apology is needed for its issue. The plan always adopted by every good teacher, of frequently recapitulating and making additions at every recapitulation, is well carried out."

NATURE—" . . . We confidently recommend it to mathematical teachers, who, we feel sure, will find it the best book of its kind for teaching purposes."

ACADEMY—"We will not say that this is the best Elementary Algebra for school use that we have come across, but we can say that we do not remember to have seen a better. . . . It is the outcome of a long experience of school teaching, and so is a thoroughly practical book."

EDUCATIONAL TIMES—" . . . A very good book. The explanations are concise and clear, and the examples both numerous and well chosen."

EDUCATIONAL NEWS—"A book of exceptional value."

OPINIONS OF TEACHERS.

"I think it decidedly the best of all books on Elementary Algebra yet published. The great merit seems to me to be that, while it is quite simple and elementary, there are no misleading and inaccurate statements which must afterwards be unlearned. I shall certainly make use of it in my classes, and hope it may come into general use throughout the country."—A. J. WALLIS, M.A., *Fellow and Lecturer of Corpus Christi College, Cambridge.*

"We have examined your Algebra very carefully; and we agree that it is as perfect as a book can be. I will introduce it at St. Paul's as soon as I can."—C. PENDLEBURY, M.A., *Senior Mathematical Master, St. Paul's School.*

"After employing it with my evening class this term, I feel it to be quite the best Elementary Algebra that has yet appeared."—R. A. HERMAN, M.A., *Fellow of Trinity College, Cambridge; Late Professor of Mathematics at University College, Liverpool.*

KEY TO NEW EDITION. Crown 8vo. 8s. 6d.

ANSWERS TO EXAMPLES IN ELEMENTARY ALGEBRA. Fcap. 8vo. Sewed. 1s.

MACMILLAN AND CO., LIMITED, LONDON.

KNIGHT, B.A.

altered, the only
ous examples has
her on in the book
alterations in the
ll cause no incon-
old form.

xcellence as com-
ded for its issue.
y recapitulating and
."

to mathematical
ok of its kind for

mentary Algebra for
do not remember
ong experience of
k."

he explanations are
chosen."

mentary Algebra
ile it is quite simple
ements which must
t in my classes, and
-A. J. WALLIS, M.A.,

e agree that it is
Paul's as soon as I
Paul's School.

eel it to be quite
l."—R. A. HERMAN,
of Mathematics at

z.

TARY ALGE-

ONDON.

Works by H. S. HALL, M.A., and S. R. KNIGHT, B.A.

HIGHER ALGEBRA. A Sequel to Elementary Algebra for
Schools. By H. S. HALL, M.A., and S. R. KNIGHT, B.A.
Fifth Edition, revised and enlarged. Crown 8vo. 7s. 6d.

The Fifth Edition contains a collection of three hundred
Miscellaneous Examples, which will be found useful for advanced
students. These Examples have been selected mainly from recent
Scholarship or Senate House Papers.

SCHOOL GUARDIAN—"We have no hesitation in saying that, in our
opinion, it is one of the best books that have been published on the
subject. . . . The authors have certainly added to their already high reputa-
tion as writers of mathematical text-books by the work now under notice, which
is remarkable for clearness, accuracy, and thoroughness."

"It is a splendid sequel to your *Elementary Algebra*, and I am very pleased to
see you have introduced the essential parts of the Theory of Equations in Chap.
XXXV., which contains all that is required of the subject for ordinary practical
purposes."—A. G. GREENHILL, M.A., *Professor of Mathematics, to the Senior Class*
of Artillery Officers, R.A. Institution, Woolwich.

ATHENÆUM—"The *Elementary Algebra* by the same authors, which has
already reached a third edition, is a work of such exceptional merit that those
acquainted with it will form high expectations of the sequel to it now issued.
Nor will they be disappointed. Of the authors' *Higher Algebra* as of their
Elementary Algebra, we unhesitatingly assert that it is by far the best
work of its kind with which we are acquainted. It supplies a want
much felt by teachers."

ACADEMY—"Is as admirably adapted for College students as its predecessor
was for schools. It is a well-arranged and well-reasoned-out treatise, and con-
tains much that we have not met with before in similar works. For instance, we
note as specially good the articles on Convergency and Divergency of Series, on
the treatment of Series generally, and the treatment of Continued Fractions.
. . . . The book is almost indispensable, and will be found to improve
upon acquaintance."

SATURDAY REVIEW—"They have presented such difficult parts of the sub-
ject as Convergency and Divergency of Series, Series generally, and Probability
with great clearness and fulness of detail. . . . No student preparing for
the University should omit to get this work in addition to any other
he may have, for he need not fear to find here a mere repetition of
the old story. We have found much matter of interest and many valuable
hints. . . . We would specially note the examples, of which there are enough,
and more than enough, to try any student's powers."

KEY. Crown 8vo. 10s. 6d.

MACMILLAN AND CO., LIMITED, LONDON.

Works by H. S. HALL, M.A., and S. R. KNIGHT, B.A.

ALGEBRA FOR BEGINNERS. By H. S. HALL, M.A., and S. R. KNIGHT, B.A., M.B., Ch.B. Globe 8vo. 2s. With Answers. 2s. 6d.

EDUCATIONAL TIMES—"Algebra for Beginners is dealt with on the same lines as the earlier and somewhat more advanced book. The learner is introduced as soon as possible to the practical and more interesting side of the subject, such as equations and problems, while work which largely consists in the manipulation and simplification of elaborate expressions is postponed till later on. The examples for practice are copious, and have been newly composed for this particular book; and, as heretofore, the explanations are clear, concise, and simply expressed. Indeed, without hesitation we pronounce this book the best of its size which we have seen."

UNIVERSITY CORRESPONDENT—"Those masters who have already adopted Messrs. Hall and Knight's *Elementary Algebra* in their schools, will welcome this new work for the use of their junior classes. . . . The numerous exercises for the student are excellent in quality and entirely new. We can unhesitatingly recommend the book to the notice of both teachers and students."

SCHOOLMASTER—"To teachers who have had experience of either the *Elementary* or the *Higher Algebra* it will only be necessary to say that this book is marked by the same qualities which have brought these works into such deserved repute. To those who are still in ignorance of these books, we can say that for clear, simple, and concise explanation, convenient order of subject-matter, and copious and well-graduated exercises, these books have, to say the least, no superiors. Quite early the student is introduced to easy problem work, which can only be looked upon as an advantage. The very numerous exercises are entirely new, so that the book might easily serve as a companion and supplement to the elementary work."

GUARDIAN—"It possesses the systematic arrangement and the lucidity which have gained so much praise for the works previously written by the authors in collaboration."

ALGEBRAICAL EXERCISES AND EXAMINATION PAPERS. With or without Answers. By H. S. HALL, M.A., and S. R. KNIGHT, B.A. Third Edition, revised and enlarged. Globe 8vo. 2s. 6d.

This book has been compiled as a suitable companion to the *Elementary Algebra* by the same authors. It consists of one hundred and twenty progressive Miscellaneous Exercises, followed by a comprehensive collection of papers set at recent examinations.

SATURDAY REVIEW—"To the exercises, one hundred and twenty in number, are added a large selection of examination papers set at the principal examinations which require a knowledge of algebra. These papers are intended chiefly as an aid to teachers, who no doubt will find them useful as a criterion of the amount of proficiency to which they must work up their pupils before they can send them in to the several examinations with any certainty of success."

SCHOOLMASTER—"We can strongly recommend the volume to teachers seeking a well-arranged series of tests in algebra."

MACMILLAN AND CO., LIMITED, LONDON.

KNIGHT, B.A.

**HALL, M.A., and
Svo. 2s. With**

ch on the same lines
er is introduced as
the subject, such as
n the manipulation
ater on. The exami-
d for this particular
and simply expressed,
ne best of its size

ave already adopted
s, will welcome this
nerous exercises for
an unhesitatingly
nd students."

f either the Elemen-
y that this book is
into such deserved
ooks, we can say
venient order of
cises, these books
udent is introduced
vantage. The very
ht easily serve as a

i the lucidity which
a by the authors in

EXAMINATION

**H. S. HALL, M.A.,
ed and enlarged.**

to the *Elementary*
d and twenty pro-
hensive collection

d twenty in number,
ncipal examinations
ended chiefly as an
rion of the amount
efore they can send
ccess."

me to teachers seek-

LONDON.

Works by H. S. HALL, M.A., and S. R. KNIGHT, B.A.

ELEMENTARY TRIGONOMETRY. By H. S. HALL, M.A.,
and S. R. KNIGHT, B.A. Third Edition, containing 300
Additional Miscellaneous Examples. Globe Svo. 4s. 6d.

EDUCATIONAL REVIEW—"The authors have that instinctive knowledge of the needs, both of the pupil and of the teacher, which only belongs to the practical teacher. . . . On the whole it is the best elementary treatise on Trigonometry we have seen."

GUARDIAN—"They are lucid and concise in exposition, their methods are simple, and the examples are judiciously selected."

LYCEUM—"It is not too much to say of this book, that it is the very best class-book that can be placed in the hands of beginners."

SPEAKER—"They here present Elementary Trigonometry so far as it can well be treated without infinite series and imaginary quantities. The authors lay a solid foundation by insisting on the thorough comprehension of trigonometrical ratios before passing on to other subjects. Logarithms and heights and distances have been treated with special care. . . . The full table of contents is a useful feature of the book."

CAMBRIDGE REVIEW—"Messrs Hall and Knight's Algebra has won them a reputation which we think their Trigonometry will sustain."

NATURE—"This book can safely be recommended to beginners, and it may, besides imparting to them a sound elementary knowledge of the subject, ingraft an intelligent interest for more advanced study."

KEY. Crown Svo. 8s. 6d.

ARITHMETICAL EXERCISES AND EXAMINATION PAPERS. With an Appendix containing Questions in LOGARITHMS AND MENSURATION. With or without Answers. By H. S. HALL, M.A., and S. R. KNIGHT, B.A. Third Edition, revised and enlarged. Globe Svo. 2s. 6d.

"In the Second Edition, the Appendix has been increased by a new series of examples, which are intended to be worked by the aid of Logarithmic Tables. In view of the increasing importance of logarithmic calculation in many examinations, this last section will be found especially useful."—*From the Preface.*

CAMBRIDGE REVIEW—"All the mathematical work these gentlemen have given to the public is of genuine worth, and these exercises are no exception to the rule. The addition of the logarithm and mensuration questions adds greatly to the value."

MACMILLAN AND CO., LIMITED, LONDON.

By H. S. HALL, M.A., and F. H. STEVENS, M.A.

A TEXT-BOOK OF EUCLID'S ELEMENTS, including Alternative Proofs, together with Additional Theorems and Exercises, classified and arranged. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military Side, Clifton College. Books I.-VI., XI., and XII., Props. 1 and 3. Globe 8vo. 4s. 6d.

Also in parts separately as follows :—

Book I., . . .	1s.	Books III.-VI., . . .	3s.
Books I. and II., . . .	1s. 6d.	Books IV.-VI., . . .	2s. 6d.
Books I.-III., . . .	2s. 6d.	Books V., VI., XI., and XII.; Props. 1 and 3, . . .	2s. 6d.
Books I.-IV., . . .	3s.	Book XI., . . .	1s.
Books II. and III., . . .	2s.		
Books III. and IV., . . .	2s.		

JOURNAL OF EDUCATION—"The most complete introduction to Plane Geometry based on Euclid's Elements that we have yet seen."

PRACTICAL TEACHER—"One of the most attractive books on Geometry that has yet fallen into our hands."

A KEY TO THE EXERCISES AND EXAMPLES CONTAINED IN A TEXT-BOOK OF EUCLID'S ELEMENTS. Books I.-VI. and XI. By H. S. HALL, M.A., and F. H. STEVENS, M.A., Masters of the Military Side, Clifton College. Crown 8vo. 8s. 6d. Books I.-IV., 6s. 6d. Books VI. and XI., 3s. 6d.

AN ELEMENTARY COURSE OF MATHEMATICS, comprising Arithmetic, Algebra, and Euclid. Globe 8vo. 2s. 6d.

By H. S. HALL and R. J. WOOD.

ALGEBRA FOR ELEMENTARY SCHOOLS. Globe 8vo. Parts I. and II., 6d. each. Cloth, 8d. each. Answers to Parts I. and II., 4d. each. Sewed.

By F. H. STEVENS, M.A.

ELEMENTARY MENSURATION. Globe 8vo. 3s. 6d.

NATURE—"The large number of original examples will be found of great assistance by teachers, and the questions, selected from papers set by the principal examining bodies, will prove of service as tests of the student's capabilities in working out mensuration problems."

MENSURATION FOR BEGINNERS, WITH THE RUDIMENTS OF GEOMETRICAL DRAWING. Globe 8vo. 1s. 6d.

EDUCATIONAL TIMES—"A considerable amount of ground is covered, and the whole is written with rare judgment and clearness."

MACMILLAN AND CO., LIMITED, LONDON.

H. STEVENS, M.A.

LEMENTS, including
ditional Theorems and
H. S. HALL, M.A., and
Military Side, Clifton
Props. 1 and 3. Globe

I.-VI., . . . 3s.
I.-VI., . . . 2s. 6d.
I., VI., XI., and
Props. 1 and 3, 2s. 6d.
" . . . 1s.

te introduction to Plane
have yet seen."
otive books on Geometry

**AND EXAMPLES
K OF EUCLID'S**
By **H. S. HALL, M.A.**,
Military Side, Clifton
I.-IV., 6s. 6d. Books

THEMATICS, com-
Globe 8vo. 2s. 6d.

WOOD.
HOOLS. Globe 8vo.
ch. Answers to Parts

M.A.
bbe 8vo. 3s. 6d.
es will be found of great
from papers set by the
as tests of the student's

**WITH THE RUDI-
WING.** Globe 8vo.

of ground is covered, and
clearness."

D, LONDON.